

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 24 (1978)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON THE GELFAND-FUKS COHOMOLOGY
Autor: Haefliger, André
Kapitel: 7. Example of a computation
DOI: <https://doi.org/10.5169/seals-49696>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Similarly, one can prove that $S_B^*(A \otimes V, B)$ is effectively a model for the space of sections Γ_G (cf. [14]).

Eventually for computations, one proves that one gets also a model for Γ_G by using instead of Ω_{M_G} a DG -algebra A as in § 5 which is a finite dimensional free B -module.

7. EXAMPLE OF A COMPUTATION

Let us consider the case where M is the n -sphere S^n , G the rotation group SO_{n+1} and E the bundle described in § 3. A model for M_G is the DG -algebra A defined by

$$A = R[p_1, \dots, p_k, s] / (s^2 - p_k) \quad d \equiv 0 \quad \text{for } n = 2k$$

$$\text{or } A = R[p_1, \dots, p_{k-1}, \chi] \otimes E(s) \quad ds = \chi \quad \text{for } n = 2k - 1$$

where $\deg p_i = 4i$ and $\deg s = n$.

A model for E_G is obtained by taking the tensor product of A with WU_n , the differential being defined by

$$dh_i = c_i - p_{i/2} \quad \text{and} \quad dc_i = 0.$$

By the way, WSO_n is also a model for E_G .

We now consider the case $n = 2$. The minimal model of E_G is the DG -algebra which begins as

$$A \otimes \Lambda(x_1, x_2, x_3, x_4, x_5, x_{12}, x_{13}, x_{23}, \dots)$$

where

$$\deg x_1 = \deg x_2 = 5, \deg x_3 = 7, \deg x_4 = \deg x_5 = 8,$$

$$\deg x_{12} = 9, \deg x_{13} = \deg x_{23} = 11,$$

etc.

(there is an infinite number of generators).

The differential is defined by

$$dx_1 = dx_2 = 0, dx_3 = -p_1^2, dx_4 = p_1 x_1, dx_5 = p_1 x_2,$$

$$dx_{12} = x_1 x_2, dx_{13} = x_1 x_3 - p_1 x_4, dx_{23} = x_2 x_3 - p_1 x_5,$$

etc.

According to the construction of § 5, a minimal model for the bundle $\Gamma_G \rightarrow B_G$ begins as

$$R[p_1] \otimes \Lambda(\bar{x}_1, \bar{x}_2, x_1, x_2, \bar{x}_3, \bar{x}_4, \bar{x}_5, x_3, \bar{x}_{12}, x_4, x_5, \dots)$$

where

$$\deg \bar{x}_1 = \deg x_i - 2, \varepsilon(x_i) = 1 \otimes x_i + s \otimes \bar{x}_i,$$

dx_i is as above and $dx_{12} = x_1x_2 + p_1\bar{x}_1\bar{x}_2$

$$d\bar{x}_1 = d\bar{x}_2 = d\bar{x}_3 = 0, d\bar{x}_4 = p_1\bar{x}_1, d\bar{x}_5 = p_1\bar{x}_2,$$

$$d\bar{x}_{12} = \bar{x}_1x_2 + x_1\bar{x}_2,$$

etc.

A basis for $H^*(\Gamma_G) = H^*(L_{S^2}, SO_3)$ is given by the classes of the cocycles

$$\bar{x}_1, \bar{x}_2, p_1, x_1, x_2, \bar{x}_3, \bar{x}_1\bar{x}_2, x_1\bar{x}_1, x_1\bar{x}_2, x_2\bar{x}_2, \bar{x}_1\bar{x}_3,$$

$$\bar{x}_2\bar{x}_3, \bar{x}_1\bar{x}_4, \bar{x}_2\bar{x}_5, \bar{x}_1\bar{x}_5 + \bar{x}_2\bar{x}_4, p_1\bar{x}_3,$$

etc.

The first multiplicative relations are

$$p_1\bar{x}_1 \sim 0, p_1\bar{x}_2 \sim 0, \bar{x}_1x_2 \sim \bar{x}_2x_1, p_1^2 \sim 0, \text{ etc.}$$

The first “exotic” class is given by the cocycle $\bar{x}_1\bar{x}_2\bar{x}_{12}$ of degree 13.

The classes \bar{x}_1 and \bar{x}_2 correspond to the classes described by Raoul in his lecture [4], for $n = 2$.

We now give an example of a general statement

THEOREM. *The kernel of the map*

$$H^*(BSO_{n+1}) \rightarrow H^*(L_{S^n}, SO_{n+1})$$

is the ideal generated by the elements which are polynomials of degree $> 2n$ in the Pontrjagin classes $p_1, \dots, p_{[n/2]}$.

As a consequence, we get exactly what is implied by the vanishing theorem of Bott [1]. For instance, for n odd, the image of the powers of the Euler class is non zero. So one can ask for examples of flat $(2k+1)$ -sphere bundles with a non zero power of the Euler class.

One can also check that the homomorphism (see end of § 3)

$$WSO_n \rightarrow C^*(L_{S^n}, SO_{n+1}, \Omega_{S^n})$$

induces an injection in cohomology.