

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 24 (1978)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: REMARKS ON THE UNIVERSAL TEICHMÜLLER SPACE
Autor: Gehring, F. W.
DOI: <https://doi.org/10.5169/seals-49698>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

REMARKS ON THE UNIVERSAL TEICHMÜLLER SPACE¹

by F. W. GEHRING²

1. INTRODUCTION

Suppose that D is a simply connected domain of hyperbolic type in the extended complex plane $\bar{\mathbf{C}} = \mathbf{C} \cup \{\infty\}$. Then the hyperbolic or noneuclidean metric ρ_D in D is given by

$$\rho_D(z) = (1 - |g(z)|^2)^{-1} |g'(z)|,$$

where g is any conformal mapping of D onto the unit disk $\{z: |z| < 1\}$. For each function φ defined in D we introduce the norm

$$\|\varphi\|_D = \sup_{z \in D} |\varphi(z)| \rho_D(z)^{-2}.$$

Next for each function f which is meromorphic and locally univalent in D we let S_f denote the Schwarzian derivative of f . At finite points of D which are not poles of f , S_f is given by

$$S_f = \left(\frac{f''}{f'} \right)' - \frac{1}{2} \left(\frac{f''}{f'} \right)^2,$$

and the definition is extended to ∞ and the poles of f by means of inversion.

Now let L denote the lower half plane $\{z = x + iy: y < 0\}$ and let $B_2 = B_2(L, 1)$ denote the complex Banach space of functions φ analytic in L with the norm

$$\|\varphi\| = \|\varphi\|_L = \sup_{z \in L} 4y^2 |\varphi(z)| < \infty.$$

Next let S denote the family of functions $\varphi = S_g$ where g is conformal in L , and let $T = T(1)$ denote the subfamily of those $\varphi = S_g$ where g has a quasiconformal extension to $\bar{\mathbf{C}}$. From [6] it follows that $\|\varphi\| \leq 6$ for all $\varphi \in S$, and hence that

$$(1) \quad T \subset S \subset B_2.$$

¹) Communicated to an International Symposium on Analysis, held in honour of Professor Albert Pfluger, ETH Zürich, 1978.

²) This research was supported in part by a grant from the U.S. National Science Foundation, Grant MCS-77-02842.

The set T is called the universal Teichmüller space. An important result due to Ahlfors and Bers shows that each Teichmüller space of a Riemann surface R or of a Fuchsian group G has a canonical embedding in the space T . See, for example, [3].

It is natural to ask if there exist relations, other than (1), between S and T as subsets of B_2 . Compactness results for conformal mappings show that S is closed in B_2 . Hence Bers asked in [2] and [3] if one can characterize S in terms of T as follows.

QUESTION. *Is S the closure of T ?*

We shall answer this question in the negative by sketching a proof for the following result.

THEOREM 1. *There exists a φ in S which does not lie in the closure of T .*

On the other hand, we have the following characterization of T in terms of S . See [4].

THEOREM 2. *T is the interior of S .*

2. REFORMULATIONS IN THE PLANE

A set $E \subset \overline{\mathbb{C}}$ is said to be a *quasiconformal circle* if there exists a quasiconformal mapping f defined in $\overline{\mathbb{C}}$ which maps the unit circle $\{z: |z| = 1\}$ onto E .

Theorems 1 and 2 are then respectively equivalent to the following two results on plane domains D .

THEOREM 3. *There exists a simply connected domain D and a positive constant δ such that $f(D)$ is not bounded by a quasiconformal circle whenever f is conformal in D with $\|S_f\|_D \leq \delta$.*

THEOREM 4. *A simply connected domain D is bounded by a quasiconformal circle if and only if there exists a positive constant δ such that f is univalent in D whenever f is meromorphic in D with $\|S_f\|_D \leq \delta$.*

We give an argument to show the equivalence of Theorems 1 and 3. Suppose first that Theorem 1 holds. Then there exists a $\varphi \in S$ and a $\delta > 0$ such that $\|\psi - \varphi\| > \delta$ for all $\psi \in T$. Choose g conformal in L with $S_g = \varphi$, let $D = g(L)$ and suppose that f is conformal in D with $\|S_f\|_D \leq \delta$. Then $h = f \circ g$ is conformal in L ,

$$(2) \quad S_h = (S_{f \circ g})(g')^2 + S_g$$

by the composition law for the Schwarzian derivative, and hence $\psi = S_h \in S$ with

$$\|\psi - \varphi\| = \|S_h - S_g\|_L = \|S_f\|_D \leq \delta.$$

Thus $\psi \notin T$, h does not have a quasiconformal extension to \bar{C} , and $\partial f(D) = \partial h(L)$ is not a quasiconformal circle. Hence Theorem 3 holds.

Suppose next that Theorem 3 holds, let $\varphi = S_g$ where g is any conformal mapping of L onto D , and choose any $\psi \in S$ with $\|\psi - \varphi\| \leq \delta$. Then $\psi = S_h$ where h is conformal in L , $f = h \circ g^{-1}$ is conformal in D and from (2) we obtain

$$\|S_f\|_D = \|S_h - S_g\|_L = \|\psi - \varphi\| \leq \delta.$$

Hence $\partial h(L) = \partial f(D)$ is not a quasiconformal circle, h does not have a quasiconformal extension to \bar{C} and $\psi \notin T$. Thus the distance from φ to T is at least δ and Theorem 1 holds.

A simple modification of the above argument yields the equivalence of Theorems 2 and 4.

Theorems 1 and 3 are immediate consequences of the following result.

THEOREM 5. *There exists a simply connected domain D and a positive constant δ such that $f(D)$ is not a Jordan domain whenever f is conformal in D with $\|S_f\|_D \leq \delta$.*

3. SPIRALS

The proof of Theorem 5 is based on two results for a class of spirals.

DEFINITION. *We say that an open arc α in C is a b -spiral from z_1 onto z_2 if α has the representation*

$$z = (z_1 - z_2)r(t)e^{it} + z_2, \quad 0 < t < \infty,$$

where $r(t)$ is positive and continuous with

$$\lim_{t \rightarrow 0} r(t) = 1, \quad \lim_{t \rightarrow \infty} r(t) = 0,$$

and where $r(t_1) \leq b r(t_2)$ for all t_1, t_2 with $|t_1 - t_2| \leq 2\pi$.

When a is a positive constant, the arc

$$\alpha = \{z = e^{(-a+i)t} : 0 < t < \infty\}$$

is an $e^{2\pi a}$ -spiral from 1 onto 0. Moreover,

$$(3) \quad k(z) |z| = c, \quad \frac{dk}{ds}(z) |z|^2 = d$$

for all $z \in \alpha$, where c and d are positive constants with $d = ac^2$, and where k and s denote the curvature and arclength of α .

The first result we need shows that a curvature condition, similar to (3), is sufficient to guarantee that an open arc is a b -spiral.

LEMMA 1. *Suppose that α is an analytic open arc with 1 and 0 as endpoints, and suppose that*

$$(4) \quad c_1 \leq k(z) |z| \leq c_2, \quad d_1 \leq \frac{dk}{ds}(z) |z|^2 \leq d_2$$

for all $z \in \alpha$, where c_1, c_2, d_1, d_2 are positive constants with $4\pi d_2 < c_1^2$. Then α is a rectifiable b -spiral from 1 onto 0 where

$$b = \frac{c_1 c_2}{c_1^2 - 4\pi d_2}.$$

The second result we require implies that when b is near 1, the points onto which two disjoint b -spirals converge either coincide or are separated by a distance greater than $\frac{1}{2b^2}$ times the diameter of the smaller spiral.

LEMMA 2. *Suppose that α and β are disjoint b -spirals from z_1 onto z_2 and from w_1 onto w_2 , respectively. If $b \in (1, 2)$, then either $z_2 = w_2$ or*

$$|z_2 - w_2| > \frac{1}{b} \min(|z_1 - z_2|, |w_1 - w_2|).$$

4. OUTLINE OF THE PROOF OF THEOREM 5

Fix $a \in \left(0, \frac{1}{8\pi}\right)$ and let $D = \overline{\mathbb{C}} - \gamma$, where

$$\gamma = \{z = \pm i e^{(-a+i)t} : 0 \leq t < \infty\} \cup \{0\}.$$

Then D is a simply connected domain which contains the disjoint $e^{2\pi a}$ -spirals

$$\alpha = \{z = e^{(-a+i)t} : 0 < t < \infty\}, \quad \beta = \{z : -z \in \alpha\}.$$

Next let f denote any conformal mapping of D which fixes the points $1, -1, \infty$. To complete the proof of Theorem 5 it is sufficient to show that there exists a positive constant $\delta = \delta(a)$ such that $f(D)$ is not a Jordan domain whenever $\|S_f\|_D \leq \delta$. This is done in three steps.

First using Lemma 1 and a normal family type argument, we can prove that there exists a $\delta_1 = \delta_1(a) > 0$ with the following property. If $\|S_f\|_D \leq \delta_1$, then $f(\alpha)$ and $f(\beta)$ are b -spirals from 1 onto z_2 and from -1 onto w_2 , respectively, where $b \in (1, 2)$. The points z_2, w_2 are the values which $f(z)$ approaches as $z \rightarrow 0$ from opposite sides of $\partial D = \gamma$.

Next theorems on quasiconformal mappings due to Ahlfors [1] and Teichmüller [8] imply the existence of a positive constant $\delta_2 = \delta_2(a) \leq \delta_1$ such that $|z_2| \leq \frac{1}{5}$ and $|w_2| \leq \frac{1}{5}$ whenever $\|S_f\|_D \leq \delta_2$.

Finally set $\delta = \delta_2$. If $\|S_f\|_D \leq \delta$, then

$$|z_2 - w_2| \leq \frac{2}{5} < \frac{4}{5b} \leq \frac{1}{b} \min(|1 - z_2|, |-1 - w_2|),$$

Lemma 2 implies that $z_2 = w_2$ and hence $f(D)$ is not a Jordan domain. A complete proof for Theorem 5 is given in [5].

5. CONCLUDING REMARKS

We have obtained Theorems 1 and 3 from the stronger conclusion in Theorem 5. We conclude by stating a result for multiply connected domains which implies Theorems 2 and 4.

Given a function φ defined in an arbitrary proper subdomain D of \mathbb{C} , we introduce the norm

$$\|\varphi\|_D^* = \sup_{z \in D} |\varphi(z)| \operatorname{dist}(z, \partial D)^2.$$

When D is simply connected, classical estimates due to Koebe and Schwarz imply that

$$\frac{1}{4} \operatorname{dist}(z, \partial D)^{-1} \leq \rho_D(z) \leq \operatorname{dist}(z, \partial D)^{-1}$$

for $z \in D$, and hence that

$$\|\varphi\|_D^* \leq \|\varphi\|_D \leq 16 \|\varphi\|_D^*.$$

Theorem 6 in [4] and a recent result due to B. Osgood [7] yield the following extension of Theorem 4.

THEOREM 6. *A finitely connected proper subdomain D of \mathbf{C} is bounded by quasiconformal circles or points if and only if there exists a positive constant δ such that f is univalent in D whenever f is meromorphic in D with $\|S_f\|_D^* \leq \delta$.*

REFERENCES

- [1] AHLFORS, L. V. Quasiconformal reflections. *Acta Math.* 109 (1963), pp. 291-301.
- [2] BERS, L. On boundaries of Teichmüller spaces and on kleinian groups I. *Ann. of Math.* 91 (1970), pp. 570-600.
- [3] ——— Uniformization, moduli, and kleinian groups. *Bull. London Math. Soc.* 4 (1972), pp. 257-300.
- [4] GEHRING, F. W. Univalent functions and the Schwarzian derivative. *Comm. Math. Helv.* 52 (1977), pp. 561-572.
- [5] ——— Spirals and the universal Teichmüller space. *Acta Math.* 141 (1978) (to appear).
- [6] KRAUS, W. Über den Zusammenhang einiger Charakteristiken eines einfach zusammenhängenen Bereiches mit der Kreisabbildung. *Mitt. Math. Sem. Giessen* 21 (1932), pp. 1-28.
- [7] OSGOOD, B. Univalence criteria in multiply connected domains. (*To appear*).
- [8] TEICHMÜLLER, O. Extremale quasikonforme Abbildungen und quadratische Differentiale. *Abh. Preuss. Akad. Wiss.* 22 (1940), pp. 1-197.

(Reçu le 15 mai 1978)

F. W. Gehring

Mathematics Department
University of Michigan
Ann Arbor, Michigan, 48104