

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 24 (1978)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: REMARKS ON THE UNIVERS AL TEICHMÜLLER SPACE
Autor: Gehring, F. W.
Kapitel: 1. Introduction
DOI: <https://doi.org/10.5169/seals-49698>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

REMARKS ON THE UNIVERSAL TEICHMÜLLER SPACE¹

by F. W. GEHRING²

1. INTRODUCTION

Suppose that D is a simply connected domain of hyperbolic type in the extended complex plane $\overline{\mathbf{C}} = \mathbf{C} \cup \{\infty\}$. Then the hyperbolic or noneuclidean metric ρ_D in D is given by

$$\rho_D(z) = (1 - |g(z)|^2)^{-1} |g'(z)|,$$

where g is any conformal mapping of D onto the unit disk $\{z : |z| < 1\}$. For each function φ defined in D we introduce the norm

$$\|\varphi\|_D = \sup_{z \in D} |\varphi(z)| \rho_D(z)^{-2}.$$

Next for each function f which is meromorphic and locally univalent in D we let S_f denote the Schwarzian derivative of f . At finite points of D which are not poles of f , S_f is given by

$$S_f = \left(\frac{f''}{f'} \right)' - \frac{1}{2} \left(\frac{f''}{f'} \right)^2,$$

and the definition is extended to ∞ and the poles of f by means of inversion.

Now let L denote the lower half plane $\{z = x + iy : y < 0\}$ and let $B_2 = B_2(L, 1)$ denote the complex Banach space of functions φ analytic in L with the norm

$$\|\varphi\| = \|\varphi\|_L = \sup_{z \in L} 4y^2 |\varphi(z)| < \infty.$$

Next let S denote the family of functions $\varphi = S_g$ where g is conformal in L , and let $T = T(1)$ denote the subfamily of those $\varphi = S_g$ where g has a quasiconformal extension to $\overline{\mathbf{C}}$. From [6] it follows that $\|\varphi\| \leq 6$ for all $\varphi \in S$, and hence that

$$(1) \quad T \subset S \subset B_2.$$

¹) Communicated to an International Symposium on Analysis, held in honour of Professor Albert Pfluger, ETH Zürich, 1978.

²) This research was supported in part by a grant from the U.S. National Science Foundation, Grant MCS-77-02842.

The set T is called the universal Teichmüller space. An important result due to Ahlfors and Bers shows that each Teichmüller space of a Riemann surface R or of a Fuchsian group G has a canonical embedding in the space T . See, for example, [3].

It is natural to ask if there exist relations, other than (1), between S and T as subsets of B_2 . Compactness results for conformal mappings show that S is closed in B_2 . Hence Bers asked in [2] and [3] if one can characterize S in terms of T as follows.

QUESTION. *Is S the closure of T ?*

We shall answer this question in the negative by sketching a proof for the following result.

THEOREM 1. *There exists a φ in S which does not lie in the closure of T .*

On the other hand, we have the following characterization of T in terms of S . See [4].

THEOREM 2. *T is the interior of S .*

2. REFORMULATIONS IN THE PLANE

A set $E \subset \overline{\mathbb{C}}$ is said to be a *quasiconformal circle* if there exists a quasiconformal mapping f defined in $\overline{\mathbb{C}}$ which maps the unit circle $\{z : |z| = 1\}$ onto E .

Theorems 1 and 2 are then respectively equivalent to the following two results on plane domains D .

THEOREM 3. *There exists a simply connected domain D and a positive constant δ such that $f(D)$ is not bounded by a quasiconformal circle whenever f is conformal in D with $\|S_f\|_D \leq \delta$.*

THEOREM 4. *A simply connected domain D is bounded by a quasiconformal circle if and only if there exists a positive constant δ such that f is univalent in D whenever f is meromorphic in D with $\|S_f\|_D \leq \delta$.*