

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 25 (1979)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** INTEGRAL REPRESENTATION THEOREMS VIA BANACH ALGEBRAS  
**Autor:** Maltese, George  
**Kapitel:** 2. Integral representation theorems for linear functionals  
**DOI:** <https://doi.org/10.5169/seals-50383>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.03.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## 2. INTEGRAL REPRESENTATION THEOREMS FOR LINEAR FUNCTIONALS

Let  $A$  be a commutative Banach algebra over  $\mathbf{C}$  and let  $\Delta$  denote the locally compact space of regular maximal ideals of  $A$ . For each  $x \in A$  we use  $\hat{x}$  to denote the Gelfand-transform; i.e.,  $\hat{x}$  is the continuous mapping from  $\Delta$  to  $\mathbf{C}$  defined by the relations:

$$\hat{x}(m) = m(x) \quad \text{for } m \in \Delta.$$

By  $C_0(\Delta)$  we shall denote the algebra of all complex-valued continuous functions on  $\Delta$  which vanish at infinity. For any subset  $\mathcal{A} \subset A$  we shall use the notation  $\hat{\mathcal{A}}$  to denote the set  $\{\hat{x} : x \in \mathcal{A}\}$ . As usual  $\|\hat{x}\|_\infty$  denotes the supremum norm.

**THEOREM 1.** Let  $f$  be a linear form on the complex commutative Banach algebra  $A$  and let  $\mathcal{A}$  be a linear subspace of  $A$ . The following two statements are equivalent:

(1) There exists a constant  $M$  such that

$$|f(x)| \leq M \|\hat{x}\|_\infty \quad \text{for every } x \in \mathcal{A}.$$

(2) There exists a bounded complex Radon measure  $\mu$  on  $\Delta$  such that

$$f(x) = \int_\Delta \hat{x}(m) d\mu(m) \quad \text{for every } x \in \mathcal{A}.$$

*Proof.* The implication (2)  $\Rightarrow$  (1) is clear with  $M = \|\mu\|$ . We shall prove (1)  $\Rightarrow$  (2). Define a mapping  $L : \hat{\mathcal{A}} \rightarrow \mathbf{C}$  by

$$L(\hat{x}) = f(x).$$

It follows from (1) that  $L$  is well-defined, and that

$$|L(\hat{x})| \leq M \|\hat{x}\|_\infty \quad \text{for every } \hat{x} \in \hat{\mathcal{A}}$$

and so  $L$  is continuous with  $\|L\| \leq M$ . Using the Hahn-Banach Theorem we can extend  $L$  to a bounded linear form  $L_0$  on  $C_0(\Delta)$  and by the Riesz Representation Theorem we obtain the existence of a bounded complex Radon measure  $\mu$  on  $\Delta$  such that

$$\| \mu \| = \| L \| = \| L_0 \| \quad \text{and}$$

$$L_0(\varphi) = \int_{\Delta} \varphi(m) d\mu(m) \quad \text{for every } \varphi \in C_0(\Delta).$$

In particular

$$f(x) = L(\hat{x}) = L_0(\hat{x}) = \int_{\Delta} \hat{x}(m) d\mu(m) \quad \text{for every } x \in \mathcal{A}.$$

*Remark:* Suppose that  $A$  has an identity and that  $\hat{A}$  is closed under complex conjugation, then since  $\hat{A}$  contains constants and separates the points of  $\Delta$ , the Stone-Weierstraß Theorem implies that  $\hat{A}$  is dense in  $C(\Delta)$ , the algebra of all complex-valued continuous functions on the compact Hausdorff space  $\Delta$ . If we impose these additional conditions on  $A$  and if we take  $\mathcal{A} = A$  in Theorem 1, we can conclude that in this case the representing measure  $\mu$  is uniquely determined.

If the algebra  $A$  has a continuous involution, one can use Theorem 1 to derive an extended version of a theorem due to Raikov [10]. We proceed to describe the situation.

Let  $A$  be a complex commutative Banach algebra with an isometric involution  $*$  and a bounded approximate identity  $\{u_\lambda\}_{\lambda \in \Lambda}$  i.e., a net satisfying the following conditions:

$$\begin{aligned} \| u_\lambda \| &\leq 1 \quad \text{for each } \lambda \in \Lambda, \\ \| u_\lambda x - x \| &\rightarrow 0 \quad \text{for each } x \in A. \end{aligned}$$

A continuous *positive* functional on  $A$  is an element  $f \in A'$  such that  $f(x^*x) \geq 0$  for every  $x \in A$ . If  $f$  is a continuous positive functional on  $A$  then the Cauchy-Schwarz inequality is valid (Dixmier [8, p. 23]) and this implies the following facts:

$$\begin{aligned} f(u_\lambda) &\rightarrow \| f \| \\ | f(x) |^2 &\leq \| f \| f(x^*x) \quad \text{for every } x \in A. \end{aligned}$$

If the involution is *symmetric*, which means  $(x^*)^\wedge = \overline{x}$  for every  $x \in A$  or, equivalently, that every  $m \in \Delta$  is a *positive* linear functional, then by modifying a classical method of Gelfand-Raikov-Silov [10; p. 62] one can prove that

$$| f(x) | \leq \| f \| \| \hat{x} \|_\infty \quad \text{for every } x \in A.$$

As a corollary to Theorem 1 and the above discussion we obtain the following extended theorem of Raikov [10; p. 64], see also Bucy-Maltese [4]):

**THEOREM 2.** Let  $A$  be a commutative Banach algebra with an isometric involution which is symmetric. Suppose that  $A$  has a bounded approximate identity and let  $f \in A'$  be a continuous positive functional. Then there exists a unique positive Radon measure  $\mu$  on  $\Delta$  such that  $\|\mu\| = \|f\|$  and

$$f(x) = \int_A \hat{x}(m) d\mu(m) \quad \text{for every } x \in A.$$

*Proof.* From the above remarks we know that

$$|f(x)| \leq \|f\| \|\hat{x}\|_\infty \quad \text{for every } x \in A.$$

By Theorem 1 there exists a complex Radon measure  $\mu$  on  $\Delta$  such that  $\|\mu\| \leq \|f\|$  and

$$f(x) = \int_A \hat{x}(m) d\mu(m) \quad \text{for every } x \in A.$$

This formula implies

$$|f(x)| \leq \|\hat{x}\|_\infty \|\mu\| \leq \|x\| \|\mu\| \quad \text{for every } x \in A,$$

so that  $\|f\| \leq \|\mu\|$  and hence  $\|f\| = \|\mu\|$ .

Since  $\hat{A}$  is a self-adjoint subalgebra of  $C_0(\Delta)$  which separates points and for each  $m \in \Delta$  contains a function  $\hat{x}$  such that  $\hat{x}(m) \neq 0$  (in fact <sup>1)</sup> there exists an element  $u_\beta$  of the approximate identity such that  $\hat{u}_\beta(m) \neq 0$ ), the Stone-Weierstraß Theorem implies the uniqueness of the measure  $\mu$ . The positivity of  $\mu$  also follows from the fact that  $\hat{A}$  is dense in  $C_0(\Delta)$ . In fact if  $p$  is a non-negative function in  $C_0(\Delta)$ , then  $p = |q|^2$  for some  $q \in C_0(\Delta)$ . Choose a sequence  $\{x_n\}$  in  $A$  such that

$$\hat{x}_n \rightarrow q.$$

<sup>1)</sup> If  $m \in \Delta$ , then  $\|m\| \neq 0$  and by the assumption of symmetry  $m$  is a positive functional. Therefore, as mentioned above,  $\|m\| = \lim_{\alpha} m(u_\alpha)$  so that there must exist some  $u_\beta$  of the approximate identity such that  $m(u_\beta) \neq 0$ .

Since  $(x_n^*)^\wedge = \overline{x_n}$  it follows that  $(x_n^*)^\wedge \rightarrow \bar{q}$  and hence

$$(x_n x_n^*)^\wedge \rightarrow |q|^2 = p.$$

This implies

$$\begin{aligned} \int_{\Delta} p(m) d\mu(m) &= \lim_n \int_{\Delta} (x_n^* x_n)^\wedge(m) d\mu(m) \\ &= \lim_n f(x_n^* x_n) \geq 0, \end{aligned}$$

so that  $\mu$  is a positive measure and this completes the proof.

If  $A$  has an identity, as is the case in Raikov's original version, the above proof can be somewhat simplified.

**THEOREM 3 (Raikov).** Let  $A$  be a complex commutative Banach algebra with an identity  $e$  and with an isometric involution which is symmetric. If  $f$  is a continuous positive functional on  $A$ , then there exists a unique positive Radon measure  $\mu$  on  $\Delta$  such that  $\|\mu\| = \|f\|$  and

$$f(x) = \int_{\Delta} \hat{x}(m) d\mu(m) \quad \text{for every } x \in A.$$

*Proof.* As above we know that

$$|f(x)| \leq \|f\| \|\hat{x}\|_{\infty} \quad \text{for every } x \in A.$$

From Theorem 1 there exists a complex Radon measure  $\mu$  on  $\Delta$  such that  $\|\mu\| \leq \|f\|$  and

$$f(x) = \int_{\Delta} \hat{x}(m) d\mu(m) \quad \text{for every } x \in A.$$

Hence  $\|\mu\| \leq \|f\| = f(e) = \mu(1) \leq \|\mu\|$  so that  $\mu(1) = \|\mu\|$  which is enough to imply that  $\mu$  is positive. The uniqueness of  $\mu$  follows as in the Remark to Theorem 1.

### 3. APPLICATIONS OF THE INTEGRAL REPRESENTATION THEOREMS

*Application 1 (Bochner's Theorem).* Let  $G$  be a locally compact abelian group and let  $\hat{G}$  denote the (locally compact) character group. Denote