

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 25 (1979)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** INTEGRAL REPRESENTATION THEOREMS VIA BANACH ALGEBRAS  
**Autor:** Maltese, George  
**Kapitel:** 3. Applications of the integral representation theorems  
**DOI:** <https://doi.org/10.5169/seals-50383>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 09.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

Since  $(x_n^*)^\wedge = \overline{x_n}$  it follows that  $(x_n^*)^\wedge \rightarrow \bar{q}$  and hence

$$(x_n x_n^*)^\wedge \rightarrow |q|^2 = p.$$

This implies

$$\begin{aligned} \int_A p(m) d\mu(m) &= \lim_n \int_A (x_n^* x_n)^\wedge(m) d\mu(m) \\ &= \lim_n f(x_n^* x_n) \geq 0, \end{aligned}$$

so that  $\mu$  is a positive measure and this completes the proof.

If  $A$  has an identity, as is the case in Raikov's original version, the above proof can be somewhat simplified.

**THEOREM 3 (Raikov).** Let  $A$  be a complex commutative Banach algebra with an identity  $e$  and with an isometric involution which is symmetric. If  $f$  is a continuous positive functional on  $A$ , then there exists a unique positive Radon measure  $\mu$  on  $\Delta$  such that  $\|\mu\| = \|f\|$  and

$$f(x) = \int_A \hat{x}(m) d\mu(m) \quad \text{for every } x \in A.$$

*Proof.* As above we know that

$$|f(x)| \leq \|f\| \|\hat{x}\|_\infty \quad \text{for every } x \in A.$$

From Theorem 1 there exists a complex Radon measure  $\mu$  on  $\Delta$  such that  $\|\mu\| \leq \|f\|$  and

$$f(x) = \int_A \hat{x}(m) d\mu(m) \quad \text{for every } x \in A.$$

Hence  $\|\mu\| \leq \|f\| = f(e) = \mu(1) \leq \|\mu\|$  so that  $\mu(1) = \|\mu\|$  which is enough to imply that  $\mu$  is positive. The uniqueness of  $\mu$  follows as in the Remark to Theorem 1.

### 3. APPLICATIONS OF THE INTEGRAL REPRESENTATION THEOREMS

*Application 1 (Bochner's Theorem).* Let  $G$  be a locally compact abelian group and let  $\hat{G}$  denote the (locally compact) character group. Denote

Haar measure on  $G$  by  $dt$  and the group algebra (with convolution as multiplication) by  $L^1(G)$ .

*Definition* : A function  $p \in L^\infty(G)$  is said to be *positive definite* provided

$$\int_G \int_G F(t) \overline{F(s)} p(t-s) dt ds \geq 0$$

for every  $F \in L^1(G)$ .

Using the natural involution  $F \rightarrow \tilde{F}$  of  $L^1(G)$  defined by  $\tilde{F}(t) = \overline{F(-t)}$ , we can rewrite the definition of positive definiteness as follows:  $p \in L^\infty(G)$  is positive definite provided

$$\int_G F^* \tilde{F}(t) p(t) dt \geq 0 \quad \text{for all } F \in L^1(G).$$

For a positive definite function  $p$  define the mapping  $P : L^1(G) \rightarrow \mathbb{C}$  as follows:

$$P(F) = \int_G F(t) p(t) dt.$$

Therefore  $P$  is a continuous positive linear functional on the symmetric involution algebra  $L^1(G)$  such that  $\|P\| = \|p\|_\infty$ . By the discussion preceding Theorem 2 and the fact that  $L^1(G)$  has an approximate identity, we know that

$$|P(F)| \leq \|P\| \|\hat{F}\|_\infty \quad \text{for every } F \in L^1(G),$$

where  $\hat{F}$  is the Fourier-Gelfand transform. By Theorem 2 there exists a unique positive Radon measure  $\mu$  on  $\Delta(L^1(G))$  such that

$$P(F) = \int_{\Delta(L^1(G))} \hat{F}(\Gamma) d\mu(\Gamma) \quad \text{for every } F \in L^1(G).$$

It is classical that  $\Delta(L^1(G))$  is homeomorphic to the locally compact character group  $\hat{G}$  under the correspondence  $\Gamma \leftrightarrow \gamma$  where

$$\Gamma(F) = \hat{F}(\gamma) \quad \text{for every } F \in L^1(G).$$

Therefore using the same symbol for the measure induced on  $\hat{G}$  by  $\mu$  under this identification, we obtain

$$P(F) = \int_{\hat{G}} \hat{F}(\gamma) d\mu(\gamma) \quad \text{for every } F \in L^1(G)$$

which implies

$$\begin{aligned}\int_G F(t) p(t) dt &= \int_{\hat{G}} d\mu(\gamma) \int_G F(t) \overline{\gamma(t)} dt \\ &= \int_G F(t) \int_{\hat{G}} \overline{\gamma(t)} d\mu(\gamma) dt.\end{aligned}$$

Since this holds for all  $F \in L^1(G)$  we conclude that

$$p(t) = \int_{\hat{G}} \overline{\gamma(t)} d\mu(\gamma) \quad \text{for almost all } t.$$

This is, of course, the famous Bochner characterization of positive definite functions ([5], [6], [7], [10], [11], [16]).

*Application 2* (A theorem of Schoenberg-Eberlein). A complex-valued function  $\psi$  defined on the locally compact abelian group  $G$  is called a *Fourier-Stieltjes transform* if there exists a bounded Radon measure  $\nu_\psi$  defined on the dual group  $\hat{G}$  such that

$$\psi(t) = \int_{\hat{G}} \overline{\gamma(t)} d\nu_\psi(\gamma) \quad \text{for almost all } t \in G.$$

*Definition*: A measurable complex-valued function  $\psi$  on  $G$  satisfies the *Schoenberg condition* provided

- (a)  $\psi$  is integrable on every compact set;
- (b) there exists a constant  $M$  such that

$$\left| \int_G F(t) \psi(t) dt \right| \leq M \sup_{\gamma} \left| \int_G F(t) \overline{\gamma(t)} dt \right|$$

for every  $F \in K(G)$ , where  $K(G)$  is the set of continuous functions on  $G$  with compact support.

The following theorem is due to Schoenberg [17] for the case  $G = \mathbf{R}$  and to Eberlein [9] in the general case.

**THEOREM.** A measurable complex-valued  $\psi$  has a representation as a Fourier-Stieltjes transform if and only if  $\psi$  satisfies the Schoenberg condition.

*Proof.* It is immediate that if  $\psi$  is a Fourier-Stieltjes transform then  $\psi$  satisfies the Schoenberg condition for the constant  $M = \|\nu_\psi\|$ .

If now  $\psi$  satisfies the Schoenberg condition define  $L : L^1(G) \rightarrow \mathbf{C}$  as usual by

$$L(F) = \int_G F(t) \psi(t) dt.$$

By hypothesis

$$|L(F)| \leq M \|\hat{F}\|_{\infty} \quad \text{for every } F \in K(G).$$

From Theorem 1 there exists a bounded Radon measure  $\nu_{\psi}$  defined on  $\hat{G}$  such that

$$L(F) = \int_{\hat{G}} \hat{F}(\gamma) d\nu_{\psi}(\gamma) \quad \text{for } F \in K(G).$$

Therefore it follows that

$$\int_G F(t) \psi(t) dt = \int_{\hat{G}} d\nu_{\psi}(\gamma) \int_G F(t) \overline{\gamma(t)} dt \quad \text{for all } F \in K(G)$$

and hence

$$\psi(t) = \int_{\hat{G}} \overline{\gamma(t)} d\nu_{\psi}(\gamma) \quad \text{for almost all } t \in G,$$

which completes the proof.

*Application 3* (Positive definite functions on abelian semigroups). Let  $S$  be an abelian semigroup, i.e. a set equipped with a composition law denoted by  $+$  such that the commutative and associative laws are valid. We shall assume the existence of a neutral element  $0$ . By  $l^1(S)$  we shall denote the real commutative Banach algebra of functions  $f : S \rightarrow \mathbb{R}$  with the property that

$$\|f\| = \sum_{a \in S} |f(a)| < \infty.$$

Multiplication in  $l^1(S)$  is defined by convolution; viz.,

$$f * g(a) = \sum_{\substack{s, t \in S \\ s+t=a}} f(s) g(t).$$

By a *character* on  $S$  we shall mean a function  $\gamma : S \rightarrow [-1, 1]$  which satisfies

- (i)  $\gamma(0) = 1$
- (ii)  $\gamma(s+t) = \gamma(s) \gamma(t)$  for all  $s, t \in S$ .

The set  $\hat{S}$  of all characters is an abelian semigroup under pointwise multiplication. If we endow  $\hat{S}$  with the topology of pointwise convergence, then  $\hat{S}$  is a compact Hausdorff space. By  $\Delta(l^1(S))$  we shall denote the

compact subset of  $l^1(S)'$  consisting of all continuous homomorphisms from  $l^1(S)$  onto  $\mathbf{R}$  endowed with the weak \* topology.

If  $\gamma \in \hat{S}$  then the mapping

$$\Gamma_\gamma : l^1(S) \rightarrow \mathbf{R}$$

defined by the relations

$$\Gamma_\gamma(f) = \sum_{s \in S} f(s) \gamma(s) \quad \text{for } f \in l^1(S)$$

is a non-trivial continuous homomorphism from  $l^1(S)$  onto  $\mathbf{R}$ , so that  $\Gamma_\gamma \in \Delta(l^1(S))$ . Conversely for every  $\Gamma \in \Delta(l^1(S))$  there exists  $\gamma \in \hat{S}$  with

$$\Gamma(f) = \sum_{s \in S} f(s) \gamma(s) \quad \text{for } f \in l^1(S).$$

It is easily verified that the mapping

$$\gamma \rightarrow \Gamma_\gamma$$

is a homeomorphism of  $\hat{S}$  onto  $\Delta(l^1(S))$ . We shall identify  $\hat{S}$  with  $\Delta(l^1(S))$  via this homeomorphism and consider the Gelfand transform  $\hat{f}$  of  $f \in l^1(S)$  as a continuous function on  $\hat{S}$  via

$$\hat{f}(\gamma) = \Gamma_\gamma(f).$$

A bounded function  $\varphi : S \rightarrow \mathbf{R}$  is called *positive definite* if

$$\sum_{s \in S} f * f(s) \varphi(s) \geq 0 \quad \text{for all } f \in l^1(S).$$

In the sequel we shall *assume* that the spectral radius formula is valid in the real Banach algebra  $l^1(S)$ ; i.e., we shall assume that

$$\|\hat{f}\|_\infty = \lim_{n \rightarrow \infty} \|f^n\|^{\frac{1}{n}}$$

for every  $f \in l^1(S)$ . One can easily prove that the spectral radius formula is true for the simple functions  $\varepsilon_a$ ,  $\varepsilon_a + \varepsilon_b$ ,  $\lambda \varepsilon_a$  and  $\varepsilon_a * \varepsilon_b$ , where for each  $a \in S$  the function  $\varepsilon_a$  is defined by the relations

$$\varepsilon_a(s) = \begin{cases} 0 & \text{for } s \neq a \\ 1 & \text{for } s = a. \end{cases}$$

In the general context of real Banach algebras, of course, the spectral radius formula need not hold.

The following theorem was demonstrated *without* the assumption of the spectral radius formula for  $l^1(S)$ . The theorem was obtained by Berg-Christensen-Ressel [3] using the Krein-Milman Theorem along with an interesting characterization of the extreme points of the convex compact set of normalized positive definite functions.

**THEOREM.** Assume that the spectral radius formula is valid for each  $f \in l^1(S)$ . If  $\varphi$  is a positive definite function on  $S$ , then there exists a unique positive Radon measure  $\mu$  on  $\hat{S}$  such that

$$\varphi(s) = \int_{\hat{S}} \gamma(s) d\mu(\gamma) \quad \text{for every } s \in S.$$

*Proof.* The functional  $L : l^1(S) \rightarrow \mathbf{R}$  defined by

$$L(f) = \sum_{s \in S} f(s) \varphi(s)$$

is positive, i.e.  $L(f * f) \geq 0$ . By our assumption of the validity of the spectral radius formula we have

$$|L(f)| \leq \varphi(0) \|\hat{f}\|_{\infty} \quad \text{for every } f \in l^1(S).$$

Exactly as in Theorem 1 and Theorem 3 there exists a positive Radon measure  $\mu$  on  $\hat{S}$  such that

$$L(f) = \int_{\hat{S}} f(\gamma) d\mu(\gamma) \quad \text{for every } f \in l^1(S).$$

Hence

$$\begin{aligned} \sum_{s \in S} f(s) \varphi(s) &= \int_{\hat{S}} \left( \sum_{s \in S} f(s) \gamma(s) \right) d\mu(\gamma) \\ &= \sum_{s \in S} f(s) \int_{\hat{S}} \gamma(s) d\mu(\gamma) \quad \text{for every } f \in l^1(S) \end{aligned}$$

and therefore we conclude that

$$\varphi(s) = \int_{\hat{S}} \gamma(s) d\mu(\gamma) \quad \text{for every } s \in S.$$

The uniqueness of  $\mu$  is a consequence of the Stone-Weierstrass Theorem: the algebra  $\widehat{l^1(S)}$  is dense in the algebra of all real continuous functions on  $\hat{S}$ .

*Application 4* (The spectral theorem for normal operators). Let  $\mathcal{H}$  be a Hilbert space and let  $\mathcal{L}(\mathcal{H})$  denote the Banach algebra of all bounded linear operators on  $\mathcal{H}$ . Consider a subalgebra  $A \subset \mathcal{L}(\mathcal{H})$  with the following properties:

- (i)  $A$  is commutative;
- (ii)  $A$  is closed;
- (iii) If  $T \in A$  then  $T^* \in A$ ;
- (iv) The identity operator belongs to  $A$ .

Let  $\Delta$  denote the maximal ideal space of  $A$ . Since each  $T \in A$  is normal it follows that  $\|T\| = \|\hat{T}\|_\infty$  for every  $T \in A$ .

For each pair of vectors  $\xi, \eta \in \mathcal{H}$  define a mapping  $L_{\xi, \eta} : A \rightarrow \mathbb{C}$  by

$$L_{\xi, \eta}(T) = (T\xi, \eta)$$

then we have

$$|L_{\xi, \eta}(T)| \leq \|T\| \cdot \|\xi\| \|\eta\| = \|\xi\| \|\eta\| \cdot \|\hat{T}\|_\infty$$

Therefore by Theorem 1 there exists a bounded complex Radon measure  $\mu_{\xi, \eta}$  on  $\Delta$  such that  $\|\mu_{\xi, \eta}\| \leq \|\xi\| \cdot \|\eta\|$  and

$$L_{\xi, \eta}(T) = \int_\Delta \hat{T} d\mu_{\xi, \eta} \quad \text{for every } T \in A.$$

An application of the Gelfand-Neumark theorem establishes the uniqueness of the measure. The usual construction of a unique resolution of the identity on the Borel sets of  $\Delta$  can be made based on this formula. A specialization of this formula to a single normal operator leads to the classical spectral theorem. We shall not give the details here since many excellent accounts exist (c.f. Berberian [1], [2], Segal-Kunze [18]). An especially lucid presentation is given in Rudin [16].

#### BIBLIOGRAPHY

- [1] BERBERIAN, S. *Notes on spectral theory*. Van Nostrand, Princeton, N.J., 1966.
- [2] ——— *Lectures in functional analysis and operator theory*. Springer Verlag, New York-Heidelberg-Berlin, 1974.