Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 25 (1979)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: INTEGRAL REPRESENTATION THEOREMS VIA BANACH ALGEBRAS

Autor: Maltese, George

Bibliographie

DOI: https://doi.org/10.5169/seals-50383

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 08.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Application 4 (The spectral theorem for normal operators). Let $\mathscr{H}$ be a Hilbert space and let $\mathscr{L}(\mathscr{H})$ denote the Banach algebra of all bounded linear operators on $\mathscr{H}$. Consider a subalgebra $A \subset \mathscr{L}(\mathscr{H})$ with the following properties:

(i) A is commutative;

(ii) A is closed;

(iii) If $T \in A$ then $T^* \in A$;

(iv) The identity operator belongs to A.

Let Δ denote the maximal ideal space of A. Since each $T \in A$ is normal it follows that $\| T \| = \| \hat{T} \|_\infty$ for every $T \in A$.

For each pair of vectors $\xi, \eta \in \mathscr{H}$ define a mapping $L_{\xi,\eta} : A \to \mathbf{C}$ by

$$L_{\xi,\eta}(T) = (T\xi, \eta)$$

then we have

$$| L_{\xi,\eta}(T) | \leqslant \| T \| \cdot \| \xi \| \; \| \eta \| = \| \xi \| \; \| \eta \| \cdot \| \hat{T} \|_\infty$$

Therefore by Theorem 1 there exists a bounded complex Radon measure $\mu_{\xi,\eta}$ on Δ such that $\| \mu_{\xi,\eta} \| \leqslant \| \xi \| \cdot \| \mu \|$ and

$$L_{\xi,\eta}(T) = \int_\Delta \hat{T} \, d\mu_{\xi,\eta} \quad \text{for every} \quad T \in A .$$

An application of the Gelfand-Neumark theorem establishes the uniqueness of the measure. The usual construction of a unique resolution of the identity on the Borel sets of Δ can be made based on this formula. A specialization of this formula to a single normal operator leads to the classical spectral theorem. We shall not give the details here since many excellent accounts exist (c.f. Berberian [1], [2], Segal-Kunze [18]). An especially lucid presentation is given in Rudin [16].

BIBLIOGRAPHY

[1] BERBERIAN, S. *Notes on spectral theory*. Van Nostrand, Princeton, N.J., 1966.

[2] —— *Lectures in functional analysis and operator theory*. Springer Verlag, New York-Heidelberg-Berlin, 1974.

[3] BERG, C., J. R. C. CHRISTENSEN and P. RESSEL. Positive definite functions on abelian semigroups. *Math. Ann. 223* (1976), pp. 253-272.
[4] BUCY, R. S. and G. MALTESE. A representation theorem for positive functionals on involution algebras. *Math. Ann. 162* (1966), pp. 364-367.
[5] —— Extreme positive definite functions and Choquet's representation theorem. *Jour. Math. Ann. Appl. 12* (1965), pp. 371-377.
[6] CHOQUET, G. *Lectures on analysis, vol. II.* W. A. Benjamin, Inc., New York, 1969.
[7] —— Deux exemples classiques de représentation intégrale. *Enseign. Math. 15* (1969), pp. 63-75.
[8] DIXMIER, J. *Les C*-algèbres et leurs représentations.* Gauthier-Villars, Paris, 1964.
[9] EBERLEIN, W. Characterizations of Fourier-Stieltjes transforms. *Duke Jour. Math. 22* (1955), pp. 465-468.
[10] GELFAND, I., D. RAIKOV and G. SILOV. *Commutative normed rings.* Chelsea, New York 1964.
[11] LOOMIS, L. *Abstract harmonic analysis.* Van Nostrand, New York, 1953.
[12] MALTESE, G. Multiplicative extensions of multiplicative functionals in Banach algebras. *Arch. Math. (Basel) 21* (1970), pp. 502-505.
[13] —— Convexity methods and the Choquet boundary in Banach algebras. *Bolletino Union Math. Ital. (5) 15-A* (1978), pp. 131-136.
[14] PHELPS, R. R. *Lectures on Choquet's theorem.* Van Nostrand, New York, 1966.
[15] RUDIN, W. *Fourier analysis on groups.* Interscience, New York, 1962.
[16] —— *Functional analysis.* McGraw-Hill, New York, 1973.
[17] SCHOENBERG, I. A remark on the preceding note by Bochner. *Bull. AMS 40* (1934), pp. 277-278.
[18] SEGAL, I. E. and R. KUNZE. *Integrals and operators.* McGraw-Hill, New York, 1968.

(Reçu le 15 février 1979)

George Maltese
Universität Münster
Münster, Germany