

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 25 (1979)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ACYCLIC MAPS
Autor: Hausmann, Jean-Claude / Husemoller, Dale
Kapitel: §1. Acyclic maps and homotopy equivalences
DOI: <https://doi.org/10.5169/seals-50372>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

§ 1. ACYCLIC MAPS AND HOMOTOPY EQUIVALENCES

We will use the terminology *CW-space* for a space having the homotopy type of a *CW-complex*. The category of *CW-spaces* is the largest category of spaces for which the Whitehead characterization of homotopy equivalences holds.

(1.1) DEFINITION. *A space X is acyclic provided the integral reduced homology $\tilde{H}_*(X) = 0$.*

In particular, an acyclic space X is path connected, its fundamental group $\pi_1(X)$ is perfect, i.e. $\pi_1(X)$ is equal to its commutator subgroup, and for any constant coefficient module L it follows that $\tilde{H}_*(X, L) = 0$. Recall that a local coefficient system L on X is a module over $\pi_1(X)$ and that

$$H_*(X, L) = H_*(C_*(X) \otimes_{\mathbf{Z}\pi_1(X)} L)$$

where $C_*(\tilde{X})$ is the chain complex over \mathbf{Z} viewed as a $\mathbf{Z}\pi_1(Y)$ -module. In general, $\tilde{H}(X, L) \neq 0$ for an acyclic space and a local coefficient system L .

(1.2) DEFINITION/PROPOSITION. *A map $f: X \rightarrow Y$ between path connected spaces is acyclic provided any of the following equivalent conditions hold:*

- (a) *The homotopy fibre F of $f: X \rightarrow Y$ is an acyclic space.*
- (b) *For any local coefficient system L on Y the induced morphism*

$$f_* : H_*(X, f^*L) \rightarrow H_*(Y, L)$$

*is an isomorphism where f^*L is the induced local system on X .*

- (c) *The induced morphism*

$$f_* : H_*(X, f^*\mathbf{Z}\pi_1(Y)) \rightarrow H_*(Y, \mathbf{Z}\pi_1(Y))$$

is an isomorphism.

- (d) *For the universal covering $\tilde{Y} \rightarrow Y$ of Y the map $X \times_Y \tilde{Y} \rightarrow \tilde{Y}$ defined by f induces an isomorphism*

$$H_*(X \times_Y \tilde{Y}) \rightarrow H_*(\tilde{Y}).$$

Proof. For (a) implies (b), we use the Serre spectral sequence for the fibration $F \xrightarrow{i} X \xrightarrow{f} Y$ where

$$E^2 = H_* (Y, H_* (F, i^* f^* L)) \Rightarrow H_* (X, f^* L).$$

Since $i^* f^* L$ is trivial on F , statement (a) gives $\tilde{H}_* (F, i^* f^* L) = 0$ and the edge morphism $H_* (X, f^* L) \rightarrow H_* (Y, L) = E^2_{*,0}$, which is induced by f , is an isomorphism.

Clearly (b) implies (c), which is a special case of (b), and for (c) implies (d) we use the following morphism of fibrations

$$\begin{array}{ccc} & \pi = \pi_1(Y) & \\ & \swarrow \quad \searrow & \\ X \times_Y \tilde{Y} & \longrightarrow & \tilde{Y} \\ \downarrow & & \downarrow \\ X & \xrightarrow{f} & Y. \end{array}$$

This induces a morphism of the Serre spectral sequences which on the E^2 -level is the given isomorphism from (c)

$$E^2 = H_* (X, f^* \mathbf{Z}\pi_1(Y)) \rightarrow H_* (Y, \mathbf{Z}\pi_1(Y)) = E^2.$$

Hence by the spectral mapping theorem $H_* (X \times_Y \tilde{Y}) \rightarrow H_* (\tilde{Y})$ is an isomorphism.

For (d) implies (a), note that $F \rightarrow X \times_Y \tilde{Y}$ is the fibre of $X \times_Y \tilde{Y} \rightarrow \tilde{Y}$. Since $H_* (X \times_Y \tilde{Y}) \rightarrow H_* (\tilde{Y})$ is an isomorphism on the horizontal edge of the spectral sequence, we see $\tilde{H}_0(F) = 0$. Moreover, assuming inductively that $\tilde{H}_j(F) = 0$ for $i < n$, we deduce that $\tilde{H}_n(F) \doteq 0$ by looking at the spectral sequence terms $E^r_{0,n}$ which is $H_n(F)$ for $r = 2$ and zero for $r > n + 1$. This completes the proof the equivalence of (a), (b), (c), and (d).

(1.3) PROPOSITION. *If $f : X \rightarrow Y$ is an acyclic map, then $f_* : \pi_1(X) \rightarrow \pi_1(Y)$ is an epimorphism with kernel a perfect normal subgroup.*

Proof. Since the fibre F of f is connected, the induced homomorphism f_* is an epimorphism, and since $\pi_1(F)$ is perfect, $\ker(f_*) = \text{im}(\pi_1(F) \rightarrow \pi_1(X))$ is perfect.

(1.4) PROPOSITION. Let $f: X \rightarrow Y$ be a map between path connected spaces. Then $\pi_i(f): \pi_i(X) \rightarrow \pi_i(Y)$ is an isomorphism for all $i \geq 0$ if and only if f is acyclic and $\pi_1(f)$ is an isomorphism.

Proof. Let $F \rightarrow X$ be the homotopy fibre of f . The second conditions say that $\pi_1(F)$ is perfect and abelian respectively. Thus $\pi_1(F) = 0$ and on simply connected spaces F the homotopy $\pi_i(F) = 0$ if and only if the homology $\tilde{H}_i(F) = 0$. The proposition follows now from an application of the homotopy exact sequence.

(1.5) COROLLARY. A map $f: X \rightarrow Y$ between path connected CW-spaces is a homotopy equivalence if and only if f is acyclic and $\pi_1(f)$ is an isomorphism.

This is an immediate application of the Whitehead criterion for homotopy equivalence applied to (1.4).

In section 3 we will see that the subgroups $\ker(\pi_1(f))$ classify acyclic maps $f: X \rightarrow Y$ from X .

(1.6) Remark. Cohomology with local coefficients can be used to characterize acyclic maps. As with homology

$$H^*(X, L) = H^*(\text{Hom}_{\mathbb{Z}\pi_1(X)}(C^*(\tilde{X}), L))$$

defines cohomology with local coefficients. Then a map $f: X \rightarrow Y$ between path connected spaces is acyclic if and only if $f^*: H^*(Y, L) \rightarrow H^*(X, f^*L)$ is an isomorphism for each local coefficient system L on Y . The direct implication is checked exactly as (a) implies (b) using cohomology in (1.2). Conversely we show that $X \times_Y \tilde{Y} \rightarrow \tilde{Y}$ defined by f induces an isomorphism $H^*(\tilde{Y}) \rightarrow H^*(X \times_Y \tilde{Y})$. This is done as (c) implies (d) in (1.2) and as in (d) implies (a) in (1.2) we have $\tilde{H}^*(F) = 0$. Using the universal coefficient theorem, we deduce that $\tilde{H}_*(F) = 0$ and F is acyclic.

The cohomology characterization of acyclic maps is useful in obstruction theory.

(1.7) Remark. Let $f: X \rightarrow Y$ be an acyclic map and \bar{Y} a connected covering of Y . Then the induced map $\bar{f}: X \times_Y \bar{Y} \rightarrow \bar{Y}$ is also acyclic. This follows directly from (1.2, (d)) or from the fact that f and \bar{f} have the same fibre. When \bar{Y} is the universal covering of Y , the space $X \times_Y \bar{Y} = \tilde{X}_N$ is the covering of X with fundamental group $N = \ker(\pi_1(f))$.