

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 25 (1979)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ACYCLIC MAPS  
**Autor:** Hausmann, Jean-Claude / Husemoller, Dale  
**Kapitel:** §6. ACYCLIC MAPS INTO A GIVEN SPACE  
**DOI:** <https://doi.org/10.5169/seals-50372>

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 16.01.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

We prove  $\tilde{X} \rightarrow F$  is a homotopy equivalence with the same argument used in (5.6) to show  $P_k$  implies  $H_k$ . Since  $F$  is also the fibre of  $X_N^+ \rightarrow [B\pi_1(X)]_N^+$  we have proved the theorem.

(5.8) *Remark.* Using (5.1), we see that for an acyclic map  $f: X \rightarrow Y$  which is  $k$ -simple for all  $k \geq 2$ , the homotopy groups  $\pi_*(Y)$  can be computed in terms of  $\pi_*(X)$  and  $\pi_*(B\pi_1(X)_N^+) \cong \pi_*(BN)^+$  for  $i \geq 2$ . Some computations of  $\pi_*(BN^+)$  for a certain perfect group  $N$  can be found for instance in [H, Chapter 7].

## § 6. ACYCLIC MAPS INTO A GIVEN SPACE

In this section we study acyclic maps  $f: X \rightarrow Y$  into a fixed space  $Y$ . Two such map  $f: X \rightarrow Y$  and  $f': X' \rightarrow Y$  are called equivalent provided there is a homotopy equivalence  $h: X \rightarrow X'$  with  $f \simeq f'h$ . Let  $AC(Y)$  denote the class of equivalence classes of acyclic  $f: X \rightarrow Y$  over  $Y$  where  $X$  and  $Y$  are  $CW$ -spaces.

(6.1) **DEFINITION.** *An extension data over a space  $Y$  is a triple  $(\Phi, i, \phi)$  where*

- (a)  $\Phi$  is an extension  $1 \rightarrow N \rightarrow G \rightarrow \pi_1(Y) \rightarrow 1$  with  $N$  perfect,
- (b)  $i: BG \rightarrow BG_N^+$  is an acyclic map with  $\ker(\pi_1(i)) = N$  (whose equivalence class is well defined by (3.5)), and
- (c)  $\phi: Y \rightarrow BG_N^+$  is a 2-connected map.

Two triples of extension data  $(\Phi, i, \phi)$  and  $(\Phi', i', \phi')$  are called equivalent provided there exists an isomorphism  $g: G \rightarrow G'$  making the following diagrams commutative (up to homotopy for the second one).

$$\begin{array}{ccccc}
 & & BG & \xrightarrow{Bg} & BG' \\
 & & i \downarrow & & \downarrow i' \\
 & & BG_N^+ & \xrightarrow{Bg^+} & B(G')_N^+ \\
 G & \xrightarrow{g} & G' & & \\
 & \searrow & \swarrow & & \\
 & & \pi_1(Y) & & \\
 & & & \nearrow \phi & \swarrow \phi' \\
 & & & Y & 
 \end{array}$$

where  $N' = g(N)$  and  $Bg^+$  is the unique homotopy equivalence determined by  $g$  with (3.1).

We denote by  $ED(Y)$  the class of equivalence classes of extension data.

(6.2) DEFINITION. *The data map  $\rho$  is the function  $\rho : AC(Y) \rightarrow ED(Y)$  which assigns to an acyclic map  $f : X \rightarrow Y$  the class  $\rho(f) = (\Phi, i, \phi)$  of extension data defined as follows :*

- (a)  $\Phi$  is the extension  $1 \rightarrow \ker \pi_1(f) \rightarrow \pi_1(X) \rightarrow \pi_1(Y) \rightarrow 1$ .
- (b) (c) With the well defined  $j : X \rightarrow BG$  for  $G = \pi_1(X)$  we form the cocartesian diagram

$$\begin{array}{ccc} X & \xrightarrow{j} & BG \\ f \downarrow & & \downarrow i \\ Y & \xrightarrow{\phi} & Y \cup_X BG \end{array}$$

Since  $f$  is acyclic,  $i$  is acyclic, and since  $\pi_1(j)$  is an isomorphism,  $\ker(\pi_1(i)) = N$ . Thus  $Y \cup_X BG$  is  $BG_N^+$  up to equivalence.

Now we have to check that the map  $\phi : Y \rightarrow Y \cup_X BG = BG_N^+$  is 2-connected. Since  $\pi_1(j)$  is an isomorphism,  $\pi_1(\phi)$  is also an isomorphism. The fact that  $\pi_2(\phi)$  is surjective comes from the diagram.

$$\begin{array}{ccccccc} \pi_2(Y) & \xleftarrow{\sim} & \pi_2(\tilde{Y}) & \xrightarrow{\sim} & H_2(\tilde{Y}) & \xleftarrow{\sim} & H_2(\tilde{X}_N) \\ \pi_2(\phi) \downarrow & & \downarrow \pi_2(\tilde{\phi}) & & \downarrow & & \searrow \\ \pi_2(BG_N^+) & \xleftarrow{\sim} & \pi_2(BN^+) & \xrightarrow{\sim} & H_2(N) & & \end{array}$$

The surjectivity on the right is a classical result of Hopf which follows easily from the Serre spectral sequence of the fibration  $\tilde{X} \rightarrow \tilde{X}_N \rightarrow BN$ .

Now using (2.5) a simple argument, left to the reader, shows that  $\rho : AC(Y) \rightarrow ED(Y)$  is well defined.

(6.3) THEOREM. *Let  $Y$  be a CW-space. The map  $\rho : AC(Y) \rightarrow ED(Y)$  surjective and its restriction to the subclass  $AC_S(Y)$  of  $AC(Y)$  of  $f : X \rightarrow Y$  which are  $k$ -simple for all  $k \geq 2$  is a bijection.*

*Proof.* To show  $\rho$  is surjective, consider extension data  $(\Phi, i, \phi)$  and form the cartesian square

$$\begin{array}{ccc} X = Y \times {}_T BG & \xrightarrow{\alpha} & BG \\ f \downarrow & & \downarrow i \\ Y & \xrightarrow{\phi} & BG_N^+ = T \end{array}$$

Now  $f$  is acyclic by (2.2), and since its fiber is the same as  $i$ , we deduce by (5.2) that  $f$  is  $k$ -simple for all  $k \geq 2$ .

Next, let  $\rho(f) = (\Phi_0, i_0, \phi_0)$  and we show this extension data is equivalent to  $(\Phi, i, \phi)$ . Using the homotopy exact sequences for  $X \rightarrow Y$  and  $BG \rightarrow BG_N^+$  and the fact that  $\phi$  is 2-connected, we deduce from the five lemma that  $\pi_1(\alpha) : \pi_1(X) \rightarrow G$  is an isomorphism. The following diagram shows that  $(\Phi_0, i_0, \phi_0)$  is equivalent to  $(\Phi, i, \phi)$  and  $\rho$  is surjective.

$$\begin{array}{ccccc} & & j & & B\pi_1(X) \\ & X & \xrightarrow{\alpha} & BG & B\pi_1(\alpha) \\ f \downarrow & & & \downarrow i & \downarrow i_0 \\ Y & \xrightarrow{\phi} & BG_N^+ & & B\pi_1(\alpha)^+ \\ & & \phi_0 & & Y \cup {}_X B\pi_1(X) \end{array}$$

Now, if  $f: X \rightarrow Y$  is an acyclic map which is  $k$ -simple for all  $k \geq 2$  and with  $\rho(f) = (\Phi, i, \phi)$ , then we form the following commutative diagram.

$$\begin{array}{ccccccc} X & \xrightarrow{d} & Y \times {}_T BG & \longrightarrow & BG & & \\ & f \searrow & \downarrow f_0 & & \downarrow i & (G = \pi_1(X)) & \\ & & Y & \xrightarrow{\phi} & BG_N^+ & & \end{array}$$

As we have seen in the proof the surjectivity of  $\rho$ , the map  $f_0$  is acyclic and  $k$ -simple for  $k \geq 2$ . The map  $d$  induces an isomorphism on the fundamental groups and on homology with  $\mathbf{Z}\pi_1(Y)$  twisted coefficients. By (5.3), the map  $d$  is a homotopy equivalence. This proves that the acyclic map  $f$  is equivalent to the induced map  $f_0$ . Thus  $\rho$  restricted to  $AC_S(U) \rightarrow ED(Y)$  is a bijection.

(6.4) *Remark.* This theorem leaves open the question of the fibres of the function.

$$\rho : AC(Y) \rightarrow ED(Y).$$

In the next theorem we factor an acyclic map by ones having simplicity properties.

(6.5) *Remark.* In theorem (6.3), if one fixes an extension  $\Phi : 1 \rightarrow N \rightarrow G \rightarrow \pi_1(Y) \rightarrow 1$ , then the same proof permits us to classify acyclic maps  $f : X \rightarrow Y$  which are  $k$ -simple for  $k > 2$  together with an identification  $d : \pi_1(X) \rightarrow G$  such that  $\Phi d = \pi_1(f)$ . The objects of  $ED(Y)$  have to be replaced by couples  $(i, \phi)$  where  $i : BG \rightarrow BG_N^+$  is as above and  $\phi : Y \rightarrow BG_N^+$  is 2-connected with the following diagram commuting up to homotopy.

$$\begin{array}{ccc} B\pi_1(Y) & \xrightarrow{B\Phi} & BG \\ \nearrow & \searrow B\Phi^+ & \downarrow i \\ Y & \xrightarrow{\phi} & BG_N^+ \end{array}$$

This is what is done implicitly in [H, Sections 2 and 4]. Observe that we are dealing here with classes which are sets.

(6.6) *LEMMA.* *Let  $X$  be a CW-space and  $N$  a perfect normal subgroup of  $\pi_1(X)$ . Let  $X \rightarrow P_n X$  denote the  $n$ th stage of the Postnikov decomposition of  $X$ . Then for all  $n \geq 1$  we have that*

- (1)  $\pi_j(X_N^+) \rightarrow \pi_j((P_n X)_N^+)$  is an isomorphism for  $j \leq n$  and an epimorphism for  $j = n + 1$ , and
- (2)  $\pi_j(A\tilde{X}_N) \rightarrow \pi_j(A(P_n \tilde{X}_N))$  is an isomorphism for  $j \leq n$  and an epimorphism for  $j = n + 1$ .

*Proof.* Consider the following homotopy commutative diagram of fibre sequences

$$\begin{array}{ccccc} T & \longrightarrow & A\tilde{X}_N & \longrightarrow & A(P_n \tilde{X}_N) \\ \downarrow & & \downarrow & & \downarrow \\ F & \longrightarrow & \tilde{X}_N & \longrightarrow & P_n \tilde{X}_N \\ \downarrow & & \downarrow & & \downarrow \\ G & \longrightarrow & (\tilde{X}_N)^+ & \longrightarrow & (P_n \tilde{X}_N)^+ \end{array}$$


Clearly  $\pi_i(F) = 0$  for  $i \leq n + 1$ . The spaces  $\tilde{X}_N$  and  $P_n \tilde{X}_N$  have the same  $(n+1)$ -skeleton and the same can be assumed for  $\tilde{X}_N^+$  and  $(P_n \tilde{X}_N)^+$ . Hence  $\pi_i(G) = 0$  for  $i \leq n + 1$ . Now (1) follows because  $G$  is the fibre of  $X_N^+ \rightarrow (P_n X)^+$ .

By comparing Serre spectral sequences, we obtain the surjectivity of

$$H_0(N, H_{n+1}(F)) \rightarrow H_0(N, H_{n+1}(G)) = H_{n+1}(G) = \pi_{n+1}(G).$$

Thus  $\pi_j(T) = 0$  for  $j \leq n$  and (2) follows.

(6.7) THEOREM. *Let  $f: X \rightarrow Y$  be a map between CW-spaces. Then there is a factorization*



such that  $\beta_i$  is  $i$ -connected and  $\alpha_i$  is an acyclic map which is  $k$ -simple for  $k > i$ .

Such a decomposition is unique up to a homotopy equivalence.

*Proof.* The  $i$ th stage  $X_i$  is defined by the cartesian diagram

$$\begin{array}{ccc} Y \times_T P_i(X) & \longrightarrow & P_i X \\ \downarrow & & \downarrow \\ Y & \longrightarrow & (P_i X)^+ = T \end{array}$$

where  $N = \ker(\pi_1(X) \rightarrow \pi_1(Y))$ . By (6.6) the map  $\beta_i$  is  $i$ -connected since the fiber of the two vertical arrows is  $A(P_n \tilde{X})_N$ . Now by (5.4) we see that  $\alpha_i$  is simple for  $k > i$ .

For two decompositions  $(X'_i)$  and  $(X''_i)$  of  $f: X \rightarrow Y$  satisfying the above conditions, we have  $P_i X'_i = P_i X''_i$  and both  $X'_i$  and  $X''_i$  map into  $X_i$ , constructed above, such that the resulting diagrams are homotopy commutative. The connectivity of the  $\beta_i$  and (5.1) shows that these maps are all homotopy equivalences. This proves the theorem.

(6.8) *Remarks.* This theorem (6.7) coincides with the Dror results for  $Y$  a point [D1, Theorem 1.3] and  $Y = S^n$  [D2]. An interesting problem is to describe the  $i$ th stage  $X_i$  in terms of invariants of  $X_{i-1}$  as in [D1] and [D2]. (See the footnote in the introduction.)

#### APPENDIX — SIMPLICITY PROPERTIES OF FIBERS

In the proof of (5.4) we used the fact that for a fibration  $F \rightarrow E \xrightarrow{f} B$  the action of  $\pi_1(F)$  on  $\text{Im}(\partial: \pi_{k+1}(B) \rightarrow \pi_k(F))$  is trivial. This assertion does not seem to be in the literature so we include a proof here.

We extend the mapping sequence of the fibration  $f$  to  $\Omega B \rightarrow F \rightarrow E \xrightarrow{f} B$  and study  $F$  as the total space of a principal fibration with fibre the  $H$ -space  $\Omega B$ . If  $G$  is an  $H$ -space, then  $\pi_1(G)$  acts trivially on  $\pi_*(G)$  because the covering transformations  $\tilde{G} \rightarrow G$  on the universal covering  $\tilde{G}$  of  $G$  are homotopic to the identity. This is proved by lifting a loop to a path in  $\tilde{G}$  and using the  $H$ -space structure on  $\tilde{G}$  to deform the identity along this path to the covering transformation defined by the homotopy class of the loop. Recall that a principal fibration is induced from  $G \rightarrow E_G \rightarrow B_G$  up to fibre homotopy equivalence.

(A.1) **PROPOSITION.** *Let  $G \rightarrow X \xrightarrow{\pi} Y$  be a principal fibration with fibre  $G$  acting on  $X$ . Then we have :*

- (a)  $\text{im}(\pi_1(G) \rightarrow \pi_1(X))$  acts trivially on  $\pi_*(X)$ , and
- (b)  $\pi_1(X)$  acts trivially on  $\text{im}(\pi_*(G) \rightarrow \pi_*(X))$ .

*Proof.* For (a) we have the following commutative diagram induced by a covering transformation  $T: \tilde{G} \rightarrow \tilde{G}$ .