Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 25 (1979)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ACYCLIC MAPS
Autor: Hausmann, Jean-Claude / Husemoller, Dale
Anhang: APPENDIX — SIMPLICITY PROPERTIES OF FIBERS
DOI: https://doi.org/10.5169/seals-50372

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 14.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

where $N = \ker(\pi_1(X) \to \pi_1(Y))$. By (6.6) the map β_i is i-connected since the fiber of the two vertical arrows is $A(P_n\tilde{X})_N$. Now by (5.4) we see that α_i is simple for $k > i$.

For two decompositions (X'_i) and (X''_i) of $f : X \to Y$ satisfying the above conditions, we have $P_iX'_i = P_iX''_i$ and both X'_i and X''_i map into X_i, constructed above, such that the resulting diagrams are homotopy commutative. The connectivity of the β_i and (5.1) shows that these maps are all homotopy equivalences. This proves the theorem.

(6.8) *Remarks.* This theorem (6.7) coincides with the Dror results for Y a point [D1, Theorem 1.3] and $Y = S^n$ [D2]. An interesting problem is to describe the ith stage X_i in terms of invariants of X_{i-1} as in [D1] and [D2]. (See the footnote in the introduction.)

Appendix — Simplicity properties of fibers

In the proof of (5.4) we used the fact that for a fibration $F \to E \xrightarrow{f} B$ the action of $\pi_1(F)$ on $\operatorname{Im}(\partial : \pi_{k+1}(B) \to \pi_k(F))$ is trivial. This assertion does not seem to be in the literature so we include a proof here.

We extend the mapping sequence of the fibration f to $\Omega B \to F \to E \xrightarrow{f} B$ and study F as the total space of a principal fibration with fibre the H-space ΩB. If G is an H-space, then $\pi_1(G)$ acts trivally on $\pi_*(G)$ because the covering transformations $\tilde{G} \to G$ on the universal covering $\tilde{G}$ of G are homotopic to the identity. This is proved by lifting a loop to a path in $\tilde{G}$ and using the H-space structure on $\tilde{G}$ to deform the identity along this path to the covering transformation defined by the homotopy class of the loop. Recall that a principal fibration is induced from $G \to E_G \to B_G$ up to fibre homotopy equivalence.

(A.1) Proposition. *Let* $G \to X \xrightarrow{\pi} Y$ *be a principal fibration with fibre* G *acting on* X. *Then we have:*

(a) $\operatorname{im}(\pi_1(G) \to \pi_1(X))$ *acts trivially on* $\pi_*(X)$, *and*

(b) $\pi_1(X)$ *acts trivially on* $\operatorname{im}(\pi_*(G) \to \pi_*(X))$.

Proof. For (a) we have the following commutative diagram induced by a covering transformation $T : \tilde{G} \to \tilde{G}$.

$$\begin{tikzcd}
 & \tilde{G} \arrow[rr] \arrow[dd] & & \tilde{G} \times \tilde{X} \arrow[rr] \arrow[dd] & & \tilde{X} \arrow[dd] \\
\tilde{G} \arrow[ur, "T"] \arrow[rr] \arrow[dr] & & \tilde{G} \times \tilde{X} \arrow[ur, "T \times 1"] \arrow[rr] \arrow[dr] & & \tilde{X} \arrow[ur, "T'"] \arrow[dr] & \\
 & G \arrow[rr] & & G \times X \arrow[rr] & & X
\end{tikzcd}$$

The covering transformation T defines T', and since T is homotopic to the identity so is T'. This proves (a).

For (b) we use the following commutative diagram where T' is any covering transformation of $\tilde{X}$.

$$\begin{tikzcd}
\tilde{G} \arrow[rr] \arrow[dr] & & \tilde{G} \times \tilde{X} \arrow[rr] \arrow[dd] & & \tilde{X} \arrow[dd] \\
 & \tilde{G} \times \tilde{X} \arrow[ur, "1 \times T'"] \arrow[rr] \arrow[dr] & & \tilde{X} \arrow[ur, "T'"] \arrow[dr] & \\
 & & G \times X \arrow[rr] & & X
\end{tikzcd}$$

Now the inclusion $i : \tilde{G} \to \tilde{X}$ is the composite of the first horizontal row, and $T'i$ and i are homotopic by $i_t(g) = g \cdot \tilde{\alpha}(t)$ where $g \in G$ and α is a lifting of the loop α corresponding to the covering transformation T'. This proves the proposition.

For a general fibration $f : E \to B$ with fibre F the mapping sequence $\Omega B \to F \to E \to B$ allows us to deduce the next proposition from the previous one.

(A.2) Proposition. *Let $f : E \to B$ be a fibration with fibre $F \to E$. Then we have:*

(a) $\operatorname{im}(\pi_2(B) \to \pi_1(F))$ *acts trivially on* $\pi_*(F)$, *and*

(b) $\pi_1(F)$ *acts trivially on* $\operatorname{im}(\partial : \pi_{i+1}(B) \to \pi_i(F))$.