

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 25 (1979)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ON LIE'S HIGHER SPHERE GEOMETRY  
**Autor:** Fillmore, Jay P.  
**Kapitel:** 1. Introduction  
**DOI:** <https://doi.org/10.5169/seals-50373>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# ON LIE'S HIGHER SPHERE GEOMETRY

by Jay P. FILLMORE

## 1. INTRODUCTION

In this paper we draw together two theories having their roots in the ideas of S. Lie over a century ago: *Lie's higher sphere geometry*, with its famous *line-sphere transformation* [5], and the theory of Lie groups, especially the description of a geometry by global Lie groups<sup>1</sup>. Indeed, not until the 1960s, with the appearance of W. M. Boothby's description of homogeneous contact manifolds [1, 2] and with the appearance of parabolic subgroups, could this connection be established. One can now say, in terms of Lie groups, that the three-dimensional complex line and sphere geometries are isomorphic and that *the real line and sphere geometries are two distinct real forms of one geometry*. Furthermore, the line-sphere transformation gives explicitly the isomorphism of the complex forms.

In Section 2 we summarize the formulation of Boothby's theory for algebraic homogeneous contact manifolds and make some observations about their real forms. The classical contact manifolds of complex co-directions in projective space and of Lie's higher sphere geometry are described in general in terms of this theory in Sections 3 and 4. Finally, in Section 5, the connection with Plücker's line geometry in three dimensions is established, and the line-sphere transformation is brought into perspective. This introduction continues with an overview of F. Klein's formulation of Lie's theory [5, 6], Boothby's theory, and their connection.

To a line in complex projective space  $P^3$  may be assigned Plücker coordinates

$$\begin{aligned}\xi_1 &= p_{12}, \quad \xi_2 = p_{31}, \quad \xi_3 = p_{23}, \\ \xi_4 &= p_{03}, \quad \xi_5 = p_{02}, \quad \xi_6 = p_{01},\end{aligned}$$

[6, §20]. These coordinates satisfy

$$\xi_1\xi_4 + \xi_2\xi_5 + \xi_3\xi_6 = 0,$$

---

<sup>1</sup> This description of Lie's higher sphere geometry in terms of Lie groups answers a question posed in 1965 by S. SASAKI [9, p. 173].

and hence lines in  $P^3$  correspond to points of a quadric  $\Omega^4$  in  $P^5$ . Two lines in  $P^3$  intersect when their corresponding points on  $\Omega^4$  are conjugate. A surface element in  $P^3$ , a point and incident plane, becomes the pencil of lines passing through the point and lying in the plane; this corresponds to a line lying in  $\Omega^4$ . The space of surface elements in  $P^3$  thus corresponds to the space of lines in  $\Omega^4$ . The projectivities of  $P^5$  which preserve the quadric  $\Omega^4$  permute the lines of  $\Omega^4$  and hence the surface elements of  $P^3$ . Moreover, these projectivities preserve the condition, between two surface elements at infinitesimally adjacent points, that a point of one lies on the plane of the other; hence they are contact transformations of  $P^3$ .

To a sphere

$$x^2 + y^2 + z^2 - 2ax - 2by - 2cz + C = 0$$

in complex Euclidean space  $E^3$ , with center at  $x = a$ ,  $y = b$ ,  $z = c$  and radius

$$r^2 = a^2 + b^2 + c^2 - C,$$

the sign of  $r$  corresponding to an "orientation", may be assigned homogeneous coordinates

$$a = \frac{\alpha}{v}, \quad b = \frac{\beta}{v}, \quad c = \frac{\gamma}{v}, \quad r = \frac{\lambda}{v}, \quad C = \frac{\mu}{v},$$

[6, §25]. These coordinates satisfy

$$\alpha^2 + \beta^2 + \gamma^2 - \lambda^2 - \mu v = 0,$$

and hence oriented spheres in  $E^3$  correspond to certain points of a quadric  $\Psi^4$  in  $P^5$ ; if spheres which are points or planes or which have centers at infinity are included, all points of  $\Psi^4$  are obtained. Two spheres in  $E^3$  are tangent at a point, orientations taken into account, when their corresponding points on  $\Psi^4$  are conjugate. An "oriented" surface element in  $E^3$ , a point and incident oriented plane, becomes the pencil of spheres tangent to the plane at the point; this corresponds to a line lying in  $\Psi^4$ . The space of oriented surface elements of  $E^3$  thus corresponds to the space of lines in  $\Psi^4$ . The projectivities of  $P^5$  which preserve the quadric  $\Psi^4$  permute the lines of  $\Psi^4$  and hence the oriented surface elements of  $E^3$ . Moreover, these projectivities are contact transformations of  $E^3$ .

The line-sphere transformation, discovered by Lie, is given by

$$\begin{aligned} \xi_1 &= \alpha + \sqrt{-1} \beta, & \xi_4 &= \alpha - \sqrt{-1} \beta, \\ \xi_2 &= \gamma + \lambda, & \xi_5 &= \gamma - \lambda, \\ \xi_3 &= \mu, & \xi_6 &= -v, \end{aligned}$$

as formulated by Klein [6, §70]. This makes correspond points of the quadric  $\Omega^4$  of signature  $(+++--)$  and points of the quadric  $\Psi^4$  of signature  $(++++--)$ . Conjugate points correspond to conjugate points, and a line in one quadric corresponds to a line in the other. Thus, surface elements in  $P^3$  correspond to oriented surface elements in  $E^3$  and this correspondence is a "contact transformation".

Now, classically a *contact transformation* in  $P^3$  or  $E^3$  is a transformation on the 5-dimensional space of surface elements which preserves, up to a non-vanishing multiple, a maximal rank Pfaffian form

$$\omega = dz - p dx - q dy,$$

[6, §63], where the coordinates  $x, y, z, p, q$  describe the surface element consisting of the plane

$$z' - z = p(x' - x) + q(y' - y)$$

at the point  $(x, y, z)$ . The condition  $\omega = 0$ , that at two infinitesimally adjacent points the point of one surface element lies on the plane of the other, is preserved by a contact transformation. The appropriate spaces for the line-sphere transformation are the 5-dimensional spaces of lines in  $\Omega^4$  and lines in  $\Psi^4$ . Exhibiting the Pfaffian forms and examining the effect of the line-sphere transformation on them may be done systematically by observing that these spaces are *homogeneous*.

Boothby's description of compact homogeneous complex contact manifolds [1, 2; and 7, §2] constructs for each type of simple complex Lie algebra  $\mathfrak{g}$ : a connected centerless simple Lie groups  $G$  having Lie algebra  $\mathfrak{g}$ , a parabolic subgroup  $P$  of  $G$ , and a Pfaffian form  $\omega$  on a principal  $\mathbb{C}^*$ -bundle over  $G/P$ , so that  $G/P$ , with  $\omega$  pulled down by local sections, is a compact complex contact manifold, homogeneous under the identity component  $G$  of the group of all its contact automorphisms. Every such contact manifold is so obtained uniquely up to isomorphism. This construction yields, for the classical simple Lie algebras:

- |                 |   |
|-----------------|---|
| $A_n$           | projective cotangent bundle of $P^n$ —the classical space of incident point-hyperplane pairs in $P^n$ , |
| $B_l$ and $D_l$ | space of lines in a quadric,  |
| $C_l$           | odd-dimensional projective space $P^{2l+1}$ ,   |

[1, (7.1)]. The isomorphism  $A_3 \simeq D_3$  arises from the description of surface elements in  $P^3$  as lines in  $\Omega^4$  by Plücker coordinates. Since the

complex quadrics  $\Omega^4$  and  $\Psi^4$  both have groups of projectivities of the type  $D_3$ , the contact manifolds of line geometry and sphere geometry, when viewed as the spaces of lines in  $\Omega^4$  and  $\Psi^4$  respectively, are necessarily *the same*, that is, isomorphic.

When Boothby's description of homogeneous contact manifolds is refined, using J. A. Wolf's theory of complex flag manifolds [8, Ch. I], to include their real forms, line geometry and sphere geometry are *no longer the same*, but, as was classically recognized [6, §25], are obtained from the real forms  $PSO(3, 3; \mathbf{R})$  and  $PSO(4, 2; \mathbf{R})$  of  $PSO(3, 3; \mathbf{C})$  and  $PSO(4, 2; \mathbf{C})$ , where the quadratic forms defining these projective special orthogonal groups are those of the quadrics  $\Omega^4$  and  $\Psi^4$ . Now,  $PSO(3, 3; \mathbf{C})$  and  $PSO(4, 2; \mathbf{C})$  are isomorphic, so the corresponding complex contact manifolds are isomorphic; in fact, these groups, are conjugate in  $PSL(6; \mathbf{C})$  by the matrix of Klein's description of the line-sphere transformation. Viewed another way,  $PSO(3, 3; \mathbf{R})$  and  $PSO(4, 2; \mathbf{R})$  correspond to two real forms of  $PSO(3, 3; \mathbf{C})$  defined by two complex conjugations. Consequently, *real line geometry and real sphere geometry are two distinct real forms of complex line geometry*. The line-sphere transformation then corresponds to an automorphism of  $PSO(3, 3; \mathbf{C})$  connecting the two complex conjugations.

