Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 25 (1979)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON LIE'S HIGHER SPHERE GEOMETRY
Autor: Fillmore, Jay P.
Kapitel: 3. CO-DIRECTIONS IN PROJECTIVE SPACE
DOI: https://doi.org/10.5169/seals-50373

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 26.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

is one-to-one on an open neighborhood U of x_0 in G/P and $(\exp X) \cdot x_0$ is identifiable as the point $(x_1, \ldots, x_n)$ and the incident hyperplane

$$x_n' - x_n = p_1 (x_1' - x_1) + \ldots + p_{n-1} (x_{n-1}' - x_{n-1}) .$$

Now, $(\exp X) \cdot x_0 \to (\exp X) \cdot b_0$ is a section of the bundle G/P_1 over U and, via this section, the form ω on G/P_1 pulls down to

$$\omega_0 ((\exp X)^{-1} d (\exp X))$$

which, when expressed in terms of $x_1, \ldots, x_n, p_1, \ldots, p_{n-1}$, will be identified with

$$dx_n - p_1 dx_1 - \ldots - p_{n-1} dx_{n-1}$$

up to a constant multiple $a \neq 0$. For this latter calculation we will use

$$(\exp X)^{-1} d (\exp X) = \frac{1 - e^{-ad\, X}}{ad\, X} (dX)$$

$$= dX - \frac{1}{2} [X, dX] + \frac{1}{6} [X, [X, dX]] - \ldots$$

[4, (10.2)], a series which is finite since $\mathfrak{m}$ is nilpotent. In fact, our choice of X will make the series for $\exp X$ themselves finite. The constant $a \neq 0$ could be made unity by using instead the section $(\exp X) \cdot x_0 \to (\exp X) g^{-1} \cdot b_0$, where g in P is chosen so that $\chi (g) = a$. This amounts to following the original section by R_a^{-1} in the bundle.

3. Co-directions in projective space

The contact structure on the $(2n-1)$-dimensional space of co-directions in complex projective space P^n, described in 2.5, is obtained when the construction of 2.10 is carried out for the simple complex Lie algebra of type A_n, $n \geqslant 1$.

3.1 Let $\mathfrak{g} = \mathfrak{sl} (n + 1; \mathbf{C})$, complex $(n+1)$ by $(n+1)$ matrices of trace zero. For Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$ take the diagonal matrices of $\mathfrak{g}$. Let δ_i, $i = 0,1, \ldots, n$ be the linear function on $\mathfrak{h}$ which assigns to $H = \operatorname{diag} (h_1, \ldots, h_n)$ in $\mathfrak{h}$ the ith diagonal element: $\delta_i (H) = h_i$. The roots of $\mathfrak{g}$ with respect to $\mathfrak{h}$ are

$$\delta_i - \delta_j \quad i, j = 0, 1, \ldots, n$$
$$\text{and } i \neq j$$

and the root vector E_α corresponding to the root α is

$$E_{\delta_i - \delta_j} = E_{ij},$$

the matrix with 1 in the ith row and j^{th} column and 0s elsewhere [4, (16.2)]. A system of simple roots is

$$\delta_0 - \delta_1, \delta_1 - \delta_2, \ldots, \delta_{n-1} - \delta_n,$$

for which the maximal root is

$$\rho = (\delta_0 - \delta_1) + (\delta_1 - \delta_2) + \ldots + (\delta_{n-1} - \delta_n) = \delta_0 - \delta_n$$

[4, App., Table E]. The Killing form of $\mathfrak{g}$ is $\langle X, Y \rangle = 2(n+1)\,\mathrm{tr}\,(XY)$, but we replace this with $\langle X, Y \rangle = \mathrm{tr}\,(XY)$ for convenience. Then the H_α in $\mathfrak{h}$ are given by

$$H_{\delta_i - \delta_j} = \mathrm{diag}(0, \ldots, 0, 1, 0, \ldots, 0, -1, 0, \ldots, 0)$$

with 1 and -1 in the ith and j^{th} entry, respectively. Especially,

$$H_\rho = \mathrm{diag}\,(1, 0, \ldots, 0, -1).$$

We have

$$\langle H_\rho, H_{\delta_i - \delta_j} \rangle \begin{cases} < 0 & j = 0 \text{ or } i = n \\ \geqslant 0 & \text{otherwise} \end{cases},$$

so that $\mathfrak{p}$ in (i) of 2.9 consists of matrices of the form

$$\begin{bmatrix} * & * & \cdots & * \\ 0 & \vdots & & \vdots \\ \vdots & * & \cdots & * \\ 0 & \cdots & 0 & * \end{bmatrix}$$

of trace zero, where the starred entries are arbitrary.

3.2 The connected centerless simple group $G = PSL\,(n+1; \mathbf{C}) = SL\,(n+1; \mathbf{C})/\{\text{center}\}$ is transitive on the space consisting of points x and incident hyperplanes u, $ux = 0$, in P^n, as in 2.5. The isotropy subgroup P of the incident point and hyperplane

$$x_0 = {}^t(1, 0, \ldots, 0), \quad u_0 = (0, \ldots, 0, 1)$$

has exactly $\mathfrak{p}$ for its Lie algebra. Hence, the homogeneous contact manifold which the construction of 2.10 gives is

$$\begin{aligned} G/P &= \text{space of incident points and hyperplanes in } P^n \\ &= \text{space of co-directions in complex } P^n. \end{aligned}$$

3.3 Let $\mathfrak{m}$ be the $(2n-1)$-dimensional supplement to $\mathfrak{p}$ in $\mathfrak{g}$ consisting of matrices of the form

$$\begin{bmatrix} 0 & & & \\ * & & 0 & \\ \vdots & & & \\ * & \cdots & * & 0 \end{bmatrix};$$

cf. 2.12. The product of any two matrices of $\mathfrak{m}$ has a nonzero entry only in the nth row and 0th column; the product of any three is zero. Set

$$X = \begin{bmatrix} 0 & & \\ x_1 & & 0 \\ \vdots & & \\ x_{n-1} & & \\ x_n - \frac{1}{2}\sum p_i x_i & p_1 \cdots p_{n-1} & 0 \end{bmatrix},$$

where the summation is over $i = 1, 2, \ldots, n-1$. X is in $\mathfrak{m}$ and

$$\exp X = 1_{n+1} + X + \frac{1}{2}X^2 = \begin{bmatrix} 1 & & & \\ x_1 & \ddots & 0 & \\ \vdots & & \ddots & \\ x_{n-1} & 0 & & \\ x_n & p_1 \cdots p_{n-1} & & 1 \end{bmatrix},$$

$$(\exp X)^{-1} = 1_{n+1} - X + \frac{1}{2}X^2 = \begin{bmatrix} 1 & & & & \\ -x_1 & \ddots & & 0 & \\ \vdots & & & & \\ -x_{n-1} & & 0 & & \\ -x_n + \sum p_i x_i & -p_1 & \dots & -p_{n-1} & 1 \end{bmatrix}.$$

The point

$$x = (\exp X) \cdot x_0 = {}^t(1, x_1, \dots, x_n)$$

is incident with the hyperplane

$$u = u_0 \cdot (\exp X)^{-1} = (-x_n + \sum p_i x_i, -p_1, \dots, -p_{n-1}, 1),$$

and the hyperplane $ux' = 0$, $x' = {}^t(1, x_1', \dots, x_n')$, is

$$x_n' - x_n = p_1(x_1' - x_1) + \dots + p_{n-1}(x_{n-1}' - x_{n-1}).$$

Thus, this choice of X establishes the classically identifiable coordinates $x_1, \dots, x_n, p_1, \dots, p_{n-1}$ on G/P.

3.8 From $\rho = \delta_0 - \delta_n$, we have $W = E_\varrho = E_{0n}$ in (iii) of 2.9 and $\omega_0(X) = \langle W, X \rangle$ is the $n0$-entry of X. The form ω on G/P is obtained as $\omega = \omega_0((\exp X)^{-1} d(\exp X))$ with

$$(\exp X)^{-1} d(\exp X) = dX - \frac{1}{2}[X, dX]$$

as in 2.12. For X as in 3.3, the only nonzero entry in $[X, dX]$ is the $n0^{th}$ and it is $\sum p_i\, dx_i = -\sum x_i\, dp_i$. Hence

$$(\exp X)^{-1} d(\exp X) = \begin{bmatrix} 0 & & & & \\ dx_1 & & & 0 & \\ \vdots & & & & \\ dx_{n-1} & & & & \\ dx_n - \sum p_i dx_i & dp_1 & \dots & dp_{n-1} & 0 \end{bmatrix},$$

and the $n0$-entry is

$$\omega = dx_n - p_1 dx_1 - \ldots - p_{n-1} dx_{n-1} .$$

This identifies the contact structure with the classical one as in 2.12.

3.5 The real contact structure on the $(2n-1)$-dimensional space of co-directions in real projective space P^n is described by viewing all quantities in the foregoing discussion as being real. Especially, G_0 of 2.11 is the connected centerless group $PSL(n+1; \mathbf{R})$ consisting of real contact automorphisms.

4. Higher sphere geometry

4.1 In complex Euclidean space E^n, the equation

$$x_1'^2 + \ldots + x_n'^2 - 2a_1 x_1' - \ldots - 2a_n x_n' + C = 0$$

describes a sphere with center $(a_1, \ldots, a_n)$ and complex radius r given by

$$r^2 = a_1^2 + \ldots + a_n^2 - C .$$

When $r \neq 0$, the two choices of sign for r is said to give two "orientations" to the sphere. Thus, the $n+2$ coordinates $a_1, \ldots, a_n, r, C$, which are related by

$$a_1^2 + \ldots + a_n^2 - r^2 - C = 0 ,$$

describe the space of oriented spheres in E^n [6, §25].

Introduce homogeneous coordinates by

$$a_i = \frac{\alpha_i}{\nu}, \; r = \frac{\lambda}{\nu}, \; C = \frac{\mu}{\nu} ,$$

$i = 1, 2, \ldots, n$. Then the oriented spheres of E^n correspond to certain points of the quadric Ψ^{n+1} in P^{n+2} described by

$$\alpha_1^2 + \ldots + \alpha_n^2 - \lambda^2 - \mu\nu = 0 .$$

The sphere corresponding to the point $(\alpha_1, \ldots, \alpha_n, \lambda, \mu, \nu)$ of Ψ^{n+1} is

$$\nu(x_1'^2 + \ldots + x_n'^2) - 2\alpha_1 x_1' - \ldots - 2\alpha_n x_n' + \mu = 0 .$$

Ordinary spheres have finite nonzero radius r, so $\nu \neq 0$. For $\nu = 0$, we obtain oriented hyperplanes. For $\lambda = 0$, we obtain point spheres or hyperplanes with isotropic hyperplane coordinate vector; these carry no