

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 25 (1979)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON LIE'S HIGHER SPHERE GEOMETRY
Autor: Fillmore, Jay P.
Kapitel: 4. HIGHER SPHERE GEOMETRY
DOI: <https://doi.org/10.5169/seals-50373>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

and the $n0$ -entry is

$$\omega = dx_n - p_1 dx_1 - \dots - p_{n-1} dx_{n-1}.$$

This identifies the contact structure with the classical one as in 2.12.

3.5 The real contact structure on the $(2n-1)$ -dimensional space of co-directions in real projective space P^n is described by viewing all quantities in the foregoing discussion as being real. Especially, G_0 of 2.11 is the connected centerless group $PSL(n+1; \mathbf{R})$ consisting of real contact automorphisms.

4. HIGHER SPHERE GEOMETRY

4.1 In complex Euclidean space E^n , the equation

$$x_1'^2 + \dots + x_n'^2 - 2a_1 x_1' - \dots - 2a_n x_n' + C = 0$$

describes a sphere with center (a_1, \dots, a_n) and complex radius r given by

$$r^2 = a_1^2 + \dots + a_n^2 - C.$$

When $r \neq 0$, the two choices of sign for r is said to give two "orientations" to the sphere. Thus, the $n+2$ coordinates a_1, \dots, a_n, r, C , which are related by

$$a_1^2 + \dots + a_n^2 - r^2 - C = 0,$$

describe the space of oriented spheres in E^n [6, §25].

Introduce homogeneous coordinates by

$$a_i = \frac{\alpha_i}{v}, \quad r = \frac{\lambda}{v}, \quad C = \frac{\mu}{v},$$

$i = 1, 2, \dots, n$. Then the oriented spheres of E^n correspond to certain points of the quadric Ψ^{n+1} in P^{n+2} described by

$$\alpha_1^2 + \dots + \alpha_n^2 - \lambda^2 - \mu v = 0.$$

The sphere corresponding to the point $(\alpha_1, \dots, \alpha_n, \lambda, \mu, v)$ of Ψ^{n+1} is

$$v(x_1'^2 + \dots + x_n'^2) - 2\alpha_1 x_1' - \dots - 2\alpha_n x_n' + \mu = 0.$$

Ordinary spheres have finite nonzero radius r , so $v \neq 0$. For $v = 0$, we obtain oriented hyperplanes. For $\lambda = 0$, we obtain point spheres or hyperplanes with isotropic hyperplane coordinate vector; these carry no

orientation. If we include these special cases as spheres of E^n , then Ψ^{n+1} is the space of all oriented spheres in E^n .

Two spheres in E^n with centers (a_1, \dots, a_n) , (a'_1, \dots, a'_n) and radii r, r' respectively are tangent, orientations taken into account, if

$$(a_1 - a'_1)^2 + \dots + (a_n - a'_n)^2 = (r - r')^2.$$

Use $a_1^2 + \dots + a_n^2 = r^2 + C$ for both spheres to obtain the condition for tangency as

$$2a_1 a'_1 + \dots + 2a_n a'_n - 2rr' - C - C' = 0$$

or, in homogeneous coordinates,

$$2\alpha_1 \alpha'_1 + \dots + 2\alpha_n \alpha'_n - 2\lambda \lambda' - \mu \nu' - \nu \mu' = 0.$$

Hence, two spheres of E^n are tangent when their corresponding points in Ψ^{n+1} are conjugate, that is, the line joining these points lies entirely in Ψ^{n+1} [6, §25].

4.2. A pencil of mutually tangent spheres in E^n corresponds to a line in Ψ^{n+1} . This pencil of spheres determines an "oriented complex co-direction" in E^n since it contains a point sphere and an incident oriented hyperplane. Corresponding to the hyperplane

$$x'_n - x_n = p_1 (x'_1 - x_1) + \dots + p_{n-1} (x'_{n-1} - x_{n-1})$$

at the point (x_1, \dots, x_n) is the line

$$\begin{bmatrix} \alpha_1 \\ \vdots \\ \alpha_{n-1} \\ \alpha_n \\ \hline \lambda \\ \mu \\ \nu \end{bmatrix} = \begin{bmatrix} x_1 \\ \vdots \\ x_{n-1} \\ x_n \\ \hline 0 \\ xx \\ 1 \end{bmatrix} + t \begin{bmatrix} -p_1 \\ \vdots \\ -p_{n-1} \\ 1 \\ \hline -\sqrt{pp+1} \\ 2(x_n - px) \\ 0 \end{bmatrix}$$

of Ψ^{n+1} , where

$$xx = \sum_{i=1}^n x_i^2, \quad px = \sum_{i=1}^{n-1} p_i x_i, \quad pp = \sum_{i=1}^{n-1} p_i^2;$$

this is the pencil of spheres

$$\sum_{i=1}^{n-1} (x'_i - x_i + t p_i)^2 + (x'_n - x_n - t)^2 = t^2 \left(\sum_{i=1}^{n-1} p_i^2 + 1 \right)$$

passing through (x_1, \dots, x_n) and having their centers on the line normal to the hyperplane at this point.

For later calculations it will be convenient to replace $-p, \dots, -p_{n-1}, 1$ by homogeneous u_1, \dots, u_{n-1}, u_n . The line in Ψ^{n+1} corresponding to the hyperplane

$$u_1(x'_1 - x_1) + \dots + u_n(x'_n - x_n) = 0$$

at the point (x_1, \dots, x_n) is then

$$\begin{array}{|c|} \hline \alpha_1 \\ \hline \vdots \\ \hline \alpha_{n-1} \\ \hline \alpha_n \\ \hline \lambda \\ \hline \mu \\ \hline \nu \\ \hline \end{array} = \begin{array}{|c|} \hline x_1 \\ \hline \vdots \\ \hline x_{n-1} \\ \hline x_n \\ \hline 0 \\ \hline xx \\ \hline 1 \\ \hline \end{array} + t \begin{array}{|c|} \hline u_1 \\ \hline \vdots \\ \hline u_{n-1} \\ \hline u_n \\ \hline -\sqrt{uu} \\ \hline 2ux \\ \hline 0 \\ \hline \end{array},$$

where

$$xx = \sum_{i=1}^n x_i^2, \quad ux = \sum_{i=1}^n u_i x_i, \quad uu = \sum_{i=1}^n u_i^2.$$

Any convenient condition may be imposed on uu .

4.3. The contact structure on the $(2n-1)$ -dimensional space of lines in Ψ^{n+1} , that is, the space of oriented co-directions in complex Euclidean space E^n , is obtained when the construction of 2.10. is carried out for the simple complex Lie algebra of type B_l or D_l , $l \geq 2$ and $l \geq 3$ respectively. However, it will be simpler to identify quantities geometrically if

- G/P = space of lines in Ψ^{n+1}
- = space of pencils of mutually tangent oriented spheres in E^n
- = space of oriented co-directions in complex E^n .

The Lie algebra \mathfrak{g} of G consists of $(n+3)$ by $(n+3)$ matrices X for which ${}^tXA + AX = 0$. The matrices of \mathfrak{g} are of the form

$$\left[\begin{array}{ccc|ccc} & & & b_1 & c_1 & d_1 \\ & & & \vdots & \vdots & \vdots \\ n \text{ by } n \text{ skew-} & & & b_{n-1} & c_{n-1} & d_{n-1} \\ \text{symmetric} & & & b_n & c_n & d_n \\ \hline b_1 \dots b_{n-1} & b_n & 0 & c & d \\ 2d_1 \dots 2d_{n-1} & 2d_n & -2d & e & 0 \\ 2c_1 \dots 2c_{n-1} & 2c_n & -2c & 0 & -e \end{array} \right].$$

P consists of those elements of G which send the subspace of \mathbb{C}^{n+3} spanned by ${}^t(0, \dots, 0, 0 \mid 0, 0, 1)$ and ${}^t(0, \dots, 0, 1 \mid -1, 0, 0)$

into itself; the Lie algebra \mathfrak{p} of P consists of those elements of \mathfrak{g} which do the same. Hence, the matrices of \mathfrak{p} are of the form

$$\left[\begin{array}{ccc|ccc} & & & b_1 & b_1 & c_1 & 0 \\ & & & \vdots & \vdots & \vdots & \vdots \\ (n-1) \text{ by } (n-1) & & & b_{n-1} & b_{n-1} & c_{n-1} & 0 \\ \text{skew-symmetric} & & & & & & \\ \hline -b_1 \dots -b_{n-1} & 0 & b_n & c_n & -d \\ b_1 \dots b_{n-1} & b_n & 0 & c & d \\ 0 \dots 0 & -2d & -2d & e & 0 \\ 2c_1 \dots 2c_{n-1} & 2c_n & -2c & 0 & -e \end{array} \right].$$

Note that \mathfrak{g} and \mathfrak{p} have dimensions $\frac{1}{2}(n+3)(n+2)$ and $\frac{1}{2}(n-1)(n-2) + 2n + 3 = \frac{1}{2}(n+3)(n+2) - 2n + 1$, respectively, in agreement with G/P having dimension $2n-1$.

4.4. For $n \geq 2$, set $n+3 = 2l+1$ or $2l$ according as n is even or odd. \mathfrak{g} is of type B_l or D_l , $l \geq 2$ and $l \geq 3$ respectively.

For Cartan subalgebra \mathfrak{h} of \mathfrak{g} take matrices of the form

$$H = \text{diag} \left[\begin{array}{c} 0 \\ \left[\begin{array}{cc} 0 & h_1 \\ -h_1 & 0 \end{array} \right], \dots, \left[\begin{array}{cc} 0 & h_{l-2} \\ -h_{l-2} & 0 \end{array} \right], \left[\begin{array}{cccc} 0 & h_{l-1} & 0 & 0 \\ h_{l-1} & 0 & 0 & 0 \\ 0 & 0 & h_l & 0 \\ 0 & 0 & 0 & -h_l \end{array} \right] \end{array} \right];$$

the first row and column occur only in case B_l , it is suppressed for case D_l . The Killing form of \mathfrak{g} is $\langle X, Y \rangle = (n+1) \text{tr}(XY)$, but we replace this with $\langle X, Y \rangle = \frac{1}{2} \text{tr}(XY)$ for convenience.

Let W in \mathfrak{p} be

$$W = \left[\begin{array}{ccc|ccc} 0 & 0 & 0 & & & \\ \hline 0 & 0 & 0 & & & \\ & & 0 & 0 & -\frac{1}{2} & 0 \\ \hline 0 & 0 & 0 & 0 & \frac{1}{2} & 0 \\ & & 0 & 0 & 0 & 0 \\ \hline & & -1 & -1 & 0 & 0 \end{array} \right]$$

For H in \mathfrak{h} we have $[H, W] = -(h_{l-1} + h_l)W$, so $\rho(H) = -(h_{l-1} + h_l)$ is a root of \mathfrak{g} with respect to \mathfrak{h} and $W = E_\rho$ is the corresponding root vector.

For X in \mathfrak{g} as described in 4.3, direct calculation shows $[X, W] = 0$ implies X is in \mathfrak{p} and $b_n + e = 0$; thus the centralizer of W in \mathfrak{g} consists of those elements of \mathfrak{p} with $b_n + e = 0$. For X in \mathfrak{p} now, the same calculation gives $[X, W] = -(b_n + e)W$, so $[X, W] = \rho(X)W$ with ρ extended to \mathfrak{p} by $\rho(X) = -(b_n + e)$. Finally, W is orthogonal to \mathfrak{p} with respect to the Killing form. Hence, (a', c', b') of 2.7 are satisfied, and W is the element of \mathfrak{g} giving the contact structure on G/P .

The origin of the element W is not immediately evident. It was obtained by determining the maximal root and corresponding root vector for Lie algebras of type B_l and D_l when the quadratic form is

$$\xi_0^2 + 2\xi_1\xi_{l+1} + \dots + 2\xi_l\xi_{2l}$$

and then passing to the form

$$\alpha_1^2 + \dots + \alpha_n^2 - \lambda^2 - \mu\nu$$

by conjugating by the element of $PSL(n+3; \mathbf{C})$ which corresponds to the "line-sphere transformation". This will be described further in the next section.

4.5. Let \mathfrak{m} be the $(2n-1)$ -dimensional supplement to \mathfrak{p} in \mathfrak{g} consisting of matrices of the form

$$\left[\begin{array}{cccc|ccc} & & & -b_1 & b_1 & 0 & d_1 \\ & & & \vdots & \vdots & \vdots & \vdots \\ & 0 & & -b_{n-1} & b_{n-1} & 0 & d_{n-1} \\ & & & \vdots & \vdots & \vdots & \vdots \\ b_1 & \dots & b_{n-1} & 0 & 0 & 0 & d_n \\ \hline b_1 & \dots & b_{n-1} & 0 & 0 & 0 & d_n \\ 2d_1 & \dots & 2d_{n-1} & 2d_n & -2d_n & 0 & 0 \\ 0 & \dots & 0 & 0 & 0 & 0 & 0 \end{array} \right],$$

$$(\exp X) \cdot \begin{bmatrix} 0 \\ \vdots \\ 0 \\ 1 \\ \hline -1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} -2b_1 \\ \vdots \\ -2b_{n-1} \\ 1 - bb \\ \hline -1 - bb \\ 4d_n - 2bd \\ 0 \end{bmatrix}$$

On this line we can identify the point sphere when $\lambda = 0$, giving

$$\begin{bmatrix} d_1 \\ \vdots \\ d_{n-1} \\ d_n + \frac{1}{2} bd \\ \hline d_n + \frac{1}{2} bd \\ dd \\ 1 \end{bmatrix} + \frac{d_n + \frac{1}{2} bd}{1 + bb} \begin{bmatrix} -2b_1 \\ \vdots \\ -2b_{n-1} \\ 1 - bb \\ \hline -1 - bb \\ 4d_n - 2bd \\ 0 \end{bmatrix} = \begin{bmatrix} x_1 \\ \vdots \\ x_{n-1} \\ x_n \\ \hline 0 \\ xx \\ 1 \end{bmatrix},$$

and the incident oriented hyperplane when $v = 0$, giving

$$\begin{bmatrix} -2b_1 \\ \vdots \\ -2b_{n-1} \\ 1 - bb \\ \hline -1 - bb \\ 4d_n - 2bd \\ 0 \end{bmatrix} = \begin{bmatrix} u_1 \\ \vdots \\ u_{n-1} \\ u_n \\ \hline -\sqrt{uu} \\ 2ux \\ 0 \end{bmatrix} .$$

These equations will be satisfied if we impose the condition $\sqrt{uu} = 1 + bb$ on uu , or

$$u_i = -2b_i, \quad u_n = 1 - bb,$$

$i = 1, 2, \dots, n-1$, and then set

$$b_i = -\frac{1}{2} u_i,$$

$$d_i = x_i - \frac{1}{2} u_i x_n, \quad i = 1, 2, \dots, n-1$$

$$d_n = \frac{1}{4} \sum_{i=1}^{n-1} u_i x_i + \frac{1}{2} x_n .$$

Thus, this choice of X establishes the classically identifiable coordinates $x_1, \dots, x_n, u_1, \dots, u_n$ on G/P as in 2.12 and 4.2.

4.6. From 2.12, the form ω on G/P is obtained as

$$\omega = \langle W, (\exp X)^{-1} d(\exp X) \rangle$$

with $(\exp X)^{-1} d(\exp X) = dX - \frac{1}{2} [X, dX] .$

Take X as in 4.5 and let the entries of dX be denoted as those of X with primes affixed. Then

$$(\exp X)^{-1} d (\exp X) = \left[\begin{array}{cccc|cccc} & & -b_1' & & b_1' & 0 & & d_1' \\ & 0 & \vdots & & \vdots & \vdots & & \vdots \\ & & -b_{n-1}' & & b_{n-1}' & 0 & & d_{n-1}' \\ b_1' \dots b_{n-1}' & & 0 & & 0 & 0 & d_n' - \frac{1}{2}c & \\ \hline b_1' \dots b_{n-1}' & & 0 & & 0 & 0 & d_n' - \frac{1}{2}c & \\ 2d_1' \dots 2d_{n-1}' & 2d_n' - c & & -2d_n' + c & 0 & & & 0 \\ 0 \dots 0 & 0 & & 0 & 0 & 0 & & 0 \end{array} \right]$$

where

$$c = \sum_{i=1}^{n-1} (b_i d_i' - d_i b_i'),$$

and consequently, from the definition of W in 4.4, $\omega = c - 2d_n'$. Using the expressions in 4.5 for $b_1, \dots, b_{n-1}, d_1, \dots, d_n$ in terms of $x_1, \dots, x_n, u_1, \dots, u_n$, we obtain

$$\omega = - \sum_{i=1}^{n-1} u_i dx_i - \left[1 - \frac{1}{4} \sum_{i=1}^{n-1} u_i^2 \right] dx_n$$

or, since $1 - \frac{1}{4} \sum_{i=1}^{n-1} u_i^2 = u_n$,

$$\omega = - (u_1 dx_1 + \dots + u_n dx_n).$$

This identifies the contact structure with the classical one as in 2.12 and 4.2.

4.7. The real contact structure on the $(2n-1)$ -dimensional space of oriented co-direction in real Euclidean space E^n is described by viewing all quantities in the foregoing discussion as being real. Especially, G_0 of 2.11 is the two-component centerless group $PSO(A; \mathbf{R})$ consisting of real contact automorphisms.