

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 25 (1979)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON LIE'S HIGHER SPHERE GEOMETRY
Autor: Fillmore, Jay P.
Kapitel: 5. The line-sphere transformation
DOI: <https://doi.org/10.5169/seals-50373>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 13.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

5. THE LINE-SPHERE TRANSFORMATION

The homogeneous contact manifold of co-directions in complex projective space P^3 , obtained from the simple complex Lie algebra of type A_3 , must coincide with that of oriented co-directions in complex Euclidean space E^3 , obtained from the algebra of type D_3 , in view of the isomorphisms $A_3 \simeq D_3$. To exhibit this explicitly, we introduce a third homogeneous contact manifold in terms of which both of these can be conveniently described, namely, the space of lines in the quadric Ω^4 in P^5 of Section 1.

5.1. We carry out the construction of 2.10 for the simple complex Lie algebras of type B_l and D_l , making the restriction to type D_3 later.

Let $\mathfrak{g} = \mathfrak{o}(A; \mathbb{C})$, complex square matrices X for which ${}^tXA + AX = 0$, where

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1_l \\ 0 & 1_l & 0 \end{bmatrix} \quad \text{in case } B_l$$

or

$$A = \begin{bmatrix} 0 & 1_l \\ 1_l & 0 \end{bmatrix} \quad \text{in case } D_l,$$

that is, the quadratic form defining \mathfrak{g} is

$$\xi_0^2 + 2\xi_1\xi_{l+1} + \dots + 2\xi_l\xi_{2l}$$

or

$$2\xi_1\xi_{l+1} + \dots + 2\xi_l\xi_{2l}$$

respectively [4, (16.3) and (16.4)].

We exhibit the details of the construction for the case of D_l . For B_l one need only carry along an additional initial row and column in the matrices, as well as the corresponding roots; the conclusions are the same.

Thus \mathfrak{g} consists of $2l$ by $2l$ matrices of the form

$$\begin{bmatrix} X_1 & X_2 \\ X_3 & -{}^tX_1 \end{bmatrix},$$

where X_1 is l by l and arbitrary and X_2 and X_3 are l by l and skew-symmetric. For Cartan subalgebra \mathfrak{h} of \mathfrak{g} take diagonal matrices H of the form

$$H = \text{diag} (h_1, \dots, h_l \mid -h_1, \dots, -h_l).$$

Let $\delta_i, i = 1, 2, \dots, l$ be the linear function on \mathfrak{h} which assigns h_i to $H: \delta_i(H) = h_i$. The roots of \mathfrak{g} with respect to \mathfrak{h} are

$$\begin{aligned} \pm \delta_i \pm \delta_j \quad i, j = 1, 2, \dots, l \\ \text{and } i \neq j \end{aligned}$$

and the root vector E_α corresponding to the root α is

$$\begin{aligned} E_{\delta_i - \delta_j} &= \begin{bmatrix} E_{ij} & 0 \\ 0 & -E_{ji} \end{bmatrix}, \quad i \neq j, \\ E_{\delta_i + \delta_j} &= \begin{bmatrix} 0 & E_{ij} - E_{ji} \\ 0 & 0 \end{bmatrix}, \quad i < j, \\ E_{-\delta_i - \delta_j} &= \begin{bmatrix} 0 & 0 \\ E_{ji} - E_{ij} & 0 \end{bmatrix}, \quad i < j, \end{aligned}$$

where E_{ij} is the l by l matrix with 1 in the i^{th} row and j^{th} column and 0s elsewhere [4, (16.3)]. A system of simple roots is

$$\delta_1 - \delta_2, \delta_2 - \delta_3, \dots, \delta_{l-1} - \delta_l, \quad \text{and} \quad -\delta_1 - \delta_2,$$

(this is not the same choice as in [4, (16.3)]), for which the maximal root is

$$\rho = -\delta_{l-1} - \delta_l,$$

[4, App., Table E]. The Killing form of \mathfrak{g} is $\langle X, Y \rangle = (2l-2) \text{tr}(XY)$, but we replace this with $\langle X, Y \rangle = \frac{1}{2} \text{tr}(XY)$ for convenience. Then the H_α in \mathfrak{h} are given by

$$H_{\pm\delta_i \pm \delta_j} = \text{diag}(0, \dots, 0, \pm 1, 0, \dots, 0, \pm 1, 0, \dots, 0 \mid \text{---}),$$

where the ± 1 s occur in the i^{th} and j^{th} entries and the second l entries are the negatives of the first l entries. Especially,

$$H_\rho = \text{diag}(0, \dots, 0, -1, -1 \mid 0, \dots, 0, 1, 1).$$

It is now straightforward to determine for which roots α we have $\langle H_\rho, H_\alpha \rangle \geq 0$ and find that \mathfrak{p} in (i) of 2.9 consists of matrices of the form

$$\left[\begin{array}{c|ccc} & \text{arbitrary} & 0 & 0 \\ & (l-2) \text{ by } (l-2) & \vdots & \vdots \\ & \text{skew-} & & \\ & \text{symmetric} & 0 & 0 \\ & 0 \text{ -----} & 0 & 0 & 0 \\ & 0 \text{ -----} & 0 & 0 & 0 \\ \hline & * \text{ -----} & * & 0 & 0 \\ & \vdots & \vdots & \vdots & \vdots \\ & \text{arbitrary} & & 0 & 0 \\ & l \text{ by } l & & * & * \\ & \text{skew-} & & & \\ & \text{symmetric} & & * & * \\ & * \text{ -----} & * & * & * \end{array} \right],$$

where the starred entries are arbitrary.

5.2. The connected centerless simple group $G = PSO(A; \mathbb{C})$ is transitive on the lines of the quadric Ω^{2l-2}

$$\xi_1 \xi_{l+1} + \dots + \xi_l \xi_{2l} = 0$$

in P^{2l-1} by Witt's theorem. The Lie algebra of the isotropy subgroup of the line l_0 joining

$${}^t(0, \dots, 0, 1, 0) \quad \text{and} \quad {}^t(0, \dots, 0, 0, 1)$$

is \mathfrak{p} . Hence

$$G/P = \text{space of lines in } \Omega^{2l-2}.$$

The element $W = E_\rho$ of \mathfrak{p} giving the contact structure on G/P , as in 2.7, is

$$W = \left[\begin{array}{c|ccc} & 0 & & 0 \\ \hline & & & \\ & 0 & & \\ & & & 0 \\ & 0 & -1 & \\ & 1 & 0 & \end{array} \right]$$

In general, the construction of 2.10 gives the $(2n-1)$ -dimensional homogeneous contact manifold of lines in the quadric Ω^{n+1} in P^{n+2} , where Ω^{n+1} is

$$\xi_0^2 + 2\xi_1 \xi_{l+1} + \dots + 2\xi_l \xi_{2l} = 0$$

in case B_l when n is even, $n+3 = 2l+1$, and Ω^{n+1} is Ω^{2l-2} above in case D_l when n is odd, $n+3 = 2l$; $n \geq 2$.

The real contact structure on the $(2n-1)$ dimensional space of lines of Ω^{n+1} in real projective space P^{n+2} is described by viewing all quantities in the foregoing discussion as being real. Especially, G_0 of 2.11 is the one- or two- component centerless group $PSO(A; \mathbf{R})$ consisting of real contact automorphisms.

5.3. The line joining $x = {}^t(x_0, x_1, x_2, x_3)$ and $y = {}^t(y_0, y_1, y_2, y_3)$ in complex projective space P^3 has Plücker coordinates $p_{ij} = x_i y_j - x_j y_i$. These coordinates are the coefficients of the bivector $x \wedge y$ with respect to the basis

$$e_1 \wedge e_2, e_3 \wedge e_1, e_2 \wedge e_3, e_0 \wedge e_3, e_0 \wedge e_2, e_0 \wedge e_1,$$

where $e_0 = {}^t(1, 0, 0, 0), \dots, e_3 = {}^t(0, 0, 0, 1)$, and satisfy

$$p_{01} p_{23} + p_{02} p_{31} + p_{03} p_{12} = 0,$$

[6, §69]. If we set

$$\begin{aligned} \xi_1 &= p_{12}, & \xi_2 &= p_{31}, & \xi_3 &= p_{23}, \\ \xi_4 &= p_{03}, & \xi_5 &= p_{02}, & \xi_6 &= p_{01}, \end{aligned}$$

we have that the lines of P^3 correspond to the points of the quadric Ω^4

$$\xi_1 \xi_4 + \xi_2 \xi_5 + \xi_3 \xi_6 = 0$$

in P^5 . Two lines of P^3 intersect exactly when their corresponding points on Ω^4 are conjugate, that is, the line joining these points lies entirely in Ω^4 .

To a point x in P^3 we associate all lines of P^3 incident with x and hence a plane lying in Ω^4 . To a plane u in P^3 we associated all lines of P^3 lying in u and hence a plane lying in Ω^4 . These two families of planes doubly rule Ω^4 . To a surface element or co-direction in P^3 , that is, a point x and incident plane u , is then associated all lines of P^3 lying in u and incident with x . In Ω^4 this corresponds to the intersection of the planes corresponding to x and u and is a line. Hence, the 5-dimensional spaces of co-directions in P^3 and lines in Ω^4 correspond naturally.

Note that the co-direction in P^3 consisting of the point $x_0 = {}^t(1, 0, 0, 0)$ and the incident plane $u_0: x_3 = 0$ in 3.2 corresponds to the line l_0 of Ω^4 joining the points ${}^t(0, 0, 0, 0, 1, 0)$ and ${}^t(0, 0, 0, 0, 0, 1)$ in 5.2. For, to the co-direction (x_0, u_0) is associated all lines of P^3 joining x_0 and a point $y = {}^t(y_0, y_1, y_2, 0)$ of u_0 ; such a line has Plücker coordinates

$$\begin{aligned} \xi_1 &= 0, & \xi_2 &= 0, & \xi_3 &= 0, \\ \xi_4 &= 0, & \xi_5 &= y_2, & \xi_6 &= y_1, \end{aligned}$$

and corresponds to a point of Ω^4 lying on l_0 .

The projectivity g in $PSL(4; \mathbf{C})$ permutes the lines of P^3 by $x \wedge y \rightarrow gx \wedge gy$, a projectivity of P^5 which preserves Ω^4 . In this way one obtains the isomorphism $A_3 \simeq D_3$:

$$PSL(4; \mathbf{C}) \simeq PSO(A; \mathbf{C}), \quad A = \begin{bmatrix} 0 & 1_3 \\ 1_3 & 0 \end{bmatrix},$$

[4, (25.8.4')]. The spaces of co-directions in P^3 and lines in Ω^4 are homogeneous under $PSL(4; \mathbf{C})$ and $PSO(A; \mathbf{C})$ respectively; hence the correspondence between these spaces is as homogeneous spaces. In fact, since (x_0, u_0) and l_0 correspond, their isotropy subgroups, as described in 3.2 and 5.2, correspond under the isomorphism.

From the isomorphism of the groups, we obtain the isomorphism of the Lie algebras $\mathfrak{sl}(4; \mathbf{C}) \simeq \mathfrak{o}(A; \mathbf{C})$, where X in $\mathfrak{sl}(4; \mathbf{C})$ is sent to the linear transformation $x \wedge y \rightarrow (Xx) \wedge y + x \wedge (Xy)$ in $\mathfrak{o}(A; \mathbf{C})$. With $X = (a_{ij})$, $i, j = 0, 1, 2, 3$, the matrix of this transformation with respect to the basis $e_i \wedge e_j$ is

$$\begin{bmatrix} a_{11} + a_{22} & -a_{23} & -a_{13} & 0 & a_{10} & -a_{20} \\ -a_{32} & a_{11} + a_{33} & -a_{12} & -a_{10} & 0 & a_{30} \\ -a_{31} & -a_{21} & a_{22} + a_{33} & a_{20} & -a_{30} & 0 \\ \hline 0 & -a_{01} & a_{02} & a_{00} + a_{33} & a_{32} & a_{31} \\ a_{01} & 0 & -a_{03} & a_{23} & a_{00} + a_{22} & a_{21} \\ -a_{02} & a_{03} & 0 & a_{13} & a_{12} & a_{00} + a_{11} \end{bmatrix};$$

Thus,

$$G = PSO(A; C), A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1_l \\ 0 & 1_l & 0 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} 0 & 1_l \\ 1_l & 0 \end{bmatrix}$$

and

$$G' = PSO(A'; C), A' = \begin{bmatrix} 2 \cdot 1_n & & & 0 \\ \hline & & & \\ & & -2 & 0 & 0 \\ & 0 & 0 & 0 & -1 \\ & & & 0 & -1 & 0 \end{bmatrix}.$$

These are groups of projectivities preserving Ω^{n+1} and Ψ^{n+1} , respectively, in P^{n+2} .

In case n is odd, the transformation which we consider is

$$\begin{aligned} \xi_1 &= \alpha_1 + \sqrt{-1} \alpha_2 & \xi_{l+1} &= \alpha_1 - \sqrt{-1} \alpha_2 \\ \xi_2 &= \alpha_3 + \sqrt{-1} \alpha_4 & \xi_{l+2} &= \alpha_3 - \sqrt{-1} \alpha_4 \\ & \vdots & & \vdots \\ \xi_{l-2} &= \alpha_{n-2} + \sqrt{-1} \alpha_{n-1} & \xi_{2l-2} &= \alpha_{n-2} - \sqrt{-1} \alpha_{n-1} \\ \xi_{l-1} &= \alpha_n + \lambda & \xi_{2l-1} &= \alpha_n - \lambda \\ \xi_l &= \mu & \xi_{2l} &= -v. \end{aligned}$$

This is a projectivity of P^{n+2} which sends the quadric Ψ^{n+1}

$$\alpha_1^2 + \dots + \alpha_n^2 - \lambda^2 - \mu v = 0$$

into the quadric Ω^{n+1}

$$2\xi_1\xi_{l+1} + \dots + 2\xi_l\xi_{2l} = 0.$$

In case n is even, the first equation of the transformation is $\xi_0 = \sqrt{2} \alpha_1$ and the remaining ones are like the above.

As before, we exhibit the details of the calculations for the case of n odd. For n even one need only carry along an additional initial row and column in the matrices; the conclusions are unchanged.

The matrix T of the transformation is

$$T = \begin{bmatrix} B & & & 0 & & \\ & 1 & 1 & 0 & 0 & \\ & 0 & 0 & 1 & 0 & \\ \bar{B} & & & 0 & & \\ & 1 & -1 & 0 & 0 & \\ 0 & & & & & \\ & 0 & 0 & 0 & -1 & \end{bmatrix},$$

where B is the $(l-2)$ by $(2l-4)$ matrix

$$B = \begin{bmatrix} 1 & \sqrt{-1} & & & & \\ & & 1 & \sqrt{-1} & & 0 \\ & & & & \ddots & \\ & 0 & & & & \\ & & & & & \\ & & & & & \\ & & & & & 1 & \sqrt{-1} \end{bmatrix}$$

and \bar{B} is its complex conjugate; T has inverse

$$T^{-1} = \frac{1}{2} \begin{bmatrix} {}^t\bar{B} & 0 & {}^tB & 0 & & \\ & 1 & 0 & & 1 & 0 \\ & 0 & 1 & 0 & 0 & -1 \\ & & 0 & 2 & & 0 \\ & & 0 & 0 & & 0 \\ & & & & & 0 & -2 \end{bmatrix}.$$

By direct calculation we ascertain the following:

- (1) $A' = {}^tTAT$ and hence $G' = T^{-1}GT$. G and G' are conjugate, but do not coincide, in $PSL(n+3; \mathbb{C})$. As a consequence, $g' = T^{-1}gT$.
- (2) $l'_0 = T^{-1}l_0$; the line l_0 in Ω^{n+1} joining

$${}^t(0, \dots, 0, 1, 0) \quad \text{and} \quad {}^t(0, \dots, 0, 0, 1)$$

is sent to the line l'_0 of Ψ^{n+1} joining

$${}^t(0, \dots, 0, 0 \mid 0, 0, 1) \quad \text{and} \quad {}^t(0, \dots, 0, 1 \mid -1, 0, 0).$$

Hence their isotropy subgroups, as in 5.2 and 4.3 are conjugate: $P' = T^{-1} P T$. As a consequence, $\mathfrak{p}' = T^{-1} \mathfrak{p} T$.

(3) The Cartan subalgebras of \mathfrak{g} and \mathfrak{g}' in 5.1 and 4.4 are conjugate: $\mathfrak{h}' = T^{-1} \mathfrak{h} T$. In fact, for

$$H = \text{diag}(h_1, \dots, h_l \mid -h_1, \dots, -h_l)$$

in \mathfrak{h} , we have

$$T^{-1} H T = \text{diag} \left[\begin{array}{cc} 0 & \sqrt{-1} h_1 \\ -\sqrt{-1} h_1 & 0 \end{array} \right], \dots, \left[\begin{array}{cc} 0 & \sqrt{-1} h_{l-2} \\ -\sqrt{-1} h_{l-2} & 0 \end{array} \right],$$

$$\left[\begin{array}{cccc} 0 & h_{l-1} & 0 & 0 \\ h_{l-1} & 0 & 0 & 0 \\ 0 & 0 & h_l & 0 \\ 0 & 0 & 0 & -h_l \end{array} \right]$$

in \mathfrak{h}' .

4) The elements W and W' of the Lie algebras which give the contact structures on G/P and G'/P' , as in 5.2 and 4.4, are conjugate: $W' = T^{-1} W T$. We conclude:

The $(2n-1)$ -dimensional manifolds of lines in Ω^{n+1} and lines in Ψ^{n+1} are isomorphic as algebraic homogeneous contact manifolds. The isomorphism is a consequence of the projectivity T carrying Ψ^{n+1} into Ω^{n+1} . T sends lines of Ψ^{n+1} into lines of Ω^{n+1} and is a contact transformation.

5.5. $G_0 = PSO(A; \mathbf{R})$ is a real form of G ; it consists of the elements of G fixed under the conjugation $g \rightarrow \bar{g}$ of G , complex conjugation of the matrix entries of g . The Cartan subalgebra \mathfrak{h} of \mathfrak{g} , as in 5.1, is stable and the maximal root $\rho = -\delta_{l-1} - \delta_l$ is real. With $P_0 = G_0 \cap P$, we obtain from 2.11 the real contact manifold

$$G_0/P_0 = \text{space of lines in } \Omega^{n+1} \text{ in real } P^{n+2},$$

a real form of G/P ; cf. 5.2. The same remarks apply to the real form $G'_0 = PSO(A'; \mathbf{R})$ of G' for the conjugation $g' \rightarrow \bar{g}'$. With $P'_0 = G'_0 \cap P'$, we obtain the real contact manifold

$$\begin{aligned} G'_0/P'_0 &= \text{space of lines in } \Psi^{n+1} \text{ in real } P^{n+2} \\ &= \text{space of pencils of mutually tangent oriented spheres in real } E^n \\ &= \text{space of oriented co-directions in real } E^n, \end{aligned}$$

a real form of G/P ; cf. 4.7.

Since $G' = T^{-1} G T$, we can exhibit G'_0/P'_0 , as well as G_0/P_0 , as a real form of the complex contact manifold G/P . $T G'_0 P^{-1}$ is the real form of $G = T G' T^{-1}$ consisting of the elements fixed under the conjugation obtained by transporting the conjugation $g' \rightarrow \bar{g}'$ of G' to G , namely

$$g \rightarrow T \overline{(T^{-1} g T)} T^{-1} = S^{-1} \bar{g} S$$

where $S = \bar{T} T^{-1}$. In the case of n odd,

$$S = \left[\begin{array}{cc|cc} 0 & 0 & 1_{l-2} & 0 \\ 0 & 1_2 & 0 & 0 \\ \hline 1_{l-2} & 0 & 0 & 0 \\ 0 & 0 & 0 & 1_2 \end{array} \right];$$

in the case of n even, S has an additional initial row and column with a 1 in their common first entry and 0s elsewhere. $S A S = A$ and $S^2 = 1_{n+3}$, so the complex conjugation $\xi \rightarrow S^{-1} \bar{\xi}$ preserves the quadric Ω^{n+1} . A point or line of Ω^{n+1} is fixed under this conjugation exactly if it is the image under T of a real point or line of Ψ^{n+1} . The latter constitute the orbit on Ω^{n+1} of $T G'_0 T^{-1}$. The isotropy subgroup in $T G'_0 T^{-1}$ of the line l_0 of Ω^{n+1} is $T G'_0 T^{-1} \cap P = T P'_0 T^{-1}$. Furthermore, the Cartan subalgebra \mathfrak{h} of \mathfrak{g} in 5.1 is stable under the conjugation $X \rightarrow S^{-1} \bar{X} S$ of \mathfrak{g} ; in fact, for

$$H = \text{diag} (h_1, \dots, h_l \mid -h_1, \dots, -h_l)$$

in \mathfrak{h} , we have

$$S^{-1} \bar{H} S = \text{diag} (-\bar{h}_1, \dots, -\bar{h}_{l-2}, \bar{h}_{l-1}, \bar{h}_l \mid \bar{h}_1, \dots, \bar{h}_{l-2}, -\bar{h}_{l-1}, -\bar{h}_l),$$

in case of n odd; the maximal root $\rho = -\delta_{l-1} - \delta_l$ is real, $\overline{\rho(S^{-1}\bar{H}S)} = \rho(H)$. Hence, the contact structure on TG'_0T^{-1}/TP'_0T^{-1} is that obtained from G/P by 2.11. We conclude:

G_0/P_0 and TG'_0T^{-1}/TP'_0T^{-1} , the latter isomorphic to G'_0/P'_0 , are two real forms of the complex contact manifold G/P .

5.6. We observed in 5.3 that the space of co-directions in complex projective space P^3 , by means of Plücker's line geometry, is isomorphic to the space of lines in the quadric Ω^4 in complex P^5 , and that this isomorphism makes real line geometry correspond to a real form of Ω^4 . We found in 5.4 and 5.5 that the space of oriented co-directions in complex Euclidean space E^3 of Lie's higher sphere geometry, which is the space of lines in the quadric Ψ^4 in complex P^5 , is isomorphic to the space of lines in the quadric Ω^4 also, and that this isomorphism makes real sphere geometry correspond to a second real form of Ω^4 . That is, real line geometry and real sphere geometry are two distinct real forms of complex line geometry. The line-sphere transformation establishes the isomorphism of the spaces of lines in Ψ^4 and lines in Ω^4 . The former places real sphere geometry in the foreground, the latter, real line geometry.

5.7. The isomorphism of 5.3 may be used to describe sphere geometry in terms of co-directions in complex P^3 . Real sphere geometry then leads to the real form $PSU(2,2)$ of $PSL(4; \mathbb{C})$.

REFERENCES

- [1] BOOTHBY, W. M. Homogeneous complex contact manifolds. *Proc. Sympos. Pure Math., Vol. III*, pp. 144-154, Amer. Math. Soc., Providence, R. I., 1961.
- [2] — A note on homogeneous complex contact manifolds. *Proc. Amer. Math. Soc.* 13 (1961), pp. 276-280.
- [3] BOREL, A. *Linear Algebraic Groups*. W. A. Benjamin, Inc., New York, 1969.
- [4] FREUDENTHAL, H. and H. de VRIES. *Linear Lie Groups*. Academic Press, New York, 1969.
- [5] KLEIN, F. *Lectures on Mathematics, Lectures II and III: Sophus Lie*. The Evanston Colloquium, MacMillan and Co., 1893. Republished by Amer. Math. Soc., New York, 1911.
- [6] — *Vorlesungen über höhere Geometrie*. Springer-Verlag, Berlin 1926. 3. Aufl. reprinted by Chelsea, New York, 1957.