

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	25 (1979)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	FIFTEEN CHARACTERIZATIONS OF RATIONAL DOUBLE POINTS AND SIMPLE CRITICAL POINTS
Autor:	Durfee, Alan H.
Kapitel:	4. Absolutely isolated double points
DOI:	https://doi.org/10.5169/seals-50375

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

functions f listed in column 1 of Table 1 have minimal resolutions as in column 3. (I believe that this first appeared in [Hirzebruch 1].) The converse follows since the singularities listed are taut [Brieskorn 2; Tjurina 3; Laufer 4]. (Two resolutions $\pi: M \rightarrow V$ and $\pi': M' \rightarrow V'$ are *topologically equivalent* if their exceptional sets are homeomorphic by a homeomorphism preserving the self-intersection numbers. A singularity V is *taut* if any other singularity with a good resolution topologically equivalent to a good resolution of V is then isomorphic to V .)

The classification of rational double points has been generalized in several ways: to rational triple points [Artin, p. 135], to elliptic singularities [Wagreich 1], and to minimally elliptic singularities [Laufer 5]. The Dynkin diagrams B_n , C_n , F_4 and G_2 occur when resolving singularities over non-algebraically closed fields [Lipman 1]. There is also a relation with simple complex Lie groups [Brieskorn 3].

4. ABSOLUTELY ISOLATED DOUBLE POINTS

There are at least three methods of resolving the singularity of the germ of a normal two-dimensional complex space V . The first method is one of local uniformization; this is originally due to Jung, and is described in detail in [Laufer 1]. The second method, due to Zariski, is to alternately blow up points and normalize. The third method (which generalizes to higher dimensions), is to blow up points and non-singular curves.

The singularity of V is *absolutely isolated* if it may be resolved by blowing up points alone, that is, it is not necessary to normalize or blow up curves. For example, the singularity of the zero locus of $f(x, y, z) = x^k + y^k + z^k$ is absolutely isolated, since it may be resolved by blowing up the origin once.

The singularity of V is a *double point* if its local ring is of multiplicity two. If V is $f^{-1}(0)$, this is equivalent to the lowest non-zero homogeneous term in the power series expansion of f being quadratic.

Characterization A4. The singularity of $f^{-1}(0)$ is an absolutely isolated double point.

The equivalence of Characterizations A1 and A4 was proved directly in [Kirby]. Later, it was shown [Tjurina 2; Lipman 1] that all rational singularities are absolutely isolated (thus showing Characterization A2 implies A4), and in [Brieskorn 1, Satz 1] that A4 implies A3.