

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 25 (1979)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: FIFTEEN CHARACTERIZATIONS OF RATIONAL DOUBLE POINTS AND SIMPLE CRITICAL POINTS
Autor: Durfee, Alan H.
Kapitel: 6. The local fundamental group
DOI: <https://doi.org/10.5169/seals-50375>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Proposition 5.2 shows that characterizations A5' and A1 are equivalent. Clearly Characterization A5' implies A5; since A5 implies A2, they are all equivalent.

COROLLARY 5.3. *Let G be a small finite subgroup of $GL(2, \mathbb{C})$. Then $G \subset SL(2, \mathbb{C})$ if and only if \mathbb{C}^2/G embeds in codimension one.*

This corollary follows from the above case-by-case analysis. J. Wahl points out that it is also possible to prove it directly, using the following two facts:

Fact 1. Let G be a small finite subgroup of $GL(2, \mathbb{C})$. Then $G \subset SL(2, \mathbb{C})$ if and only if the singularity of \mathbb{C}^2/G is Gorenstein.

This is a special case of [Watanabe]. A germ of a normal two-dimensional complex space is *Gorenstein* if there is a nowhere-vanishing holomorphic two-form on its regular points.

Fact 2. Let V be the germ at \mathfrak{v} of a two-dimensional rational singularity. Then V is Gorenstein if and only if V embeds in codimension 1.

Proof. Any singularity embedded in codimension one is Gorenstein. Conversely, suppose V is Gorenstein. Let $\pi: M \rightarrow V$ be the minimal resolution of V , and let $E_1 \cup \dots \cup E_s = \pi^{-1}(\mathfrak{v})$ be its exceptional set as in Section 3. Since V is Gorenstein, there is a divisor K on M (the *canonical class*) satisfying the adjunction formula. Furthermore $K \cdot E_i \geq 0$ for all i since the resolution is minimal, so $K \leq 0$ [Artin, bottom of p. 130]. If $K < 0$, then $-K > 0$, so arithmetic genus p of $-K$ satisfies $p(-K) \leq 0$ [Artin, Proposition 1]. On the other hand, $p(-K) = 1 - \chi(-K) = 1$ by the Riemann-Roch Theorem, a contradiction. Hence $K = 0$. Thus $K \cdot E_i = 0$ for all i , so V is a double point and embeds in codimension one, as in the proof that Characterization A3 implies Characterization A2.

6. THE LOCAL FUNDAMENTAL GROUP

Let V be the germ of a normal two-dimensional complex analytic space with an isolated singularity at \mathfrak{v} . Without loss of generality, we may assume that V is a *good neighborhood* of \mathfrak{v} , that is, that there is a neighborhood basis V_i of \mathfrak{v} in V such that each $V_i - \mathfrak{v}$ is a deformation retract of $V - \mathfrak{v}$ [Prill]. The *local fundamental group* of V at \mathfrak{v} is then defined as $\pi_1(V - \mathfrak{v})$. This group is trivial if and only if V is nonsingular at \mathfrak{v} [Mumford].

PROPOSITION 5.1 (bis). *The following statement is equivalent to those listed above.*

(d) *The local fundamental group of V is finite.*

It is shown in [Prill, p. 381; Brieskorn 2, p. 344] that conditions (a) and (d) are equivalent.

Characterization A6. The local fundamental group of $f^{-1}(0)$ is finite. Thus Characterizations A5 and A6 are equivalent.

There is an algorithm for computing the local fundamental group of V from a resolution [Mumford], and singularities V with finite, nilpotent and solvable local fundamental group have been classified [Brieskorn 2; Wagreich 2]. When V is a complete intersection, this classification is particularly simple [Durfee 2, Proposition 3.3].

7. VOLUME

Let $f(x, y, z)$ be the germ at the origin $\mathbf{0}$ of a complex analytic function, and suppose that $f(\mathbf{0}) = 0$ and that the origin is an isolated critical point of f . There is an $\varepsilon > 0$ such that $f^{-1}(0)$ intersects all spheres of radius ε' about $\mathbf{0}$ transversally for $0 < \varepsilon' \leq \varepsilon$. (See Section 12.) For $t \in \mathbf{C}$, let

$$V_t = f^{-1}(t) \cap D_\varepsilon^6$$

where D_ε^6 is the closed disk of radius ε about $\mathbf{0}$. The function $f(x, y, z)$ takes the constant value t on V_t , so $\frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy + \frac{\partial f}{\partial z} dz \equiv 0$ there. Hence a nowhere-vanishing holomorphic two-form ω_t on V_t may be defined by the equivalent expressions

$$\omega_t = \frac{dy \wedge dz}{\partial f / \partial x} = \frac{dz \wedge dx}{\partial f / \partial y} = \frac{dx \wedge dy}{\partial f / \partial z},$$

Characterization A7. The integral $\int_{V_0} \omega_0 \wedge \bar{\omega}_0$ is finite.

Note that the form $\omega_0 \wedge \bar{\omega}_0$ takes positive real values. The equivalence of Characterizations A2 and A7 is due to Laufer, and follows easily from his expression for the geometric genus in terms of forms [Laufer 2, Corollary 3.6].

A different formulation of this characterization is due to E. Looijenga (unpublished): Let $\Delta(r) = \{t \in \mathbf{C} : t < r\}$, let

$$X(r) = f^{-1}(\Delta(r)) \cap D_\varepsilon^6$$

and let $\text{vol}(X(r))$ be its volume in \mathbf{C}^3 .