

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 25 (1979)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** FIFTEEN CHARACTERIZATIONS OF RATIONAL DOUBLE POINTS AND SIMPLE CRITICAL POINTS  
**Autor:** Durfee, Alan H.  
**Kapitel:** 9. CHARACTERIZATIONS UNDER RIGHT AND CONTACT EQUIVALENCE  
**DOI:** <https://doi.org/10.5169/seals-50375>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 05.03.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

The fundamental lemmas used in the classification are as follows:

*Lemma 8.2.* If  $\mathfrak{m}^{k+1} \subset \mathfrak{m}^2 \Delta f$  then  $f$  is  $k$ -determined.

For the proof, see [Arnold 1, Lemma 3.2; Zeeman, Theorem 2.9; Siersma, p. 8]. Note that  $\mathfrak{m}^{k-1} \subset \Delta f$  implies that  $\mathfrak{m}^{k+1} \subset \mathfrak{m}^2 \Delta f$ . The *corank* of  $f$  is defined as  $n + 1$  minus the rank of the Hessian matrix  $\{(\partial^2 f / \partial z_i \partial z_j)(\mathbf{0})\}$ . The proof of part (a) of the following lemma may be found in [Arnold 1, Lemma 4.1; Siersma Lemma 3.2].

*Splitting Lemma 8.3.* (a) Let  $f(z_0, \dots, z_n) \in \mathcal{F}$  be of corank  $r + 1$ . Then there is a  $g(z_0, \dots, z_r) \in \mathfrak{m}^3$  such that

$$f(z_0, \dots, z_n) \sim g(z_0, \dots, z_r) + z_{r+1}^2 + \dots + z_n^2.$$

(b) Let  $g(z_0, \dots, z_r)$  and  $g'(z_0, \dots, z_r) \in \mathcal{F} \cap \mathfrak{m}^3$ . If

$$g(z_0, \dots, z_r) + z_{r+1}^2 + \dots + z_n^2 \sim g'(z_0, \dots, z_r) + z_{r+1}^2 + \dots + z_n^2$$

then

$$g(z_0, \dots, z_r) \sim g'(z_0, \dots, z_r).$$

The classification proceeds by low corank and low Milnor number. A germ of corank 0 is right equivalent to  $z_0^2 + \dots + z_n^2$ , a germ of corank 1 and Milnor number  $k > 1$  is right equivalent to  $z_0^{k+1} + z_1^2 + \dots + z_n^2$ , and so forth. The proofs are not hard [Arnold 1, Zeeman, Siersma]. Table 2, for instance, includes all right-equivalence classes of germs with Milnor number  $\mu \leq 9$ .

## 9. CHARACTERIZATIONS UNDER RIGHT AND CONTACT EQUIVALENCE

*Characterization B1.* The germ  $f$  is right equivalent to one of the germs in Table 2a.

*Characterization B2.* The germ  $f$  is contact equivalent to one of the germs in Table 2a.

When  $n = 2$ , Characterization B2 is the same as Characterization A1. Clearly Characterization B1 implies Characterization B2. Since all the germs in Table 2a are weighted homogeneous (§16), the converse follows from the next lemma.

*Lemma 9.1.* Let  $g$  be a weighted homogeneous polynomial, and suppose that a germ  $f \in \mathcal{F}$  is contact equivalent to  $g$ . Then  $f$  is right equivalent to  $g$ .

*Proof.* To say that  $f$  is contact equivalent to  $g$  means that there is a germ of an analytic isomorphism  $h: (\mathbb{C}^{n+1}, \mathbf{0}) \rightarrow (\mathbb{C}^{n+1}, \mathbf{0})$  and a function  $u: \mathbb{C}^{n+1} \rightarrow \mathbb{C}$  with  $u(\mathbf{0}) \neq 0$  such that  $f = u \cdot (g \circ h)$ . Let  $h = (h^0, \dots, h^n)$  be the components of  $h$ , and suppose that  $g$  is weighted homogeneous with weights  $(w_0, \dots, w_n)$ . Then,

$$\begin{aligned} f(z_0, \dots, z_n) &= u(z_0, \dots, z_n) \cdot g(h^0(z_0, \dots, z_n), \dots, h^n(z_0, \dots, z_n)) \\ &= g((u(z_0, \dots, z_n))^{1/w_0} h^0(z_0, \dots, z_n), \dots, \\ &\quad (u(z_0, \dots, z_n))^{1/w_n} h^n(z_0, \dots, z_n)). \end{aligned}$$

Hence  $f$  is right equivalent to  $g$ .

## 10. DEGENERATION

Let  $J_k$  be the set of  $k$ -jets of germs in  $\mathcal{O}$ . There is a projection of  $\mathcal{O}$  to  $J_k$  by mapping germs to their power series expansion truncated in degree  $k$ . The ring  $\mathcal{O}$  becomes a topological space by letting a basis of open sets be inverse images of open sets in  $J_k$ , for all  $k$ .

The group of germs of analytic automorphisms fixing  $\mathbf{0}$  acts on  $\mathcal{O}$ , and the orbits of this action (*right-equivalence orbits*) are the right-equivalence classes. Similarly, there is a contact equivalence group which acts on  $\mathcal{O}$ , and the orbits of this action (*contact-equivalence orbits*) are the contact equivalence classes [Mather, §2]. A right-equivalence orbit is always contained in a contact-equivalence orbit; Lemma 9.1 says that the right-equivalence orbit of a germ  $f$  in Table 2a or b equals its contact-equivalence orbit.

A subset  $A$  of  $\mathcal{O}$  is said to *right* (or *contact*) *degenerate* to a subset  $B$  of  $\mathcal{O}$  if the closure of the right (or contact) equivalence orbit of  $A$  contains  $B$ . If  $A$  degenerates to  $B$ , then  $B$  *simplifies* to  $A$  (written  $A \leftarrow B$ ). Right degeneracy is also called *adjacency*. For example, when  $n = 0$ , the germ  $z_0^k$  right degenerates to the germ  $z_0^l$  for  $k < l$ , since the one-parameter family  $tz_0^k + (1-t)z_0^l$  is  $z_0^l$  when  $t = 0$ , and is right-equivalent to  $z_0^k$  when  $t \neq 0$ . All germs of low codimension can be arranged according to right degeneracy in fascinating tables [Arnold 3; Siersma]. Table 3 lists some (but not all) of the simplifications that occur. The following proposition is a principal consequence of the work on degeneration.

### PROPOSITION 10.1.

- (i) *The germs in Table 2a right simplify only to each other.*
- (ii) *The germs in Table 2b right simplify only to the germs in Table 2a.*