Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 25 (1979)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: FIFTEEN CHARACTERIZATIONS OF RATIONAL DOUBLE POINTS AND SIMPLE CRITICAL POINTS

Autor: Durfee, Alan H.

Kapitel: 12. The quadratic form

DOI: https://doi.org/10.5169/seals-50375

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 19.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

12. The quadratic form

Let $f(z_0, ..., z_n)$ be a germ with $f(\mathbf{0}) = 0$ and an isolated critical point at $\mathbf{0}$ (that is, a germ in $\mathscr{F}$). There is an $\varepsilon > 0$ such that $f^{-1}(0)$ intersects all spheres of radius ε' about $\mathbf{0}$ transversally for $0 < \varepsilon' \leqslant \varepsilon$. For suitably small $\delta > 0$, $f^{-1}(\delta')$ intersects the closed disk D_ε^{2n+2} of radius ε transversally for all $|\delta'| \leqslant \delta$. Let

$$F = f^{-1}(\delta) \cap D_\varepsilon^{2n+2}$$

be the *Milnor fiber* of f [Milnor 1]. The set F is a smooth real $2n$-manifold with boundary whose diffeomorphism type is independent of the choice of ε and δ. Furthermore, F is $(n-1)$-connected, and the Milnor number μ of §7 is the rank of $H_n(F)$. The Milnor number is zero if and only if the germ f has a regular point at $\mathbf{0}$ [Milnor 1, Corollary 7.3]. The *intersection pairing* $(\ ,\ )$ of F is the integral bilinear form $H_n(F) \times H_n(F) \to \mathbf{Z}$ defined by sending (x, y) to $(x' \cup y')[F]$, where x' and y' in $H^n(F, \partial F)$ are Lefschetz duals to x and y, and $[F]$ in $H_{2n}(F, \partial F)$ is the orientation class of F given by the underlying complex structure. The intersection pairing is symmetric if n is even, and skew symmetric if n is odd. For example, the germ $f(z_0, ..., z_n) = z_0^2 + ... + z_n^2$ has $H_n(F)$ a free cyclic group with generator e, and $(e, e) = 2(-1)^{n/2}$ or 0 according as n is even or odd. There are many methods of computing the intersection pairing in special cases.

By a tensor product theorem [Gabrielov 1; Sakamoto], the Milnor numbers of $f(z_0, ..., z_n)$ and $f(z_0, ..., z_n) + z_{n+1}^2 + ... + z_m^2$ are equal. The *quadratic form* of $f(z_0, ..., z_n)$ is defined to be the intersection pairing of the germ $f(z_0, ..., z_n) + z_{n+1}^2 + ... + z_m^2$ where $m \equiv 2 \pmod 4$. This is independent of the choice of m. For example, if $n \equiv 0 \pmod 4$ then the quadratic form of f is the negative of its intersection pairing; all this follows from the tensor product theorem. See also [Kauffmann and Neumann].

A germ f *topologically degenerates* to a germ g if there is an $\eta > 0$ and a family h_t of germs for $\{t \in \mathbf{C} : |t| < 2\eta\}$ with $h_\eta \sim f$, $h_0 \sim g$, and h_t of constant Milnor number for $t \neq 0$. Compare [Lê and Ramanujam]. Clearly right degeneracy implies topological degeneracy.

Lemma 12.1 [Tjurina 1, Theorem 1]. If f topologically degenerates to g, then there is an injection of $H_n(F_f)$ into $H_n(F_g)$ (where F_f is the Milnor fiber of f, and F_g is the Milnor fiber of g), and this injection preserves the intersection pairing. In particular, if g topologically degenerates to f as well, then the intersection pairings of f and g are isomorphic.

Characterization B5. The quadratic form of f is negative definite.

The equivalence of Characterizations B1 and B5 is proved in [Tjurina 1]. By explicit computation the quadratic forms of the germs in Table 2a are shown to be negative definite, and those of Table 2b are shown to be negative semi-definite. (In fact, the quadratic form of a germ in Table 2a is isomorphic to the intersection pairing of its minimal resolution, and the quadratic form of a germ of type $\tilde{E}_k$ in Table 2b is isomorphic to the quadratic form of E_k plus a two-dimensional zero form.) The result then follows from Proposition 10.1 and Lemma 12.1. When $n = 2$, the Milnor fiber F is in fact diffeomorphic to the minimal resolution M of $f^{-1}(0)$, since the singularity of $f^{-1}(0)$ is an absolutely isolated double point [Brieskorn 1, Theorem 4; Tjurina 1, Lemma 1].

When $n = 2$, the equivalence of Characterizations A2 and B5 follows from the following result [Durfee 2, Proposition 3.1].

THEOREM 12.2. *Twice the geometric genus p of $f^{-1}(0)$ equals the number of positive plus the number of zero diagonal elements in a diagonalization of the intersection pairing over the real numbers.*

The classification of germs according to signature of the quadratic form has been extended in [Arnold 3]; see also [Durfee 2, Proposition 3.3].

13. NEARBY MORSE FUNCTIONS

A *deformation* of a germ $f \in \mathscr{F}$ is a germ $g: \mathbf{C}^{n+1} \times \mathbf{C} \to \mathbf{C}$ with $g(z, 0) = f(z)$. Choose ε and δ for f as in §11. Then choose $\eta > 0$ such that for all $|t| < \eta$ and $|\delta'| \leqslant \delta$, the set $\{z \in \mathbf{C}^{n+1} : g(z, t) = \delta'\}$ intersects S_ε^{2n+1} transversally and the critical values of $g(z, t)$ for fixed t are less than δ in absolute value. A germ $\bar{f}$ is a *nearby Morse function* to f if $\bar{f}$ has only non-degenerate critical points in D_ε^{2n+2} and there is a deformation g and a t_0 with $|t_0| < \eta$ such that $\bar{f}(z) = g(z, t_0)$.

Characterization B6. There is a nearby Morse function to f with one or two critical values.

In fact, the nearby Morse function has one critical value if and only if f is right equivalent to A_2, since the quadratic form diagram is connected (§14). This surprising characterization is in [A'Campo 2II], where it is shown that Characterization B1 implies B6, and B6 implies B7 below.