

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 25 (1979)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** TERMWISE AVERAGES OF TWO DIVERGENT SERIES  
**Autor:** Ash, J. Marshall / Sexton, Harlan  
**DOI:** <https://doi.org/10.5169/seals-50377>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 15.03.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# TERMWISE AVERAGES OF TWO DIVERGENT SERIES

by J. Marshall ASH and Harlan Sexton

*Definition.* For  $a, b > 0$ , let  $M_\infty(a, b) = \max\{a, b\}$ ,  $M_r(a, b) = \left(\frac{a^r + b^r}{2}\right)^{\frac{1}{r}}$  for finite non-zero  $r$ ,  $M_0(a, b) = \sqrt{ab}$ , and  $M_{-\infty}(a, b) = \min\{a, b\}$ .

*Definition.* The sequence  $\{a_n\}$ ,  $n = 1, 2, \dots$ , is *convex* if  $a_{n+2} - 2a_{n+1} + a_n \geq 0$ ,  $n = 1, 2, \dots$ .

If  $\{a_n\}$  is convex then the union of the line segments  $(n, a_n) (n+1, a_{n+1})$ ,  $n = 1, 2, \dots$ , is the graph of a continuous convex function on the interval  $[1, \infty)$ .

Let  $\{a_n\}$ ,  $\{b_n\}$  be sequences of positive numbers. If  $\sum a_n$  and  $\sum b_n$  are finite, so is  $\sum_n M_r(a_n, b_n)$  since  $\sum M_\infty(a_n, b_n) < \sum (a_n + b_n) < \infty$  and  $M_r(a, b)$  is an increasing function of  $r$ ,  $-\infty \leq r \leq \infty$  [Hardy, Littlewood, Polya, *Inequalities*, Cambridge Univ. Press, Cambridge (1973), pp. 15, 26]. If either  $\sum a_n = \infty$  or  $\sum b_n = \infty$  and if  $r > 0$ , then  $\sum_n M_r(a_n, b_n) = \infty$  since

$$\begin{aligned} \sum M_r(a_n, b_n) &= \sum \left(\frac{a_n^r + b_n^r}{2}\right)^{\frac{1}{r}} \geq 2^{-1/r} \sum M_\infty(a_n, b_n) \\ &\geq 2^{-1/r} \max\{\Sigma a_n, \Sigma b_n\} = \infty. \end{aligned}$$

If however,  $r \leq 0$ , the situation is entirely different.

**THEOREM.** *There are convex monotonically decreasing to zero sequences  $\{a_n\}$ ,  $\{b_n\}$  with  $\sum a_n = \sum b_n = \infty$ , such that  $\sum_n M_r(a_n, b_n) < \infty$  for  $-\infty \leq r \leq 0$ .*

*Remark 1.* The most interesting special cases are  $r = -\infty$  where the conclusion is  $\sum \min\{a_n, b_n\} < \infty$  and  $r = 0$  where the conclusion is  $\sum \sqrt{a_n b_n} < \infty$ . Since  $\lim_{r \rightarrow 0+} M_r(a, b) = M_0(a, b)$  [ibid, p. 15] the theorem coupled with its preceding remarks form a classification with a tidy "dividing line".



and similarly  $B_j$  will be the first point of the  $j$ -th block of  $\beta$ 's,  $B'_j$  will lie on the axis below the last point of the  $\beta$ 's  $j$ -th block,  $m_j$  will be the line connecting  $B_j$  with  $B_{j+1}$ , and  $m'_j$  will be the line connecting  $B_j$  with  $B'_j$ .

We will construct the  $\alpha$ 's and  $\beta$ 's one block at a time. Let  $\{c_n\}$  be any convergent sequence with all  $c_n > 0$ . Let  $\alpha_1 = \alpha_2 = 1$  and  $\beta_1 = c_1^2$  so that the first block of  $\alpha$ 's has length 2 and the first block of  $\beta$ 's has length 1. Continue inductively as follows. We suppose that after the  $n$ -th stage  $n$  blocks of  $\alpha$ 's and  $\beta$ 's have been chosen so that

- (1)  $\sum \alpha \geq n + 1, \sum \beta \geq n - 1, \alpha \searrow, \beta \searrow$
- (2)  $A'_n > B'_n > A'_{n-1}$  (We identify the point  $A'_n$  with its first co-ordinate; here  $A'_0$  is taken to be zero.)
- (3)  $\sum_{j=1}^{B'_n} \sqrt{\alpha_j \beta_j} \leq c_1 + \dots + c_n$
- (4) The polygonal paths  $l_1 l_2 \dots l_{n-1} l'_n$  and  $m_1 m_2 \dots m'_n$  are convex.

To reach the next stage of the construction first pick  $A'_n - B'_n$   $\beta$ 's all equal to  $B$  where  $B > 0$  is so small that  $\sum_{j=B'_n+1}^{A'_n} \sqrt{\alpha_j B} \leq \frac{1}{2} c_{n+1}$  and so

small that  $B < \beta_{B'_n}$ . Then pick sufficiently many more  $\beta$ 's of this same size  $B$  so that there are now more  $\beta$ 's than  $\alpha$ 's, and so that  $\sum \beta \geq n$ , and so that the path  $m_1 \dots m_n m'_{n+1}$  is convex. In much the same manner we now pick

$B'_{n+1} - A'_n$   $\alpha$ 's all equal to  $A$  where  $A$  is so small that  $\sum_{j=A'_n+1}^{B'_{n+1}} \sqrt{AB} = (B'_{n+1} - A'_n) \sqrt{AB} \leq \frac{1}{2} c_{n+1}$  and so small that  $A < \alpha_{A'_n}$ . Then pick

sufficiently many more  $\alpha$ 's of this same size  $A$  so that there are now more  $\alpha$ 's than  $\beta$ 's, and so that  $\sum \alpha \geq n + 2$ , and so that the path  $l_1 \dots l_n l'_{n+1}$  is convex. Now (1)-(4) hold with  $n$  replaced by  $n + 1$ .

Following a suggestion of Andrejs Treibergs we complete the proof as follows. Define the sequence  $\{a_k\}$  ( $\{b_k\}$ ) as the projection of the  $\alpha$ 's ( $\beta$ 's) down onto their supporting lines  $l_j$  ( $m_j$ ).




Figure 2

Then  $\sum \sqrt{a_k b_k} < \sum \sqrt{\alpha_k \beta_k} \leq \sum c_n < \infty$ . Clearly  $\{a_k\}$  and  $\{b_k\}$  are convex and monotonically decreasing to zero, and  $\sum a_k > \sum a_k^* = \frac{1}{2} \sum \alpha_k = \infty$  since  $\sum_{k=r}^s a_k^* = \frac{1}{2} \sum_{k=r}^s \alpha_k$ . Similarly,  $\sum b_k = \infty$ .

(Reçu le 29 juin 1978)

J. Marshall Ash  
Harlan Sexton

Department of Mathematics  
De Paul University  
2323 North Seminary Av.  
Chicago, Illinois 60614

Stanford University  
Stanford, California 94305