

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 26 (1980)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE FAST SKEW-CLOSURE ALGORITHM
Autor: Fischer, M. J. / Paterson, M. S.
Kurzfassung
DOI: <https://doi.org/10.5169/seals-51079>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 16.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

THE FAST SKEW-CLOSURE ALGORITHM ¹⁾

by M. J. FISCHER and M. S. PATERSON

ABSTRACT. A subtle matrix algorithm is explored and generalized. Originally used for transitive closures of symmetric Boolean matrices, this $O(n^2)$ algorithm computes a closure operation which is of interest for asymmetric and non-Boolean matrices too. The correctness of a generalized form of the algorithm is shown. The monoid generated by "skew-closure" and some of the more usual closures is investigated.

1. INTRODUCTION

The algorithm which forms the principal theme of this paper is of interest for several reasons. It is of mysterious ancestry; we have been unable to trace any published source which refers to it. It came to us by oral tradition at least seven years ago, when it impressed us with its speed and by the non-triviality of establishing its correctness. Further, whereas it seemed intended to be applied to Boolean matrices of symmetric and reflexive relations, the result of an application to more general matrices invited analysis.

The operation achieved by the algorithm we have termed "skew-closure". This closure is related to the more customary symmetric-and-transitive closure, and belongs to a very natural class of closure operations which we elaborate a little. In the interests of finding which matrix operations can be done equally rapidly, the monoid generated by several of these simple closures is treated in some detail. While this is finite, we later display a pair of slightly more complicated closures which together yield an infinite monoid.

¹⁾ Presented at the *Symposium über Logik und Algorithmik* in honour of Ernst SPECKER, Zürich, February 1980.