

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 27 (1981)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ON THE GENUS OF GENERALIZED FLAG MANIFOLDS  
**Autor:** Glover, Henry H. / Mislin, Guido  
**Kapitel:** §1. Genus and self maps  
**DOI:** <https://doi.org/10.5169/seals-51749>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 19.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

**THEOREM.** Let  $M = M(n_1, \dots, n_k)$  be a generalized flag manifold for which the Conjecture C holds. Then

$$G(M) = \{[M]\}.$$

In particular the Grassmann manifolds  $G_p(\mathbb{C}^{p+q})$  for  $p \neq q$  and the flag manifolds  $U(n)/T^n$  are all generically rigid.

### §1. GENUS AND SELF MAPS

Let  $P$  denote a fixed set of primes. Two  $P$ -sequences

$$S_1, S_2 : P \rightarrow E(X_0)$$

are called *equivalent*, if there exist maps  $h(0) \in E(X_0)$  and

$$h(p) \in \text{im}(E(X_p) \xrightarrow{\text{can}} E(X_0))$$

such that for all  $p \in P$  one has

$$h(0) S_1(p) = S_2(p) h(p).$$

**Definition 1.1.** We denote by  $P\text{-Seq}(E(X_0))$  the set of equivalence classes of  $P$ -sequences in  $E(X_0)$ .

If  $P$  is a finite set of primes and  $X$  a nilpotent space of finite type, then there is a canonical map

$$\theta : G(X) \rightarrow P\text{-Seq}(E(X_0)).$$

It is defined as follows. Let  $Y \in G(X)$  and  $P = \{p_1, \dots, p_n\}$ . Then the localization  $Y_P$  is a pull-back of maps  $X_{p_i} \xrightarrow{\lambda_i} X_0$ , i.e.  $Y_P \simeq \text{hoinvlim} \{X_{p_i} \xrightarrow{\lambda_i} X_0\}$ . The maps  $\lambda_i$  induce equivalences  $\bar{\lambda}_i \in E(X_0)$  and we put

$$\theta(Y) = \{[\bar{\lambda}_1, \bar{\lambda}_2, \dots, \bar{\lambda}_n]\}.$$

If  $Y_P$  may also be represented by  $\text{hoinvlim} \{X_{p_i} \xrightarrow{\mu_i} X_0\}$ , then there exist maps  $h(0) \in E(X_0)$  and  $\tilde{h}(p_i) \in E(X_{p_i})$ ,  $i \in \{1, \dots, n\}$  rendering the diagrams

$$\begin{array}{ccc}
 X_{p_i} & \xrightarrow{\tilde{h}(p_i)} & X_{p_i} \\
 \downarrow \lambda_i & & \downarrow \mu_i \\
 X_0 & \xrightarrow{h(0)} & X_0
 \end{array}$$

homotopy commutative and thus inducing  $\text{hoinvlim } \{\lambda_i\} \simeq \text{hoinvlim } \{\mu_i\}$ . Hence

$$\{[\bar{\lambda}_1, \dots, \bar{\lambda}_n]\} = \{[\bar{\mu}_1, \dots, \bar{\mu}_n]\} \in P\text{-Seq}(E(X_0))$$

and therefore  $\theta$  is well defined.

**LEMMA 1.2.** Let  $X$  be a nilpotent space of finite type and let  $P$  denote a finite set of primes. Then

$$\theta: G(X) \rightarrow P\text{-Seq}(E(X_0))$$

is surjective with fibers of the form

$$\theta^{-1}(\theta(Y)) = \{Z \in G(X) \mid Z_P \simeq Y_P\}.$$

*Proof.* Let  $P = \{p_1, \dots, p_n\}$  and

$$\{[\bar{f}_1, \dots, \bar{f}_n]\} \in P\text{-Seq}(E(X_0)).$$

Let  $e_i: X_{p_i} \rightarrow X_0$  denote the canonical maps and put

$$f_i = \bar{f}_i \circ e_i: X_{p_i} \rightarrow X_0.$$

Define  $W = \text{hoinvlim } \{f_i\}$ ;  $W$  comes equipped with a canonical map  $f: W \rightarrow X_0$ . Let  $Z$  be the homotopy pull back of  $W \xrightarrow{f} X_0 \xleftarrow{\text{can}} X_{\bar{P}}$ , where  $\bar{P}$  denotes the set of primes complementary to  $P$ . Then  $Z \in G(X)$  and

$$\theta(Z) = \{[\bar{f}_1, \dots, \bar{f}_n]\};$$

thus  $\theta$  is surjective. It is clear from the definition of  $\theta$  that for  $Y, Z \in G(X)$  one has  $\theta(Y) = \theta(Z)$  if and only if  $Y_P \simeq Z_P$ .

The next lemma provides a sufficient condition for  $\theta$  to be monic “at the basepoint”.

LEMMA 1.3. Let  $X$  be a nilpotent space of finite type. Suppose that there exists a finite set of primes  $P$  with complement  $\bar{P}$  such that

- a)  $Y \in G(X)$  implies  $Y_{\bar{P}} \simeq X_{\bar{P}}$
- b) every  $f \in E(X_0)$  can be written as  $f = f_1 \circ f_2$  with  $f_1 \in \text{im}(E(X_P) \xrightarrow{\text{can}} E(X_0))$  and  $f_2 \in \text{im}(E(X_{\bar{P}}) \rightarrow E(X_0))$ .

Then for  $\theta: G(X) \rightarrow P\text{-Seq}(E(X_0))$  as above, one has  $\theta^{-1}(\theta(X)) = \{X\}$ .

*Proof.* Let  $Y \in G(X)$  with  $\theta(Y) = \theta(X)$ . Then  $Y_P \simeq X_P$  by the definition of  $\theta$ , and  $Y_{\bar{P}} \simeq X_{\bar{P}}$  by assumption. Hence  $Y$  may be obtained as a homotopy pull back of the form

$$\begin{array}{ccc} & X_P & \\ Y \swarrow & \alpha \searrow & \\ & X_0 & \\ & \beta \nearrow & \\ & X_{\bar{P}} & \end{array}$$

If  $\alpha$  induces  $\bar{\alpha} \in E(X_0)$  and if  $\gamma = \bar{\alpha}^{-1} \circ \beta$ , then  $Y$  is also a pull back of the form

$$\begin{array}{ccc} & X_P & \\ Y \swarrow & \text{can} \searrow & \\ & X_0 & \\ & \gamma \nearrow & \\ & X_{\bar{P}} & \end{array}$$

Let  $\bar{\gamma} \in E(X_0)$  be the map induced by  $\gamma$  and write  $\bar{\gamma} = f_1 f_2$  with

$$f_1 \in \text{im}(E(X_P) \rightarrow E(X_0)), \quad f_2 \in \text{im}(E(X_{\bar{P}}) \rightarrow E(X_0)).$$

Choose a lift  $\tilde{f}_1^{-1} \in E(X_P)$  of  $f_1^{-1}$  and a lift  $\tilde{f}_2 \in E(X_{\bar{P}})$  of  $f_2$ . Then  $f_1^{-1} \circ \gamma = \text{can} \circ \tilde{f}_2$  and one can form a commutative diagram,

$$\begin{array}{ccccc}
 & & f_1^{-1} & & \\
 & X_P & \xrightarrow{\hspace{2cm}} & X_P & \\
 & \searrow \text{can} & & \swarrow \text{can} & \\
 & X_0 & \xrightarrow{f_1^{-1}} & X_0 & \\
 & \nearrow \gamma & & \swarrow \text{can} & \\
 X_{\bar{P}} & \xrightarrow{\tilde{f}_2} & X_{\bar{P}} & &
 \end{array}$$

which shows that  $Y \simeq X$ .

## §2. THE CASE OF GENERALIZED FLAG MANIFOLDS

The following result is an easy consequence of [F].

**LEMMA 2.1.** Let  $M$  be a generalized flag manifold. Then the following holds.

- a) If  $g(\lambda) \in Gr(M_0)$  is a grading map with  $\lambda \in \mathbf{Z}_Q^*$  for some (not necessarily finite) set of primes  $Q$ , then  $g(\lambda)$  lifts to a homotopy equivalence  $\tilde{g}(\lambda): M_Q \rightarrow M_Q$ .
- b) Let  $P$  be an arbitrary set of primes with complement  $\bar{P}$ . Then every

$$f \in \langle Gr(M_0), N(H)/H \rangle$$

may be written in the form  $f = f_1 \circ f_2$  with

$$f_1 \in \text{im}(E(M_P) \rightarrow E(M_0))$$

and

$$f_2 \in \text{im}(E(M_{\bar{P}}) \rightarrow E(M_0)).$$

*Proof.* Let  $\lambda = k/l$  with  $k$  and  $l$  relatively prime integers. Then  $g(k)$  and  $g(l)$  lift to equivalences

$$\tilde{g}(k), \tilde{g}(l): M_Q \rightarrow M_Q$$