

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 27 (1981)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ON THE GENUS OF GENERALIZED FLAG MANIFOLDS  
**Autor:** Glover, Henry H. / Mislin, Guido  
**Kapitel:** §2. The case of generalized flag manifolds  
**DOI:** <https://doi.org/10.5169/seals-51749>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 19.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

$$\begin{array}{ccccc}
 & & f_1^{-1} & & \\
 & X_P & \xrightarrow{\hspace{2cm}} & X_P & \\
 & \searrow \text{can} & & \swarrow \text{can} & \\
 & X_0 & \xrightarrow{f_1^{-1}} & X_0 & \\
 & \nearrow \gamma & & \swarrow \text{can} & \\
 X_{\bar{P}} & \xrightarrow{\tilde{f}_2} & X_{\bar{P}} & &
 \end{array}$$

which shows that  $Y \simeq X$ .

## §2. THE CASE OF GENERALIZED FLAG MANIFOLDS

The following result is an easy consequence of [F].

**LEMMA 2.1.** Let  $M$  be a generalized flag manifold. Then the following holds.

- a) If  $g(\lambda) \in Gr(M_0)$  is a grading map with  $\lambda \in \mathbf{Z}_Q^*$  for some (not necessarily finite) set of primes  $Q$ , then  $g(\lambda)$  lifts to a homotopy equivalence  $\tilde{g}(\lambda): M_Q \rightarrow M_Q$ .
- b) Let  $P$  be an arbitrary set of primes with complement  $\bar{P}$ . Then every

$$f \in \langle Gr(M_0), N(H)/H \rangle$$

may be written in the form  $f = f_1 \circ f_2$  with

$$f_1 \in \text{im}(E(M_P) \rightarrow E(M_0))$$

and

$$f_2 \in \text{im}(E(M_{\bar{P}}) \rightarrow E(M_0)).$$

*Proof.* Let  $\lambda = k/l$  with  $k$  and  $l$  relatively prime integers. Then  $g(k)$  and  $g(l)$  lift to equivalences

$$\tilde{g}(k), \tilde{g}(l): M_Q \rightarrow M_Q$$

since necessarily  $k, l \in \mathbf{Z}_Q^*$  (compare [F]). Thus  $\tilde{g}(k)\tilde{g}(l)^{-1}$  is a lift of  $g(\lambda)$ . For b) we note that  $f = g(\rho) \circ \sigma$  for some  $\rho \in \mathbf{Q}^*$  and

$$\sigma \in N(H)/H.$$

If we write  $\rho = \rho_1 \cdot \rho_2$  with  $\rho_1 \in \mathbf{Z}_P^*$  and  $\rho_2 \in \mathbf{Z}_{\bar{P}}^*$ , then

$$f = g(\rho_1) \cdot (g(\rho_2) \sigma)$$

and we may choose

$$f_1 = g(\rho_1), f_2 = g(\rho_2) \sigma.$$

Since  $\sigma$  lifts even to  $E(M)$ , we infer by using a) that  $f_1$  and  $f_2$  lift as desired.

A final step towards proving the Theorem formulated in the introduction consists in the following.

**LEMMA 2.2.** Let  $M$  be a generalized flag manifold for which Conjecture C holds. Then for every finite set of primes  $P$ ,

$$P\text{-Seq}(E(M_0)) = \{[1, 1, \dots, 1]\}.$$

*Proof.* Let  $\{\mu_1, \dots, \mu_n\} \in P\text{-Seq}(E(M_0))$ , where  $P = \{p_1, \dots, p_n\}$  and

$$\mu_i \in \text{im}(E(M_{p_i}) \rightarrow E(M_0))$$

for all  $i$ . Then  $\mu_i = g(\lambda_i) \circ \sigma_i$  with  $\lambda_i \in \mathbf{Q}^*$  and

$$\sigma_i \in N(H)/H \subset E(M_0).$$

Define  $\lambda \in \mathbf{Q}^*$  by  $\lambda = \prod p_i^{m_i}$ , where  $m_i \in \mathbf{Z}$  is such that  $p_i^{m_i} \lambda_i \in \mathbf{Z}_{p_i}^*$ . Then  $g(\lambda) \mu_i = g(\lambda \lambda_i) \sigma_i$  with  $\lambda \lambda_i \in \mathbf{Z}_{p_i}^*$ . By Lemma 2.1 a) we know that  $g(\lambda \lambda_i)$  lifts to  $M_{p_i}$ , and since  $\sigma_i$  lifts even to  $M$  we conclude that

$$h(p_i) = g(\lambda \lambda_i) \sigma_i \in \text{im}(E(M_{p_i}) \rightarrow E(M_0))$$

for all  $i$ . The equation

$$g(\lambda) \mu_i = h(p_i), i \in \{1, \dots, n\}$$

show that  $\{\mu_1, \dots, \mu_n\} = \{[1, \dots, 1]\} \in P\text{-Seq}(E(M_0))$ .

The proof of the main Theorem:

Let  $M$  be a generalized flag manifold for which the Conjecture C holds. Since  $M$  is a formal space we can find for every  $N \in G(M)$  a rational equivalence

$f(N): N \rightarrow M$ . Let  $P(M)$  denote the set of primes which appear in any of the orders of

$$\ker(f(N)_*: H_*(N; \mathbf{Z}) \rightarrow H_*(M; \mathbf{Z}))$$

or  $\text{coker } f(N)_*$ ,  $N$  ranging over  $G(M)$ . The set  $P(M)$  is finite, since each  $\ker f(N)_*$  and  $\text{coker } f(N)_*$  is finite and since  $G(M)$  is a finite set by [W]. Consider now the map

$$\theta: G(M) \rightarrow P\text{-Seq } E(M_0)$$

with respect to this finite set of primes  $P(M) = P$ . Since  $P$  is finite,

$$P\text{-Seq } (E(M_0))$$

consists of only one element (Lemma 2.2). It remains to show that

$$\theta^{-1}(\theta(M)) = \{M\}.$$

For this we apply Lemma 1.3. Note that  $N \in G(M)$  implies  $N_{\bar{P}} \simeq M_{\bar{P}}$  since  $f(N): N \rightarrow M$  is a  $\bar{P}$ -equivalence. Moreover, the condition b) of 1.3 is satisfied in view of Lemma 2.1 b). Therefore we conclude that  $G(M) = \{[M]\}$  and the proof is completed.

*Note added in proof.* Since this paper went to press, we have been informed that Conjecture C has been proved for the case  $k = 2, n_1 = n_2$ , by M. Hoffman: "Cohomology endomorphisms of complex flag manifolds", Ph.D. dissertation, MIT 1981. As a consequence, it follows that all complex Grassmann manifolds are generically rigid.