

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 27 (1981)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SPECKER'S MATHEMATICAL WORK FROM 1949 TO 1979
Autor: Wang, Hao
Kapitel: Topology and recursive analysis
DOI: <https://doi.org/10.5169/seals-51741>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 21.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

notes that about half of Bernays' papers may be classified as philosophical: this difference between them may be partly due to the different historical periods in which they live.

ENJOYMENT OF INTERACTION

Among Specker's publications, several papers seem to have been stimulated primarily by the enjoyment of personal interaction. Thus the paper 1949c dealt with a problem of Sikorski who was visiting Zürich then, while the paper 1964 continued the study to more elaborate cases (3 and 18 in the above list). The papers 1957b and 1961b (10 and 16 in the above list) seem to belong to the class of papers which are provoked by the infinite supply of problems from Erdős. Paper 11 answers a problem raised by Mostowski. A most obviously playful paper is 1978b (30) which gives, for the recognition problem, the generating problem, and the counting problem of the partition of finite sets, algorithms programmable on the "toy" computer HP-25.

Several of these papers contain clever constructions which stimulate extensions and generalizations. For example, the paper 10 gives the Specker graph which shows:

$$\omega^3 \rightarrow (2, \omega^3)^2 \text{ and } \omega^3 \not\rightarrow (3, \omega^3)^2.$$

This leads to the function $f(n)$ such that $f(n) < \omega$,

$$\omega^n \rightarrow (f(n)-1, \omega^3)^2 \text{ and } \omega^n \not\rightarrow (f(n), \omega^3)^2.$$

Eva Nosal many years later showed that $f(n) = 2^{n-2} + 1$ for $n \geq 3$, *J. London Math. Soc. (2)*, 8 (1974), 306-310.

TOPOLOGY AND RECURSIVE ANALYSIS

It is interesting to observe that Specker's early papers of 1949 and 1950 have continued to interest mathematicians over the years. For example, the paper 2 gives a bounded increasing recursive sequence of rational numbers that does not converge to a recursive real number. In a recent paper by M. I. Kanovič, such sequences are called Specker sequences, and the complexity of "limit candidates" for a Specker sequence is studied with the result that the larger the complexity of the candidate, the closer it is to

the actual (more recursive) limit. The reference is: *Soviet Math. Dokl.*, vol. 15 (1974), No. 1, pp. 299-303.

The three papers on recursive analysis are remarkable in staying away from more controversial issues concerning constructivity. In particular, the first paper is probably the earliest use of recursive functions to elucidate constructive analysis.

Of the three papers on topology, I can out of ignorance have very little to say. The paper 1950b was done when Specker visited the Institute for Advanced Study. In it he is able to make interesting contributions to group theory as a topologist. It contains elegant ideas which received further development, for instance, almost two decades later in G. Nöbeling, "Verallgemeinerung eines Satzes von Herrn E. Specker", *Inventiones math.*, 6 (1968), 41-55. Let F be the Abelian group of sequences of integers with $\{a_n\} + \{b_n\} = \{a_n + b_n\}$. Specker shows that every countable subgroup of F is free and that F contains a nonfree subgroup of cardinality aleph-one (hence, F itself is not free). Furthermore, let F_b be the subgroup of F consisting of all bounded sequences of integers. Then it is shown that every subgroup of F_b with cardinality aleph-one is a free group. This last theorem is generalized by Nöbeling to show that for an arbitrary set X (rather than just the set of integers), the group of all bounded sequences from X is free and possesses a characteristic basis.

Specker's interest in the group F and its subgroups is suggested by the problem of determining the algebraic structure of the first cohomology group of an infinite complex, a problem studied in his dissertation 1949a. The paper 1950a considers end lattices and introduces what is known as Specker compactification which is investigated extensively, for instance, in Herbert Abels, Specker-Kompaktifizierung von lokal kompakten topologischen gruppen, *Math. Z.*, 135 (1974), 325-361.

AN EXAMPLE OF BEGINNING WITH CONCRETE PROBLEMS

In March 1966, Specker lectured in England on his result that Ramsey's theorem does not hold in recursive set theory. Afterwards, he was persuaded to present it at the Logic Colloquium of 1969 (and publish it as 24 in the above list). More exactly, Specker proves that there is a recursive (Σ_0) partition of the 2-elements sets of natural numbers which possesses no recursively enumerable (Σ_1) infinite sets of indiscernibles and that for every recursive partition there is always a Δ_3 set of indiscernibles. This suggests