

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 27 (1981)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: IDENTITIES FOR PRODUCTS OF GAUSS SUMS OVER FINITE FIELDS
Autor: Evans, Ronald J.
Kapitel: 4. Proof of (2)
DOI: <https://doi.org/10.5169/seals-51748>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

3. THEOREMS OF STICKELBERGER AND DAVENPORT-HASSE

We will make use of the following three classical formulas. First [3, (0.8)],

$$(6) \quad G_{f^m} \left(\chi^{\frac{q^m-1}{q-1}} \right) = G_f(\chi)^m,$$

where χ is a character on $GF(q^m)$. Next [3, (0.9)],

$$(7) \quad 1 = \frac{\chi^l(l)}{G_f(\chi^l)} \prod_{j=0}^{l-1} \frac{G_f(\chi\psi^j)}{G_f(\psi^j)},$$

where χ, ψ are characters on $GF(q)$ and ψ has order l . Finally [8], [5, p. 25]

$$(8) \quad \frac{G_f(\chi^\alpha)}{(\zeta - 1)^{s(\alpha)}} \equiv \frac{1}{\gamma(\alpha)} \equiv \frac{(\zeta - 1)^{\alpha - s(\alpha)}}{\alpha!} \pmod{P},$$

where α is an integer, $0 \leq \alpha < q - 1$; $s(\alpha)$ denotes the sum of the p -adic digits of α ; $\gamma(\alpha)$ denotes the product of the factorials of the p -adic digits of α ; P is a prime ideal above p in the ring $\mathcal{O} = \mathbb{Z}[\omega]$, where $\omega = \exp(2\pi i/p(q-1))$; and χ is the character on $\mathcal{O}/P \approx GF(q)$ of order $q - 1$ which maps the coset $\omega + P$ to $\bar{\omega}$.

4. PROOF OF (2)

Let η denote the right side of (2). We must show that $\eta = 1$. Let $\delta = \frac{q^n - 1}{q - 1}$, $\theta = w^{k-1}(cw + i_j)$. Using (6), we have

$$\eta^n = \frac{\chi^{ln}(l) G_{f^n}(\chi^\delta)}{G_{f^n}(\chi^{\delta l})} \prod_{j=1}^e \prod_{k=1}^r \prod_{c=1}^{w^r - k} \frac{G_{f^n}^n(\chi^\delta \psi^\theta)}{G_{f^n}^n(\psi^\theta)}.$$

Consider a fixed pair j, k . For each $a \in \{1, 2, \dots, n\}$, $G_{f^n}(\psi^\theta) = G_{f^n}(\psi^{\theta q^a})$, so

$$\prod_{c=1}^{w^r - k} G_{f^n}(\psi^\theta) = \prod_{c=1}^{w^r - k} G_{f^n}(\psi^{w^{k-1}(cw + i_j q^a)}).$$

Similarly,

$$\prod_{c=1}^{w^r - k} G_{f^n}(\chi^\delta \psi^\theta) = \prod_{c=1}^{w^r - k} G_{f^n}(\chi^\delta \psi^{w^{k-1}(cw + i_j q^a)}).$$

Thus

$$(9) \quad \eta^n = \frac{\chi^{ln}(l) G_{fn}(\chi^\delta)}{G_{fn}(\chi^{\delta l})} \prod_{j=1}^{l-1} \frac{G_{fn}(\chi^\delta \psi^j)}{G_{fn}(\psi^j)}.$$

Since $n \equiv \delta \pmod{q-1}$, $\chi^{ln}(l) = \chi^{\delta l}(l)$. Therefore, by (7), the right side of (9) equals 1, so

$$(10) \quad \eta^n = 1.$$

By the definition of η and of Gauss sums,

$$\eta^l \equiv \frac{\chi^{l^2}(l) \bar{\chi}^l(l) G_f(\chi^l)}{G_f^l(\chi^l)} \prod_{j=1}^e \prod_{k=1}^r \prod_{c=1}^{w^{r-k}} \frac{\bar{\chi}^{\delta l}(l) G_{fn}(\chi^{\delta l})}{1} \pmod{w},$$

so

$$\eta^l \equiv \frac{\chi^{l^2 - l - l\delta(l-1)/n}(l) G_{fn}^{(l-1)/n}(\chi^{\delta l})}{G_f^{l-1}(\chi^l)} \pmod{w}.$$

By (6), $G_{fn}(\chi^{\delta l}) = G_f^n(\chi^l)$; hence

$$(11) \quad \eta^l \equiv 1 \pmod{w}.$$

Thus w divides the norm $N(\eta^l - 1)$. By (10), η^l is an n -th root of unity. Thus if $\eta^l - 1 \neq 0$, then $N(\eta^l - 1)$ divides n , which contradicts the fact that $w \nmid n$. Therefore $\eta^l = 1 = \eta^n$, so since $(l, n) = 1$, $\eta = 1$.

5. PROOF OF (3)

Let η denote the right side of (3). We assume that $0 < \alpha < q - 1$. To see that this presents no loss of generality, we now show that η is unchanged when α is replaced by $\alpha + (q-1)j$, where j is an integer. Clearly $G_f(\chi^\alpha)$ and $\chi^\alpha(l)$ are unchanged, since the restriction $\chi|_{GF(q)}$ has order $q - 1$. Finally, $G_{fl}(\chi^{\alpha\beta})$ is also unchanged, as

$$(12) \quad G_{fl}(\chi^{\alpha\beta}) = G_{fl}(\chi^{\alpha\beta q^{\alpha j}}) = G_{fl}(\chi^{\beta(\alpha + j(q-1))}),$$

where α_j is defined by $\alpha_j \alpha \equiv j \pmod{l}$, $\alpha_j \geq 0$.

Let $\psi = \chi^{\beta(q-1)}$. Using (6), we have

$$\eta^l = \frac{G_{fl}(\chi^{\alpha\beta l})}{\chi^{\alpha l}(l) G_{fl}(\chi^{\alpha\beta})} \prod_{j=1}^{l-1} G_{fl}(\psi^j).$$