Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 28 (1982)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: RFDUCIBILITY BY ALGEBRAIC PROJECTIONS
Autor: Valiant, L. G.
Anhang: Appendix 2
DOI: https://doi.org/10.5169/seals-52240

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 08.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

$$T_{m+s+1}(\mathbf{x}, \mathbf{c})\, T_{m+s+1}(\mathbf{x}, \mathbf{c}')\, S_{m+r+1}(\mathbf{x}, \mathbf{e})$$

where $(\mathbf{c}, \mathbf{c}', \mathbf{e})$ is a fixed (0, 1)-vector of $2s+r+3$ elements. But any such vector can be specified by a conjunction of $2s+r+3$ Boolean literals. Consider the disjunction of the r such conjunctions and let $R(\mathbf{c}, \mathbf{c}', \mathbf{e})$ be the polynomial that simulates this Boolean formula at (0, 1) values. Then clearly

$$Q_m(x) = \sum T(\mathbf{x}, \mathbf{c})\, T(\mathbf{x}, \mathbf{c}')\, S(\mathbf{x}, \mathbf{e})\, R(\mathbf{c}, \mathbf{c}', \mathbf{e}) ,$$

where summation is over $(\mathbf{c}, \mathbf{c}', \mathbf{e}) \in \{0, 1\}^{2s+r+3}$.
Let $A(C, d)$ be the upper bound over every homogeneous polynomial having degree d and homogeneous program complexity C, of the minimal size of formula needed to define it in Definition 4. Then the above recursive expression ensures that

$$A(C, d) \leqslant 3A(3C+d, \lfloor d/2 \rfloor + 1) + 0(C) .$$

Clearly also $A(C, 1) \leqslant 2C$. Hence if d is p-bounded in m then so is the solution to this recurrence. □

Appendix 2

For completeness we describe here a direct proof of the p-definability of HC in the sense of Definition 1. $HC_{n \times n}(x_{i,j})$ will be the projection under

$$\sigma(u_{k,m}) = 1 \quad \text{for} \quad 1 \leqslant k, m \leqslant n$$

of the polynomial in $\{x_{i,j}, u_{k,m}\}$ defined by

$$Q_{n \times n}(y_{i,j}) \cdot Q_{n \times n}(z_{k,m}) \cdot R^1 \ldots R^n$$

with the association $y_{i,j} \leftrightarrow x_{i,j}$ and $z_{k,m} \leftrightarrow u_{k,m}$. Here $Q_{n \times n}$ is the polynomial that defines the permanent in §3. Its first occurrence with argument y plays exactly the same role as in the permanent and ensures a cycle cover. The intention of $z_{k,m}$ is to denote whether the kth node in the circuit (starting from node 1, say) is node m. $Q_{n \times n}(z_{k,m})$ ensures that this intention is realised. For each k R^k captures the fact that if $z_{k,m}$ and $z_{k+1,r}$ are both 1 then $y_{m,r}$ must be also. In Boolean notation we require

$$y_{m,r} \vee (\bar{z}_{k,m} \vee \bar{z}_{k+1,r}) .$$

As is well known such Boolean formulae can be simulated by polynomials at $\{0, 1\}$ values (e.g. see Proposition 2 in [13]). To guarantee just one monomial for each cycle we fix $R^1 = z_{11}$. □