

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	28 (1982)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
 Artikel:	ON THE NUMBER OF RESTRICTED PRIME FACTORS OF AN INTEGER. III
Autor:	Norton, Karl K.
Kapitel:	§4. Proofs of Theorem 1.14 and related results
DOI:	https://doi.org/10.5169/seals-52232

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 10.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

THEOREM 3.20. Suppose there exist real numbers $\delta > 0$, $x_0 \geq 2$ such that (3.14) holds. If $x \geq c_{29}(\delta)$ and $x_0 \leq y \leq \delta(\log x)(\log_2 x)^{-1}$, then

$$S(x, y; E, \omega) \gg x \exp \{-\delta^{-1}(y \log y + \log y + 2)\}.$$

Proof: In the notation of the preceding proof, (3.17) holds, and trivially $\log n_r \leq r \log p_r$. If $y \geq x_0$, then $p_r > r \geq x_0$ and $r = \pi(p_r; E) \geq p_r^\delta$, so

$$\begin{aligned} \log n_r &\leq \delta^{-1} r \log r \leq \delta^{-1} (y+1) (\log y + y^{-1}) \\ &\leq \delta^{-1} (y \log y + \log y + 2). \end{aligned} \quad (3.21)$$

But $\log y \leq \log_2 x - \log_3 x$, so $\log n_r < \log x$ if $x \geq c_{29}(\delta)$. Hence (3.19) holds, and the result follows from (3.17) and (3.21). Q.E.D.

§4. PROOFS OF THEOREM 1.14 AND RELATED RESULTS

We begin by quoting the following easy result from [13, pp. 689-690]:

LEMMA 4.1. For $x \geq 1$ and $z \geq 1$,

$$\sum_{n \leq x} z^{\omega(n; E)} \leq x \prod_{p \leq x, p \in E} \{1 + (z-1)p^{-1}\}.$$

To put this in a more convenient form, we prove

LEMMA 4.2. If $x \geq 1$ and $w \geq -2$, then (cf. (1.2))

$$\prod_{p \leq x, p \in E} (1 + wp^{-1}) \leq e^{wE(x)}. \quad (4.3)$$

If $1 \leq w \leq x$, then

$$\begin{aligned} &\prod_{p \leq x, p \in E} (1 + wp^{-1}) \\ &= \exp \{w(E(x) - E(w)) + O(w/\log(2w))\}. \end{aligned} \quad (4.4)$$

Proof: (4.3) follows immediately from the inequalities

$$0 \leq 1 + wp^{-1} \leq \exp(wp^{-1}).$$

To get (4.4), we first write

$$\begin{aligned} \prod_{p \leq x, p \in E} (1 + wp^{-1}) &\leq \prod_{p \leq w} (2wp^{-1}) \cdot \prod_{w < p \leq x, p \in E} \exp(wp^{-1}) \\ &= \exp \{w(E(x) - E(w)) + \pi(w) \log(2w) - \theta(w)\}, \end{aligned}$$

where $\pi(w) = \sum_{p \leq w} 1$ and $\theta(w) = \sum_{p \leq w} \log p$. Since $\pi(t) \ll t/\log(2t)$ for $t \geq 1$, we have

$$\begin{aligned}\theta(w) &= \int_1^w (\log t) d\pi(t) = \pi(w) \log w - \int_1^w \pi(t) t^{-1} dt \\ &= \pi(w) \log w + O(w/\log(2w)),\end{aligned}$$

and it follows that the right-hand side of (4.4) is an upper bound for the left-hand side. On the other hand, since $\log(1+y) = y + O(y^2)$ for $y > 0$, we have

$$\begin{aligned}\prod_{p \leq x, p \in E} (1+wp^{-1}) &\geq \prod_{w < p \leq x, p \in E} \exp \{wp^{-1} + O(w^2 p^{-2})\} \\ &= \exp \{w(E(x) - E(w)) + O(w^2 \sum_{p > w} p^{-2})\}.\end{aligned}$$

But

$$\sum_{p > w} p^{-2} = \int_w^{+\infty} t^{-2} d\pi(t) < 2 \int_w^{+\infty} t^{-3} \pi(t) dt \ll (w \log(2w))^{-1},$$

and (4.4) follows. Q.E.D.

COROLLARY 4.5. If $x \geq 1$ and $z \geq 1$, then

$$\sum_{n \leq x} z^{\omega(n; E)} \leq x e^{(z-1)E(x)}. \quad (4.6)$$

If $1 \leq z \leq x$, then

$$\begin{aligned}\sum_{n \leq x} z^{\omega(n; E)} \\ \leq x \exp \{(z-1)(E(x) - E(z)) + c_{30}z/\log(2z)\}.\end{aligned} \quad (4.7)$$

Note that if $1 \leq z < 2$, then (4.7) follows from (4.6).

THEOREM 4.8. Let $x \geq 1, v > 0, 1 \leq \alpha \leq x$. Define $\Lambda = \Lambda(x, v; E)$ by (1.22). Then

$$S(x, \alpha v; E, \omega) \leq x \exp \{(\alpha - 1 - \alpha \log \alpha)v - \alpha E(\alpha) + c_{31}\Lambda\alpha\}.$$

Proof: Suppose $1 \leq z \leq x$. Then

$$\sum_{n \leq x} z^{\omega(n; E)} \geq \sum_{n \leq x, \omega(n; E) > \alpha v} z^{\omega(n; E)} \geq z^{\alpha v} S(x, \alpha v; E, \omega).$$

Combining this result with (4.7), we get

$$\begin{aligned}S(x, \alpha v; E, \omega) &\leq x \exp \{(z-1)(v+\Lambda) - zE(z) - \alpha v \log z \\ &\quad + c_{32}z/\log(2z)\}.\end{aligned} \quad (4.9)$$

In practice, we think of v as being a good approximation to $E(x)$, so that Λ is small compared to v . We want to minimize the right-hand side of (4.9) approximately, and for simplicity, we choose z so as to minimize the expression $(z - 1)v - \alpha v \log z$, i.e., we take $z = \alpha$. With this value of z , we get the result from (4.9). Q.E.D.

LEMMA 4.10. *Suppose that there exists a real number $\gamma(E) > 0$ such that (1.7) holds. Then there is a real number $\delta(E)$ such that*

$$E(x) = \gamma(E) \log_2 x + \delta(E) + O_E(1/\log x) \quad \text{for } x \geq 2. \quad (4.11)$$

Proof : Write

$$E(x) = \int_1^x t^{-1} d\pi(t; E),$$

integrate by parts, and use (1.7). Q.E.D.

From Theorem 4.8 and Lemma 4.10, we get

COROLLARY 4.12. *Suppose that there exists a real number $\gamma(E) > 0$ such that (1.7) holds. Let $x \geq 3$, $2 \leq \alpha \leq x$. Then*

$$\begin{aligned} & S(x, \alpha \gamma(E) \log_2 x; E, \omega) \\ & \leq x \exp \{(\alpha - 1 - \alpha \log \alpha) \gamma(E) \log_2 x - \alpha \gamma(E) \log_2 \alpha + c_{33}(E) \alpha\}. \end{aligned}$$

Using (1.8), it is easy to show that Corollary 4.12 actually holds for all $\alpha \geq 2$, but it is also clear from (1.8) that

$$S(x, \alpha \gamma(E) \log_2 x; E, \omega) = 0$$

whenever α is somewhat greater than $(\log x)(\log_2 x)^{-2}$.

The upper bound given in Corollary 4.12 compares favorably with the theorem of Delange (Theorem 1.17 above), and our result is more general and holds for a much wider range of α . Our proof is also much simpler than Delange's. Unfortunately, our lower bound (1.13) is much smaller than the upper bound in Corollary 4.12.

Theorem 1.14 is proved in the same way as Theorem 4.8, but we use (4.6) instead of (4.7), apply Lemma 4.10, and take $z = y(\gamma(E) \log_2 x)^{-1}$.

We conclude this section by generalizing the Erdős-Nicolas result (1.10).

THEOREM 4.13. Suppose that there exists a real number $\gamma(E) > 0$ such that (1.7) holds. Let $\varepsilon > 0$, and suppose that $x \geq c_{34}(E, \varepsilon)$ and

$$(\log_2 x)^2 (\log x)^{-1} \leq \alpha \leq 1 + \{1 + \log \gamma(E) - \varepsilon\} (\log_2 x)^{-1}.$$

Then

$$\begin{aligned} x^{1-\alpha} \exp \left\{ -c_{35}(E) \frac{\log x}{\log_2 x} \right\} &\leq S(x, \alpha (\log x) (\log_2 x)^{-1}; E, \omega) \\ &\leq x^{1-\alpha} \exp \left\{ \frac{2\alpha (\log x) \log_3 x}{\log_2 x} + c_{36}(E) \frac{\log x}{\log_2 x} \right\}. \end{aligned}$$

This can be obtained from Theorems 1.11 and 1.14 (take

$$y = \alpha (\log x) (\log_2 x)^{-1}$$

and use the inequalities

$$\log_2 y \leq \log_3 x, y \geq \log_2 x \geq \gamma(E) \log_2 x.$$

Theorem 4.13 should be compared with Theorem 1.6.

§5. PROOFS OF THEOREM 1.21 AND RELATED RESULTS

In estimating $S(x, y; E, \Omega)$ (defined by (1.1)), we do not need any assumption such as (1.7). Hence we emphasize that throughout the remainder of this paper, E is merely assumed to be any nonempty set of primes. (We shall sometimes assume explicitly that E has at least two members.) The smallest member of E will always be denoted by p_1 (and the smallest member of $E - \{p_1\}$, if it exists, by p_2). When x and v are positive real numbers, the function $\Lambda = \Lambda(x, v; E)$ is always defined by (1.22).

The subsequent work depends heavily on the following elementary lemma [13, p. 690]:

LEMMA 5.1. If $x > 0$ and $1 \leq z < p_1$, then

$$\sum_{n \leq x} z^{\Omega(n; E)} < p_1 (p_1 - z)^{-1} x e^{(z-1)E(x) + 4z}.$$