

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 28 (1982)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE REPRESENTATION THEORY OF $SL(2, \mathbb{R})$, A NON-INFINITESIMAL APPROACH
Autor: Koornwinder, Tom H.
Kapitel: 2.2. The principal series
DOI: <https://doi.org/10.5169/seals-52233>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

2.2. THE PRINCIPAL SERIES

Let G be a connected noncompact real semisimple Lie group with finite center. Let $G = KAN$ be an Iwasawa decomposition. For $g \in G$ write $g = u(g)\exp(H(g))n(g)$, where $u(g) \in K$, $H(g) \in \mathfrak{a}$ (the Lie algebra of A) and $n(g) \in N$. Let $\rho \in \mathfrak{a}^*$ be half the sum of the positive roots. Let M be the centralizer of A in K . For $\xi \in \hat{M}$, $\lambda \in \mathfrak{a}_\mathbb{C}^*$ the *principal series* representation $\pi_{\xi, \lambda}$ of G is obtained by inducing the (not necessarily unitary) finite-dimensional irreducible representation $man \rightarrow e^{\lambda(\log a)}\xi(m)$ of the subgroup MAN . In the so-called compact picture we have the following realization of $\pi_{\xi, \lambda}$ (cf. WALLACH [45, §8.3]):

$$(2.2) \quad (\pi_{\xi, \lambda}(g)f)(k) = e^{-(\rho+\lambda)(H(g^{-1}k))} f(u(g^{-1}k)), \\ f \in L^2_\xi(K, \mathcal{H}(\xi)), \quad k \in K, g \in G.$$

Here the Hilbert space $L^2_\xi(K, \mathcal{H}(\xi))$ consists of all $\mathcal{H}(\xi)$ -valued L^2 -functions f on K such that $f(km) = \xi(m^{-1})f(k)$, $k \in K$, $m \in M$. The representation $\pi_{\xi, \lambda}$ is a K -unitary Hilbert representation. It is unitary if $\lambda \in ia^*$. By Frobenius reciprocity, $\pi_{\xi, \lambda}$ is K -finite and $\pi_{\xi, \lambda}$ is K -multiplicity free if each $\delta \in \hat{K}$ is M -multiplicity free.

Let us now specialize the above results to $G = SL(2, \mathbf{R})$. It is convenient to work with the group $G = SU(1, 1)$, isomorphic to $SL(2, \mathbf{R})$:

$$(2.3) \quad G := \left\{ g_{\alpha, \beta} = \begin{pmatrix} \alpha & \beta \\ \bar{\beta} & \bar{\alpha} \end{pmatrix}; \quad \alpha, \beta \in \mathbf{C}, |\alpha|^2 - |\beta|^2 = 1 \right\}.$$

Let

$$(2.4) \quad K := \left\{ u_\theta = \begin{pmatrix} e^{\frac{1}{2}i\theta} & 0 \\ 0 & e^{-\frac{1}{2}i\theta} \end{pmatrix}; \quad 0 \leq \theta < 4\pi \right\},$$

$$(2.5) \quad A := \left\{ a_t = \begin{pmatrix} ch_{\frac{1}{2}t} & sh_{\frac{1}{2}t} \\ sh_{\frac{1}{2}t} & ch_{\frac{1}{2}t} \end{pmatrix} = \exp t \begin{pmatrix} 0 & \frac{1}{2} \\ \frac{1}{2} & 0 \end{pmatrix}; \quad t \in \mathbf{R} \right\},$$

$$(2.6) \quad N := \left\{ n_z = \begin{pmatrix} 1 + \frac{1}{2}iz & -\frac{1}{2}iz \\ \frac{1}{2}iz & 1 - \frac{1}{2}iz \end{pmatrix}; \quad z \in \mathbf{R} \right\}.$$

Then $G = KAN$ is an Iwasawa decomposition for $G = SU(1, 1)$, $\rho(\log a_t) = \frac{1}{2}t$ and $M = \{u_0, u_{2\pi}\}$. \hat{M} consists of the two one-dimensional representations

$$(2.7) \quad u_\theta \rightarrow e^{i\xi\theta}, \quad u_\theta \in M, \xi = 0 \text{ or } \frac{1}{2}.$$

Let $L_\xi^2(K)$ consist of all $f \in L^2(K)$ such that $f(u_\psi + 2\pi) = f(u_\psi)$ or $-f(u_\psi)$ according to whether $\xi = 0$ or $\frac{1}{2}$, respectively.

Now, by using explicit expressions for the factors in the Iwasawa decomposition of $g_{\alpha, \beta}^{-1} u_\psi$ (cf. TAKAHASHI [39, §1]) we can write (2.2) in the case $G = U(1, 1)$ as follows:

$$(2.8) \quad (\pi_{\xi, \lambda}(g_{\alpha, \beta})f)(u_\psi) := |\bar{\alpha}e^{\frac{1}{2}i\psi} - \beta e^{-\frac{1}{2}i\psi}|^{-2\lambda-1} f(u_\psi),$$

$$\psi' := 2 \arg(\bar{\alpha}e^{\frac{1}{2}i\psi} - \beta e^{-\frac{1}{2}i\psi}), \quad g_{\alpha, \beta} \in G, u_\psi \in K, f \in L_\xi^2(K),$$

$$\xi = 0 \text{ or } \frac{1}{2}, \lambda \in \mathbf{C}.$$

On putting $g_{\alpha, \beta} := u_\theta \in K$ we get

$$(2.9) \quad (\pi_{\xi, \lambda}(u_\theta)f)(u_\psi) = f(u_\psi - \theta), \quad f \in L_\xi^2(K), u_\theta, u_\psi \in K,$$

which again shows that $\pi_{\xi, \lambda}$ is K -unitary. \hat{K} consists of the representations

$$(2.10) \quad \delta_n(u_\theta) := e^{in\theta}, \quad u_\theta \in K,$$

where n runs through the set $\frac{1}{2}\mathbf{Z}$, i.e., $2n \in \mathbf{Z}$. An orthogonal basis for $L_\xi^2(K)$ is given by the functions

$$(2.11) \quad \phi_n(u_\psi) := e^{-in\psi}, \quad u_\psi \in K,$$

where n runs through the set $\mathbf{Z} + \xi := \{m + \xi \mid m \in \mathbf{Z}\}$. Then

$$(2.12) \quad \pi_{\xi, \lambda}(u_\theta)\phi_n = \delta_n(u_\theta)\phi_n, \quad u_\theta \in K, n \in \mathbf{Z} + \xi.$$

Thus $\pi_{\xi, \lambda}$ is K -multiplicity free,

$$(2.13) \quad \mathcal{M}(\pi_{\xi, \lambda}) = \{\delta_n \in \hat{K} \mid n \in \mathbf{Z} + \xi\},$$

the ϕ_n 's form a K -basis for $L_\xi^2(K)$ and the canonical matrix elements of $\pi_{\xi, \lambda}$ are

$$(2.14) \quad \pi_{\xi, \lambda, m, n}(g) = (\pi_{\xi, \lambda}(g)\phi_n, \phi_m), \quad g \in G, m, n \in \mathbf{Z} + \xi.$$

Because of the Cartan decomposition $G = KAK$, $\pi_{\xi, \lambda, m, n}$ is completely determined by its restriction to A . It follows from (2.8) and (2.11) that

$$(\pi_{\xi, \lambda}(a_t)\phi_n)(u_\psi) = |ch_{\frac{1}{2}t} e^{\frac{1}{2}i\psi} - sh_{\frac{1}{2}t} e^{-\frac{1}{2}i\psi}|^{-2\lambda+2n-1}$$

$$\cdot (ch_{\frac{1}{2}t} e^{\frac{1}{2}i\psi} - sh_{\frac{1}{2}t} e^{-\frac{1}{2}i\psi})^{-2n}.$$

Hence

$$(2.15) \quad \pi_{\xi, \lambda, m, n}(a_t) = (ch\frac{1}{2}t)^{-2\lambda-1} \\ \cdot \frac{1}{4\pi} \int_0^{4\pi} (1 - th\frac{1}{2}t e^{i\psi})^{-\lambda+n-1/2} (1 - th\frac{1}{2}t e^{-i\psi})^{-\lambda-n-1/2} e^{i(m-n)\psi} d\psi, \\ t \in \mathbf{R}, m, n \in \mathbf{Z} + \xi.$$

The following symmetry is evident from (2.15):

$$(2.16) \quad \pi_{\xi, \lambda, -m, -n}(a_t) = \pi_{\xi, \lambda, m, n}(a_t).$$

2.3. CALCULATION OF THE CANONICAL MATRIX ELEMENTS

Let us calculate the integral (2.15). In view of (2.16) we can suppose $m \geq n$. The binomial expansion

$$(2.17) \quad (1-z)^{-a} = \sum_{k=0}^{\infty} \frac{(a)_k}{k!} z^k, \quad |z| < 1, a \in \mathbf{C},$$

where

$$(2.18) \quad (a)_k := a(a+1)\dots(a+k-1) = \frac{\Gamma(a+k)}{\Gamma(a)},$$

can be substituted for the first two factors in the integrand of (2.15). Now interchange the order of summation and integration and perform the integration in each term. Then we obtain ($m \geq n$)

$$(2.19) \quad \pi_{\xi, \lambda, m, n}(a_t) = \frac{(\lambda + n + \frac{1}{2})_{m-n}}{(m-n)!} (sh\frac{1}{2}t)^{m-n} (ch\frac{1}{2}t)^{n-m-2\lambda-1} \\ \cdot {}_2F_1(\lambda + m + \frac{1}{2}, \lambda - n + \frac{1}{2}; m - n + 1; (th\frac{1}{2}t)^2),$$

where the ${}_2F_1$ denotes a *hypergeometric series*, defined by

$$(2.20) \quad {}_2F_1(a, b; c; z) := \sum_{k=0}^{\infty} \frac{(a)_k (b)_k}{(c)_k k!} z^k, \quad |z| < 1, a, b, c \in \mathbf{C},$$

cf. [10, Vol. I, Ch. 2].

The expression (2.20) is clearly symmetric in a and b . As a function of z , the ${}_2F_1$ has an analytic continuation to a one-valued function on $\mathbf{C} \setminus [1, \infty)$. Application of the transformation formulas