

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 28 (1982)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE REPRESENTATION THEORY OF $SL(2, \mathbb{R})$, A NON-INFINITESIMAL APPROACH
Autor: Koornwinder, Tom H.
Kapitel: 2.3. CALCULATION OF THE CANONICAL MATRIX ELEMENTS
DOI: <https://doi.org/10.5169/seals-52233>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Hence

$$(2.15) \quad \pi_{\xi, \lambda, m, n}(a_t) = (ch\frac{1}{2}t)^{-2\lambda-1} \\ \cdot \frac{1}{4\pi} \int_0^{4\pi} (1 - th\frac{1}{2}t e^{i\psi})^{-\lambda+n-1/2} (1 - th\frac{1}{2}t e^{-i\psi})^{-\lambda-n-1/2} e^{i(m-n)\psi} d\psi, \\ t \in \mathbf{R}, m, n \in \mathbf{Z} + \xi.$$

The following symmetry is evident from (2.15):

$$(2.16) \quad \pi_{\xi, \lambda, -m, -n}(a_t) = \pi_{\xi, \lambda, m, n}(a_t).$$

2.3. CALCULATION OF THE CANONICAL MATRIX ELEMENTS

Let us calculate the integral (2.15). In view of (2.16) we can suppose $m \geq n$. The binomial expansion

$$(2.17) \quad (1-z)^{-a} = \sum_{k=0}^{\infty} \frac{(a)_k}{k!} z^k, \quad |z| < 1, a \in \mathbf{C},$$

where

$$(2.18) \quad (a)_k := a(a+1)\dots(a+k-1) = \frac{\Gamma(a+k)}{\Gamma(a)},$$

can be substituted for the first two factors in the integrand of (2.15). Now interchange the order of summation and integration and perform the integration in each term. Then we obtain ($m \geq n$)

$$(2.19) \quad \pi_{\xi, \lambda, m, n}(a_t) = \frac{(\lambda + n + \frac{1}{2})_{m-n}}{(m-n)!} (sh\frac{1}{2}t)^{m-n} (ch\frac{1}{2}t)^{n-m-2\lambda-1} \\ \cdot {}_2F_1(\lambda + m + \frac{1}{2}, \lambda - n + \frac{1}{2}; m - n + 1; (th\frac{1}{2}t)^2),$$

where the ${}_2F_1$ denotes a *hypergeometric series*, defined by

$$(2.20) \quad {}_2F_1(a, b; c; z) := \sum_{k=0}^{\infty} \frac{(a)_k (b)_k}{(c)_k k!} z^k, \quad |z| < 1, a, b, c \in \mathbf{C},$$

cf. [10, Vol. I, Ch. 2].

The expression (2.20) is clearly symmetric in a and b . As a function of z , the ${}_2F_1$ has an analytic continuation to a one-valued function on $\mathbf{C} \setminus [1, \infty)$. Application of the transformation formulas

$$(2.21) \quad {}_2F_1(a, b; c; z) = (1-z)^{-b} {}_2F_1\left(c-a, b; c; \frac{z}{z-1}\right) \\ = (1-z)^{-a} {}_2F_1\left(a, c-b; c; \frac{z}{z-1}\right)$$

(cf. [10, Vol. I, §2.1 (22)]) to (2.19) yields ($m \geq n$):

$$(2.22) \quad \pi_{\xi, \lambda, m, n}(a_t) \\ = \frac{(\lambda + n + \frac{1}{2})_{m-n}}{(m-n)!} (sh\frac{1}{2}t)^{m-n} (ch\frac{1}{2}t)^{-m-n} {}_2F_1(\lambda - n + \frac{1}{2}, -\lambda - n + \frac{1}{2}; m-n+1; -(sh\frac{1}{2}t)^2) \\ = \frac{(\lambda + n + \frac{1}{2})_{m-n}}{(m-n)!} (sh\frac{1}{2}t)^{m-n} (ch\frac{1}{2}t)^{m+n} {}_2F_1(\lambda + m + \frac{1}{2}, -\lambda + m + \frac{1}{2}; m-n+1; -(sh\frac{1}{2}t)^2).$$

It is more elegant to express the hypergeometric functions in (2.22) in terms of *Jacobi functions* $\phi_{\mu}^{(\alpha, \beta)}$ ($\mu, \alpha, \beta \in \mathbf{C}, \alpha \notin \{-1, -2, \dots\}$), which are defined on \mathbf{R} by

$$(2.23) \quad \phi_{\mu}^{(\alpha, \beta)}(t) \\ := {}_2F_1\left(\frac{1}{2}(\alpha + \beta + 1 + i\mu), \frac{1}{2}(\alpha + \beta + 1 - i\mu); \alpha + 1; -(sht)^2\right)$$

(cf. KOORNWINDER [36, §2]). Clearly,

$$(2.24) \quad \phi_{\mu}^{(\alpha, \beta)}(0) = 1,$$

$$(2.25) \quad \phi_{\mu}^{(\alpha, \beta)}(t) = \phi_{\mu}^{(\alpha, \beta)}(-t),$$

$$(2.26) \quad \phi_{\mu}^{(\alpha, \beta)}(t) = \phi_{-\mu}^{(\alpha, \beta)}(t).$$

The function $\phi_{\mu}^{(\alpha, \beta)}$ satisfies the differential equation

$$(2.27) \quad (\Delta_{\alpha, \beta}(t))^{-1} \frac{d}{dt} \left(\Delta_{\alpha, \beta}(t) \frac{du(t)}{dt} \right) \\ = -(\mu^2 + (\alpha + \beta + 1)^2)u(t),$$

where

$$\Delta_{\alpha, \beta}(t) := (sht)^{2\alpha+1} (cht)^{2\beta+1},$$

and $u := \phi_{\mu}^{\alpha, \beta}$ is the unique solution of (2.27) which is regular at $t = 0$ and satisfies $u(0) = 1$. For fixed $\alpha > -1, \beta \in \mathbf{R}$, Jacobi functions $\phi_{\mu}^{(\alpha, \beta)}$ form a continuous orthogonal system with respect to the measure $\Delta_{\alpha, \beta}(t)dt, t > 0$.

Substitution of (2.23) and (2.22) yields ($m \geq n$):

$$\begin{aligned}
 (2.28) \quad & \pi_{\xi, \lambda, m, n}(a_t) \\
 &= \frac{(\lambda + n + \frac{1}{2})_{m-n}}{(m-n)!} (sh \frac{1}{2}t)^{m-n} (ch \frac{1}{2}t)^{-m-n} \phi_{2i\lambda}^{(m-n, -m-n)}(\frac{1}{2}t) \\
 &= \frac{(\lambda + n + \frac{1}{2})_{m-n}}{(m-n)!} (sh \frac{1}{2}t)^{m-n} (ch \frac{1}{2}t)^{m+n} \phi_{2i\lambda}^{(m-n, m+n)}(\frac{1}{2}t).
 \end{aligned}$$

Application of (2.16) gives a similar result in the case $m < n$. Finally we conclude:

THEOREM 2.1. *The canonical matrix elements $\pi_{\xi, \lambda, m, n}(a_t)$ ($\lambda \in \mathbf{C}$; $\xi = 0$ or $\frac{1}{2}$; $m, n \in \mathbf{Z} + \xi$; $t \in \mathbf{R}$) of $SU(1, 1)$ can be expressed in terms of Jacobi functions by*

$$(2.29) \quad \pi_{\xi, \lambda, m, n}(a_t) = \frac{c_{\xi, \lambda, m, n}}{(|m-n|)!} (sh \frac{1}{2}t)^{|m-n|} (ch \frac{1}{2}t)^{m+n} \phi_{2i\lambda}^{(|m-n|, m+n)}(\frac{1}{2}t),$$

where

$$(2.30) \quad c_{\xi, \lambda, m, n} := \begin{cases} (\lambda + n + \frac{1}{2})_{m-n} & \text{if } m \geq n, \\ (\lambda - n + \frac{1}{2})_{n-m} & \text{if } n \geq m. \end{cases}$$

In view of (2.24), formulas (2.29) and (2.30) describe the asymptotics of $\pi_{\xi, \lambda, m, n}$ near $t = 0$.

2.4. NOTES

2.4.1. The principal series of representations was first written down for $SL(2, \mathbf{R})$ by BARGMANN [2], for $SL(2, \mathbf{C})$ by GELFAND & NAIMARK [18], and for a general noncompact semisimple Lie group by HARISH-CHANDRA [21, §12].

2.4.2. BARGMANN [2, §10] already obtained explicit expressions in terms of hypergeometric functions for the canonical matrix elements of the irreducible unitary representations of $SL(2, \mathbf{R})$. He solved the differential equation satisfied by these matrix elements, which is obtained from the Casimir operator. VILENKIN [43, Ch. VI, §3] gives a derivation of these expressions which is similar to our derivation in §2.4, starting from the integral representation (2.15).

2.4.3. It follows from the present paper that the spherical functions for $SL(2, \mathbf{R})$ can be expressed as Jacobi functions of order $(\alpha, \beta) = (0, 0)$. More generally, the spherical functions on any noncompact real semisimple Lie group