

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 28 (1982)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE REPRESENTATION THEORY OF $SL(2, \mathbb{R})$, A NON-INFINITESIMAL APPROACH
Autor: Koornwinder, Tom H.
Kapitel: 2.4. Notes
DOI: <https://doi.org/10.5169/seals-52233>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 10.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Substitution of (2.23) and (2.22) yields ($m \geq n$):

$$(2.28) \quad \begin{aligned} & \pi_{\xi, \lambda, m, n}(a_t) \\ &= \frac{(\lambda + n + \frac{1}{2})_{m-n}}{(m-n)!} (sh \frac{1}{2}t)^{m-n} (ch \frac{1}{2}t)^{-m-n} \phi_{2i\lambda}^{(m-n, -m-n)} (\frac{1}{2}t) \\ &= \frac{(\lambda + n + \frac{1}{2})_{m-n}}{(m-n)!} (sh \frac{1}{2}t)^{m-n} (ch \frac{1}{2}t)^{m+n} \phi_{2i\lambda}^{(m-n, m+n)} (\frac{1}{2}t). \end{aligned}$$

Application of (2.16) gives a similar result in the case $m < n$. Finally we conclude:

THEOREM 2.1. *The canonical matrix elements $\pi_{\xi, \lambda, m, n}(a_t)$ ($\lambda \in \mathbf{C}$; $\xi = 0$ or $\frac{1}{2}$; $m, n \in \mathbf{Z} + \xi$; $t \in \mathbf{R}$) of $SU(1, 1)$ can be expressed in terms of Jacobi functions by*

$$(2.29) \quad \pi_{\xi, \lambda, m, n}(a_t) = \frac{c_{\xi, \lambda, m, n}}{(|m-n|)!} (sh \frac{1}{2}t)^{|m-n|} (ch \frac{1}{2}t)^{m+n} \phi_{2i\lambda}^{(|m-n|, m+n)} (\frac{1}{2}t),$$

where

$$(2.30) \quad c_{\xi, \lambda, m, n} := \begin{cases} (\lambda + n + \frac{1}{2})_{m-n} & \text{if } m \geq n, \\ (\lambda - n + \frac{1}{2})_{n-m} & \text{if } n \geq m. \end{cases}$$

In view of (2.24), formulas (2.29) and (2.30) describe the asymptotics of $\pi_{\xi, \lambda, m, n}$ near $t = 0$.

2.4. NOTES

2.4.1. The principal series of representations was first written down for $SL(2, \mathbf{R})$ by BARGMANN [2], for $SL(2, \mathbf{C})$ by GELFAND & NAIMARK [18], and for a general noncompact semisimple Lie group by HARISH-CHANDRA [21, §12].

2.4.2. BARGMANN [2, §10] already obtained explicit expressions in terms of hypergeometric functions for the canonical matrix elements of the irreducible unitary representations of $SL(2, \mathbf{R})$. He solved the differential equation satisfied by these matrix elements, which is obtained from the Casimir operator. VILENKO [43, Ch. VI, §3] gives a derivation of these expressions which is similar to our derivation in §2.4, starting from the integral representation (2.15).

2.4.3. It follows from the present paper that the spherical functions for $SL(2, \mathbf{R})$ can be expressed as Jacobi functions of order $(\alpha, \beta) = (0, 0)$. More generally, the spherical functions on any noncompact real semisimple Lie group

of rank 1 (i.e., $\dim(A) = 1$) can be written as Jacobi functions of certain order (cf. HARISH-CHANDRA [23, §13]). This motivated FLENSTED-JENSEN [14] to study harmonic analysis for Jacobi function expansions of quite general order (α, β) , $\alpha \geq \beta \geq -\frac{1}{2}$. This research was continued in several papers by Flensted-Jensen and the author.

3. THE IRREDUCIBLE SUBQUOTIENT REPRESENTATIONS OF THE PRINCIPAL SERIES

3.1. SUBQUOTIENT REPRESENTATIONS

We start with the definition and some general properties and next derive an irreducibility criterium (Theorem 3.2) and a decomposition theorem 3.3.

Let G be a lcsc. group and let τ be a Hilbert representation of G . Let \mathcal{H}_0 be a closed subspace of $\mathcal{H}(\tau)$ and let P_0 be the orthogonal projection from $\mathcal{H}(\tau)$ onto \mathcal{H}_0 . Define

$$(3.1) \quad \tau_0(g)v := P_0\tau(g)v, \quad g \in G, v \in \mathcal{H}_0.$$

Then $\tau(g) \in \mathcal{L}(\mathcal{H}_0)$ for each $g \in G$, $\tau_0(e) = id.$, and $g \rightarrow \tau_0(g)v : G \rightarrow \mathcal{H}_0$ is continuous for each $v \in \mathcal{H}_0$. If also

$$(3.2) \quad \tau_0(g_1g_2) = \tau_0(g_1)\tau_0(g_2), \quad g_1, g_2 \in G,$$

then τ_0 is a Hilbert representation of G on \mathcal{H}_0 and it is called a *subquotient representation* of τ . Formula (3.2) is clearly valid if \mathcal{H}_0 is an *invariant subspace* of $\mathcal{H}(\tau)$, i.e., if $\tau(g)v \in \mathcal{H}_0$ for all $g \in G$, $v \in \mathcal{H}_0$. In that case, τ_0 is called a *subrepresentation* of τ .

LEMMA 3.1. *Let \mathcal{H}_0 be a closed subspace of $\mathcal{H}(\tau)$, let \mathcal{H}_2 be the closed G -invariant subspace of $\mathcal{H}(\tau)$ which is generated by \mathcal{H}_0 and let $\mathcal{H}_1 := \mathcal{H}_2 \cap \mathcal{H}_0^\perp$. Then τ_0 is a subquotient representation if and only if \mathcal{H}_1 is G -invariant.*