

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 28 (1982)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE REPRESENTATION THEORY OF $SL(2, \mathbb{R})$, A NON-INFINITESIMAL APPROACH
Autor: Koornwinder, Tom H.
Kapitel: 3.3. Notes
DOI: <https://doi.org/10.5169/seals-52233>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 10.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Thus $c_{\xi, \lambda, m, n}$ has block matrix

$$\begin{array}{ccc} n \leq -\lambda - \frac{1}{2} & -\lambda + \frac{1}{2} \leq n \leq \lambda - \frac{1}{2} & n \geq \lambda + \frac{1}{2} \\ \begin{matrix} m \leq -\lambda - \frac{1}{2} \\ -\lambda + \frac{1}{2} \leq m \leq \lambda - \frac{1}{2} \\ m \geq \lambda + \frac{1}{2} \end{matrix} & \left(\begin{array}{ccc} * & * & 0 \\ 0 & * & 0 \\ 0 & * & * \end{array} \right) & \end{array}$$

where each starred block has all entries nonzero.

- (d) $c_{\xi, \lambda, m, n} \neq 0 \Leftrightarrow \lambda + \frac{1}{2} \leq m \leq -\lambda - \frac{1}{2}$ or $m, n \leq \lambda - \frac{1}{2}$
or $m, n \geq \lambda + \frac{1}{2}$. □

The finite-dimensional representation occurring in the above classification are the representations $\pi_{\xi, \lambda}^0 (\lambda + \xi \in \mathbf{Z} + \frac{1}{2}, \lambda \neq 0)$.

3.3. NOTES

3.3.1. In the case of the unitary principal series (λ imaginary), Theorem 3.4 was first proved by BARGMANN [2, sections 6 and 7]. See van DIJK [9, Theorem 4.1] for the statement and (infinitesimal) proof of our Theorem 3.4 in the general case. A proof of Theorem 3.4 similar to our proof was earlier given by BARUT & PHILLIPS [3, §II (4)].

3.3.2. Theorem 3.4 in the case of imaginary and nonzero λ is contained in a general theorem by BRUHAT [5, Theorem 7; 2]: For $\xi \in \hat{M}$, $\lambda \in i\mathbf{a}$, the principal series representation $\pi_{\xi, \lambda}$ of G (cf. (2.2)) is irreducible if $s \cdot \lambda \neq \lambda$ for all $s \neq e$ in the Weyl group for (G, K) .

3.3.3. GELFAND & NAIMARK [18, §5.4, Theorem 1] proved the irreducibility of the unitary principal series for $SL(2, \mathbf{C})$ by a global method different from ours, working in a noncompact realization and calculating the “matrix elements” of the representation with respect to a (continuous) \overline{N} -basis.

3.3.4. Analogues of Theorems 3.2 and 3.3 can be formulated in the case of non-abelian K , cf. [27, Theorem 3.3]. In that case the canonical matrix elements $\tau_{\gamma, \delta}$ are matrix-valued functions. By using this method, NAIMARK [34, Ch. 3, §9, No. 15] examined the irreducibility of the nonunitary principal series for $SL(2, \mathbf{C})$, see also KOSTERS [28].

3.3.5. Further applications of the irreducibility criterium in Theorem 3.2 can be found in MILLER [32, Lemmas 3.2 and 4.5] for the Euclidean motion group of \mathbf{R}^2 and for the harmonic oscillator group, TAKAHASHI [39, §3.4] for the discrete series of $SL(2, \mathbf{R})$ and [41, p. 560, Cor. 2] for the spherical principal series of $F_{4(-20)}$.

3.3.6. The method of this section does not show in an *a priori* way that a K -multiplicity free principal series representation has only finitely many irreducible subquotient representations. Actually, this property holds quite generally, cf. WALLACH [45, Theorem 8.13.3].

4. EQUIVALENCES BETWEEN IRREDUCIBLE SUBQUOTIENT REPRESENTATIONS OF THE PRINCIPAL SERIES

4.1. NAIMARK EQUIVALENCE

In this subsection we derive a criterium (Theorem 4.5) for Naimark equivalence of K -multiplicity free representations. Lemmas 4.3 and 4.4 are preparations for its proof.

Let G be an lcsc. group.

Definition 4.1. Let σ and τ be Hilbert representations of G . The representation σ is called *Naimark related* to τ if there is a closed (possibly unbounded) injective linear operator A from $\mathcal{H}(\sigma)$ to $\mathcal{H}(\tau)$ with domain $\mathcal{D}(A)$ dense in $\mathcal{H}(\sigma)$ and range $\mathcal{R}(A)$ dense in $\mathcal{H}(\tau)$ such that $\mathcal{D}(A)$ is σ -invariant and $A\sigma(g)v = \tau(G)Av$ for all $v \in \mathcal{D}(A)$, $g \in G$. Then we use the notation $\sigma \simeq \tau$ or $\overset{A}{\sigma} \simeq \tau$.

Naimark relatedness is not necessarily a transitive relation (cf. WARNER [48, p. 242]). However, we will see that it becomes an equivalence relation (called *Naimark equivalence*) when restricted to the class of unitary representations or of K -multiplicity free representations, K abelian.

Two unitary representations σ and τ of G are called *unitarily equivalent* if there is an isometry A from $\mathcal{H}(\sigma)$ onto $\mathcal{H}(\tau)$ such that $A\sigma(g)v = \tau(g)Av$ for all $v \in \mathcal{H}(\sigma)$, $g \in G$. Clearly unitary equivalence is an equivalence relation.