Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 28 (1982)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE REPRESENTATION THEORY OF SL(2, R), A NON-INFINITESIMAL APPROACH

Autor: Koornwinder, Tom H.

Kapitel: 5.2. Spherical functions of type

DOI: https://doi.org/10.5169/seals-52233

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 10.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

5.2. Spherical functions of type δ

Let G be a unimodular lcsc. group with compact subgroup K. Let

$$K^* := \{(k, k) \in G \times K \mid k \in K\}.$$

Let $\delta \in \hat{K}$ and let τ be a K-unitary representation of G. Then $\tau \otimes \check{\delta}$ ($\check{\delta}$ the contragredient representation to δ) is a K^*-unitary representation of $G \times K$ on $\mathscr{H}(\tau) \otimes \mathscr{H}(\delta)$.

Lemma 5.1. *The multiplicity of* δ *in* $\tau|_K$ *is equal to the multiplicity of the representation* 1 *of* K^* *in* $\tau \otimes \check{\delta}|_{K^*}$. τ *is irreducible iff* $\tau \otimes \check{\delta}$ *is irreducible.* τ *is unitary iff* $\tau \otimes \check{\delta}$ *is unitary.*

This can be proved immediately. By using the results summarized in §5.1 we conclude that $(G \times K, K^*)$ is a Gelfand pair if there exists a continuous involutive homomorphism α on G such that for each $(g, k) \in G \times K$ we have $\alpha(g) = k_1 g^{-1} k_2$, $\alpha(k) = k_1 k^{-1} k_2$ for certain $k_1, k_2 \in K$. Furthermore, if $(G \times K, K^*)$ is a Gelfand pair and if the irreducible representation τ of G is unitary or K-finite then τ is K-multiplicity free. In particular, this applies to $SU(1, 1)$:

Proposition 5.2. *If* $G = SU(1, 1)$ *then* $(G \times K, K^*)$ *is a Gelfand pair.*

Proof. For $g \in SU(1, 1)$ define $\alpha(g) := {}^t(g^{-1})$. Then α is a continuous involutive automorphism on G and $\alpha(a_t) = a_{-t}$ on A, $\alpha(u_\theta) = u_{-\theta}$ on K. Since $G = KAK$, α has the required properties. □

Let $(G \times K, K^*)$ be a Gelfand pair. Identify $G \times \{e\}$ with G. A spherical function on $G \times K$ is completely determined by its restriction to G. By using the results mentioned in §5.1 we obtain the following properties. First, a continuous function ϕ on G is the restriction to G of a spherical function on $G \times K$ iff $\phi \neq 0$ and

$$\phi(x)\phi(y) = \int_K \phi(xkyk^{-1})dk, \qquad x, y \in G.$$

Next, let

$$\begin{aligned} &I_c(G) \ (\text{or } I_c^\infty(G)) \\ &:= \{f \in C_c(G) \ (\text{or } C_c^\infty(G)) \mid f(kgk^{-1}) = f(g), \\ &\qquad g \in G, k \in K\}. \end{aligned}$$

These are commutative topological algebras under convolution and their characters are precisely of the form (5.1), where ϕ is a spherical function on $G \times K$. If ϕ is a spherical function on $G \times K$ then there is a $\delta \in \hat{K}$ such that for all $x \in G$ the function $k \to \phi(xk)$ on K belongs to δ. Then δ is called a *spherical function of type* δ on G (with respect to K), cf. GODEMENT [19]. It is funny that spherical functions of type δ are on the one hand generalizations of ordinary spherical functions for (G, K), on the other hand restrictions to G of ordinary spherical functions for $(G \times K, K^*)$.

For convenience, we take a one-dimensional $\delta \in \hat{K}$. Then a spherical function ϕ on $G \times K$ is of type δ iff

$$\phi(xk) = \phi(kx) = \delta(k)\phi(x), \qquad x \in G, k \in K .$$

Let

$$\begin{aligned} & I_{c,\delta}(G) \ (\text{or } I^{\infty}_{c,\delta}(G)) \\ := & \{f \in C_c(G) \ (\text{or } C^{\infty}_c(G)) \mid f(xk) = f(kx) \\ & = \delta(k)f(x), x \in G, k \in K\} . \end{aligned}$$

These are closed subalgebras of $I_c(G)$ (or $I^{\infty}_c(G)$) and their characters are precisely of the form (5.1), where ϕ is a spherical function of type δ. Finally, if τ is a K-unitary representation of G and if $\mathscr{H}(\tau)$ contains a unit vector v satisfying $\tau(k)v = \delta(k)v$, unique up to a constant factor, then $x \to (\tau(x)v, v)$ is a spherical function of type δ.

5.3. THE GENERALIZED ABEL TRANSFORM

Let G be a connected noncompact real semisimple Lie group with finite center. Use the notation of §2.2. For given Haar measures dk, da, dn on K, A, N, respectively, normalize the Haar measure on G such that

$$\int_G f(g)dg = \int_{K \times A \times N} f(kan)e^{2\rho(\log a)}\, dk\, da\, dn, \ f \in C_c(G) \tag{5.2}$$

(cf. HELGASON [25, Ch. X, Prop. 1.11]). Note the property

$$\int_N f(n)dn = e^{2\rho(\log a)} \int_N f(ana^{-1})dn, \ f \in C_c(N), \ a \in A \tag{5.3}$$

(cf. [25, Ch. X, proof of Prop. 1.11]).