

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 28 (1982)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** THE REPRESENTATION THEORY OF  $SL(2, \mathbb{R})$ , A NON-INFINITESIMAL APPROACH  
**Autor:** Koornwinder, Tom H.  
**Kapitel:** 5.3. The generalized Abel transform  
**DOI:** <https://doi.org/10.5169/seals-52233>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 10.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

These are commutative topological algebras under convolution and their characters are precisely of the form (5.1), where  $\phi$  is a spherical function on  $G \times K$ . If  $\phi$  is a spherical function on  $G \times K$  then there is a  $\delta \in \hat{K}$  such that for all  $x \in G$  the function  $k \rightarrow \phi(xk)$  on  $K$  belongs to  $\delta$ . Then  $\delta$  is called a *spherical function of type  $\delta$*  on  $G$  (with respect to  $K$ ), cf. GODEMENT [19]. It is funny that spherical functions of type  $\delta$  are on the one hand generalizations of ordinary spherical functions for  $(G, K)$ , on the other hand restrictions to  $G$  of ordinary spherical functions for  $(G \times K, K^*)$ .

For convenience, we take a one-dimensional  $\delta \in \hat{K}$ . Then a spherical function  $\phi$  on  $G \times K$  is of type  $\delta$  iff

$$\phi(xk) = \phi(kx) = \delta(k)\phi(x), \quad x \in G, k \in K.$$

Let

$$\begin{aligned} & I_{c, \delta}(G) \text{ (or } I_{c, \delta}^\infty(G)) \\ & := \{f \in C_c(G) \text{ (or } C_c^\infty(G)) \mid f(xk) = f(kx) \\ & \quad = \delta(k)f(x), x \in G, k \in K\}. \end{aligned}$$

These are closed subalgebras of  $I_c(G)$  (or  $I_c^\infty(G)$ ) and their characters are precisely of the form (5.1), where  $\phi$  is a spherical function of type  $\delta$ . Finally, if  $\tau$  is a  $K$ -unitary representation of  $G$  and if  $\mathcal{H}(\tau)$  contains a unit vector  $v$  satisfying  $\tau(k)v = \delta(k)v$ , unique up to a constant factor, then  $x \rightarrow (\tau(x)v, v)$  is a spherical function of type  $\delta$ .

### 5.3. THE GENERALIZED ABEL TRANSFORM

Let  $G$  be a connected noncompact real semisimple Lie group with finite center. Use the notation of §2.2. For given Haar measures  $dk, da, dn$  on  $K, A, N$ , respectively, normalize the Haar measure on  $G$  such that

$$(5.2) \quad \int_G f(g)dg = \int_{K \times A \times N} f(kan)e^{2\rho(\log a)} dk da dn, \quad f \in C_c(G)$$

(cf. HELGASON [25, Ch. X, Prop. 1.11]). Note the property

$$(5.3) \quad \int_N f(n)dn = e^{2\rho(\log a)} \int_N f(ana^{-1})dn, \quad f \in C_c(N), a \in A$$

(cf. [25, Ch. X, proof of Prop. 1.11]).

For  $\lambda \in \mathfrak{a}_C^*$  let  $U^\lambda$  be the representation of  $G$  induced by the one-dimensional representation  $an \rightarrow e^{\lambda(\log a)}$  of the subgroup  $AN$ :

$$(5.4) \quad (U^\lambda(g)f)(k) := e^{-(\rho + \lambda)H(g^{-1}k)} f(u(g^{-1}k)), \quad f \in L^2(K), \quad g \in G, \quad k \in K.$$

The representation  $U^\lambda$  is easily seen to split as a direct sum of principal series representations  $\pi_{\xi, \lambda}$ .  $U^\lambda$  restricted to  $K$  is the left regular representation of  $K$ .

Let  $\delta \in \hat{K}$ . For convenience, suppose that  $\delta$  is one-dimensional. The *generalized Abel transform*  $f \rightarrow F_f^\delta : I_{c, \delta}(G) \rightarrow C_c(A)$  is defined by

$$(5.5) \quad F_f^\delta(a) := e^{\rho(\log a)} \int_N f(an) dn, \quad a \in A.$$

If  $G = SU(1, 1)$  and  $\delta = 1$  then this transform can be rewritten as the classical Abel transform, cf. §5.4.

**PROPOSITION 5.3.** *The mapping  $f \rightarrow F_f^\delta$  is a continuous homomorphism (with respect to convolution on  $G$  and  $A$ , respectively) from  $I_{c, \delta}^\infty(G)$  to  $C_c^\infty(A)$ . Furthermore,*

$$(5.6) \quad \int_A F_f^\delta(a) e^{-\lambda(\log a)} da = \int_G f(g) (U^\lambda(g^{-1})\delta, \delta) dg, \quad f \in I_{c, \delta}^\infty(G), \quad \lambda \in \mathfrak{a}_C^*,$$

where  $(., .)$  denotes the inner product on  $L^2(K)$ .

*Proof.* The continuity is immediate. The homomorphism property follows easily from (5.2) and (5.3) (cf. WARNER [49, pp. 34, 35]). For the proof of (5.6) substitute (5.4) into the right hand side of (5.6):

$$\begin{aligned} \int_G f(g) (U^\lambda(g^{-1})\delta, \delta) dg &= \int_G \int_K f(g) e^{-(\rho + \lambda)H(gk)} \delta((u(gk))^{-1}k) dk dg \\ &= \int_G f(g) e^{-(\rho + \lambda)H(g)} \delta((u(g))^{-1}) dg \\ &= \int_{K \times A \times N} f(kan) e^{(\rho - \lambda)\log a} \delta(k^{-1}) dk da dn \\ &= \int_A \int_N f(an) e^{(\rho - \lambda)\log a} dn da \\ &= \int_A F_f^\delta(a) e^{-\lambda(\log a)} da. \end{aligned}$$

□

Now let  $G = SU(1, 1)$ . Write  $F_f^n(t)$  and  $I_{c,n}^\infty(G)$  instead of  $F_f^{\delta_n}(a_t)$  and  $I_{c,\delta_n}^\infty(G)$ , respectively. If  $n \in \mathbf{Z} + \xi$  then (5.5) and (5.6) take the form

$$(5.7) \quad F_f^n(t) = e^{\frac{1}{2}t} \int_{-\infty}^{\infty} f(a_t n_z) dz$$

and

$$(5.8) \quad \int_{-\infty}^{\infty} F_f^n(t) e^{-\lambda t} dt = \int_G f(g) \pi_{\xi, \lambda, n, n}(g^{-1}) dg, \quad f \in I_{c,n}^\infty(G), \lambda \in \mathbf{C},$$

where  $dg = (2\pi)^{-1} e^t d\theta dt dz$  if  $g = u_\theta a_t n_z$ .

#### 5.4. THE MAIN THEOREM

It is the purpose of this section to prove:

**THEOREM 5.4.** *Let  $\tau$  be an irreducible  $K$ -unitary representation of  $SU(1, 1)$  which is  $K$ -finite or unitary. Then  $\tau$  is Naimark equivalent to an irreducible subrepresentation of some principal series representation  $\pi_{\xi, \lambda}$ .*

By Proposition 5.2  $\tau$  is  $K$ -multiplicity free. If  $\delta_n \in \mathcal{M}(\tau)$  then write  $\tau_{n,n}$  instead of  $\tau_{\delta_n, \delta_n}$ . In view of Theorem 4.5 and Remark 4.8 it is sufficient for the proof of Theorem 5.4 to show that for some  $\delta_n \in \mathcal{M}(\tau)$ , for some  $\lambda \in \mathbf{C}$  and for  $\xi \in \{0, \frac{1}{2}\}$  with  $n \in \mathbf{Z} + \xi$  we have

$$(5.9) \quad \tau_{n,n} = \pi_{\xi, \lambda, n, n}.$$

Both sides of (5.9) are spherical functions of type  $\delta_n$ . Then (5.9) holds if the corresponding characters on  $I_{c,n}^\infty(G)$  are equal. Hence Theorem 5.4 will follow from

**PROPOSITION 5.5.** *Let  $G = SU(1, 1)$ ,  $n \in \frac{1}{2}\mathbf{Z}$ . Let  $\alpha$  be a continuous character on  $I_{c,n}^\infty(G)$ . Then*

$$(5.10) \quad \alpha(f) = \int_G f(g) \pi_{\xi, \lambda, n, n}(g^{-1}) dg, \quad f \in I_{c,n}^\infty(G),$$

for some  $\lambda \in \mathbf{C}$  and for  $\xi \in \{0, \frac{1}{2}\}$  such that  $n \in \mathbf{Z} + \xi$ .