

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 28 (1982)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** THE REPRESENTATION THEORY OF  $SL(2, \mathbb{R})$ , A NON-INFINITESIMAL APPROACH  
**Autor:** Koornwinder, Tom H.  
**Kapitel:** 5.5. COMPLETION OF THE PROOF OF THE MAIN THEOREM  
**DOI:** <https://doi.org/10.5169/seals-52233>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 27.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

Now substitute (5.8) into the right hand side of (5.10). Thus, for the proof of Prop. 5.5 we have to show that each continuous character  $\alpha$  on  $I_{c,n}^\infty(G)$  takes the form

$$(5.11) \quad \alpha(f) = \int_{-\infty}^{\infty} F_f^n(t) e^{-\lambda t} dt, \quad f \in I_{c,n}^\infty(G).$$

for some  $\lambda \in \mathbf{C}$ . In §5.5 we will prove:

**THEOREM 5.6.** *Let  $G = SU(1, 1)$ ,  $n \in \frac{1}{2}\mathbf{Z}$ . The mapping  $f \rightarrow F_f^n$  is a topological algebra isomorphism from  $I_{c,n}^\infty(G)$  onto  $\mathcal{D}_{\text{even}}(\mathbf{R})$ , the algebra of even  $C^\infty$ -functions with compact support on  $\mathbf{R}$ .*

Thus, in view of (5.11) we are left to prove:

**PROPOSITION 5.7.** *The continuous characters on  $\mathcal{D}_{\text{even}}(\mathbf{R})$  have the form*

$$h \rightarrow \int_{-\infty}^{\infty} h(t) e^{-\lambda t} dt$$

for some  $\lambda \in \mathbf{C}$ .

## 5.5. COMPLETION OF THE PROOF OF THE MAIN THEOREM

By the discussion in §5.4 we reduced the proof of Theorem 5.4 to the task of proving Theorem 5.6 and Prop. 5.7. Theorem 5.6 was partly proved in Prop. 5.3. It is left to prove that  $f \rightarrow F_f^n$  is injective on  $I_{c,n}^\infty(G)$  with image  $\mathcal{D}_{\text{even}}(\mathbf{R})$  and that the inverse mapping is continuous. In order to establish this we identify both  $I_{c,n}^\infty(G)$  and  $\mathcal{D}_{\text{even}}(\mathbf{R})$ , considered as topological vector spaces, with  $\mathcal{D}([1, \infty))$  and we rewrite  $f \rightarrow F_f^n$  as a mapping from  $\mathcal{D}([1, \infty))$  onto itself. This mapping turns out to be a known integral transformation, for which an inverse transformation can be explicitly given. First note:

**LEMMA 5.8.** *The formula*

$$(5.12) \quad f(x) = g(x^2)$$

defines an isomorphism of topological vector spaces  $f \rightarrow g$  from  $\mathcal{D}_{\text{even}}(\mathbf{R})$  onto  $\mathcal{D}([0, \infty))$ .

*Proof.* Clearly, if  $g \in \mathcal{D}([0, \infty))$  then  $f \in \mathcal{D}_{\text{even}}(\mathbf{R})$  and the mapping  $g \rightarrow f$  is continuous. Conversely, let  $f \in \mathcal{D}_{\text{even}}(\mathbf{R})$  and let  $g$  be defined by (5.12). By complete induction with respect to  $n$  we prove:  $g^{(n)}(0)$  exists and there is a function  $f_n \in \mathcal{D}_{\text{even}}(\mathbf{R})$  such that

$$f_n(x) = g^{(n)}(x^2), x \in \mathbf{R},$$

and  $f \rightarrow f_n: \mathcal{D}_{\text{even}}(\mathbf{R}) \rightarrow \mathcal{D}_{\text{even}}(\mathbf{R})$  is continuous. Indeed, suppose this is proved up to  $n - 1$ . Then

$$2x(g^{(n-1)})'(x^2) = f'_{n-1}(x) = \int_0^x f''_{n-1}(y)dy,$$

so

$$g^{(n)}(x^2) = \frac{1}{2} \int_0^1 f''_{n-1}(tx)dt = : f_n(x). \quad \square$$

For  $f \in I_{c,n}^\infty(G)$  define

$$(5.13) \quad \tilde{f}(x) := f \left( \begin{pmatrix} x & (x^2-1)^{\frac{1}{2}} \\ (x^2-1)^{\frac{1}{2}} & x \end{pmatrix} \right), x \in [1, \infty).$$

For  $f \in I_{c,n}^\infty(G)$  define

$$(5.14) \quad \tilde{h}(ch\frac{1}{2}t) := h(t), t \in \mathbf{R}.$$

LEMMA 5.9. *The mapping  $f \rightarrow \tilde{f}$  defined by (5.13) is an isomorphism of topological vector spaces from  $I_{c,n}^\infty(G)$  onto  $\mathcal{D}([1, \infty))$ . The mapping  $h \rightarrow \tilde{h}$  defined by (5.35) is an isomorphism of topological vector spaces from  $\mathcal{D}_{\text{even}}(\mathbf{R})$  onto  $\mathcal{D}([1, \infty))$ .*

*Proof.* The second statement follows from Lemma 5.8. For the proof of the first statement introduce global real analytic coordinates on  $G$  by the mapping

$$(z, \phi) \rightarrow \begin{pmatrix} e^{\frac{1}{2}i\phi}(1+|z|^2)^{\frac{1}{2}} & z \\ \bar{z} & e^{-\frac{1}{2}i\phi}(1+|z|^2)^{\frac{1}{2}} \end{pmatrix}$$

from  $C \times (\mathbf{R}/4\pi\mathbf{Z})$  onto  $G$ . If  $g \in \mathcal{D}([1, \infty))$  and

$$f \left( \begin{pmatrix} e^{\frac{1}{2}i\phi}(1+|z|^2)^{\frac{1}{2}} & z \\ \bar{z} & e^{-\frac{1}{2}i\phi}(1+|z|^2)^{\frac{1}{2}} \end{pmatrix} \right) := e^{in\phi}g((1+|z|^2)^{\frac{1}{2}})$$

then  $f \in I_{c,n}^\infty(G)$ ,  $\tilde{f} = g$  and the mapping  $g \rightarrow f$  is continuous. Conversely, if  $f \in I_{c,n}^\infty(G)$  then  $f$ , as a function of  $z$  and  $\phi$ , is radial in  $z$ , so the function

$$z \rightarrow f \left( \begin{pmatrix} (1+z^2)^{\frac{1}{2}} & z \\ z & (1+z^2)^{\frac{1}{2}} \end{pmatrix}, z \in \mathbf{R}, \right.$$

belongs to  $\mathcal{D}_{\text{even}}(\mathbf{R})$ . Now make the transformation  $z = (x^2 - 1)^{\frac{1}{2}}$  and apply Lemma 5.8. It follows that  $\tilde{f} \in \mathcal{D}([1, \infty))$  and that the mapping  $f \rightarrow \tilde{f}$  is continuous.  $\square$

Define the Chebyshev polynomial  $T_n(x)$  by

$$(5.15) \quad T_n(\cos \theta) := \cos n\theta.$$

It follows from (5.7) that, for  $f \in I_{c,n}^\infty(G)$ :

$$\begin{aligned} F_f^n(t) &= e^{\frac{1}{2}t} \int_{-\infty}^{\infty} f \left( \begin{pmatrix} ch_{\frac{1}{2}t} + \frac{1}{2}ize^{\frac{1}{2}t} & * \\ * & * \end{pmatrix} \right) dz \\ &= e^{\frac{1}{2}t} \int_{-\infty}^{\infty} \tilde{f}(|ch_{\frac{1}{2}t} + \frac{1}{2}ize^{\frac{1}{2}t}|) \left( \frac{ch_{\frac{1}{2}t} + \frac{1}{2}ize^{\frac{1}{2}t}}{|ch_{\frac{1}{2}t} + \frac{1}{2}ize^{\frac{1}{2}t}|} \right)^{2n} dz \\ &= e^{\frac{1}{2}t} \int_0^{\infty} \tilde{f}(|ch_{\frac{1}{2}t} + \frac{1}{2}ize^{\frac{1}{2}t}|) T_{2|n|} \left( \frac{ch_{\frac{1}{2}t}}{|ch_{\frac{1}{2}t} + \frac{1}{2}ize^{\frac{1}{2}t}|} \right) dz, \end{aligned}$$

so

$$F_f^n(t) = 2 \int_{ch_{\frac{1}{2}t}}^{\infty} \tilde{f}(y) T_{2|n|}(y^{-1}ch_{\frac{1}{2}t})(y^2 - ch^2\frac{1}{2}t)^{-\frac{1}{2}} y dy.$$

This formula shows that  $F_f^n$  is even on  $\mathbf{R}$ , so  $F_f^n \in \mathcal{D}_{\text{even}}(\mathbf{R})$ . Now, by (5.14):

$$(5.16) \quad \tilde{F}_f^n(x) = 2 \int_x^{\infty} \tilde{f}(y) T_{2|n|}(y^{-1}x)(y^2 - x^2)^{-\frac{1}{2}} y dy, \quad x \in [1, \infty].$$

For  $n = 0$ , (5.16) takes the form

$$\tilde{F}_f^0(x) = 2 \int_x^{\infty} \tilde{f}(y)(y^2 - x^2)^{-\frac{1}{2}} y dy:$$

The problem of inverting this just means to solve the Abel integral equation, as was pointed out by GODEMENT [20]. Indeed, we get

$$\tilde{f}(y) = -\pi^{-1} \int_y^\infty \frac{d}{dx} \tilde{F}_f^0(x)(x^2 - y^2)^{-\frac{1}{2}} dx.$$

For general  $n$ , we can use an inversion formula obtained by DEANS [7, (30)], see also MATSUSHITA [30, §2.3] and KOORNWINDER [27, §5.9]:

THEOREM 5.10. For  $m = 0, 1, 2, \dots, g \in \mathcal{D}([1, \infty))$ ,  $x \in [1, \infty)$  define

$$(5.17) \quad (A_m g)(x) := 2 \int_x^\infty g(y) T_m(y^2 - x^2)^{-\frac{1}{2}} y dy,$$

$$(5.18) \quad (B_m g)(x) := -\pi^{-1} \int_x^\infty g'(y) T_m(x^{-1}y)(y^2 - x^2)^{-\frac{1}{2}} dy.$$

Then  $A_m$  and  $B_m$  map  $\mathcal{D}([1, \infty))$  into itself and  $A_m B_m = id, B_m A_m = id$ .

This theorem shows that  $f \rightarrow F_f^n$  is a linear bijection from  $I_{c,n}^\infty(G)$  onto  $\mathcal{D}_{\text{even}}(\mathbf{R})$ . Finally in order to prove the continuity of the inverse mapping, we show that  $B_m$  is continuous. Just expand  $T_m(x^{-1}y)$  as a polynomial and use that

$$\begin{aligned} & \left(x^{-1} \frac{d}{dy}\right)^P \int_x^\infty h(y)(y^2 - x^2)^{-\frac{1}{2}} y dy \\ &= \int_x^\infty \left(y^{-1} \frac{d}{dy}\right)^P h(y)(y^2 - x^2)^{-\frac{1}{2}} y dy \end{aligned}$$

by the properties of the Weyl fractional integral transform (cf. [11, Ch. 13]). This completes the proof of Theorem 5.6.

*Proof of proposition 5.7.* Extend  $\alpha$  to a continuous linear functional on  $\mathcal{D}(\mathbf{R})$ , for instance by putting  $\alpha(f) = 0$  if  $f$  is odd. Choose  $f_1 \in \mathcal{D}_{\text{even}}(\mathbf{R})$  such that  $\alpha(f_1) \neq 0$ . Let

$$(\lambda(y)f_1)(x) := f_1(x - y), \quad x, y \in \mathbf{R}.$$

By the continuity and homomorphism property of  $\alpha$  we have, for  $f \in \mathcal{D}_{\text{even}}(\mathbf{R})$ :

$$\alpha(f_1)\alpha(f) = \alpha(f_1 * f) = \int_{-\infty}^{\infty} \alpha(\lambda(y)f_1)f(y)dy.$$

Hence

$$\alpha(f) = \int_{-\infty}^{\infty} f(y)\beta(y)dy, f \in \mathcal{D}_{\text{even}}(\mathbf{R}),$$

where

$$\beta(y) := \frac{1}{2}(\alpha(f_1))^{-1}(\alpha(\lambda(y)f_1) + \alpha(\lambda(-y)f_1)).$$

Then  $\beta$  is even and it is a continuous function by the continuity of  $\alpha$ . It follows from the homomorphism property of  $\alpha$  and from the fact that  $\beta$  is even, that

$$\beta(x)\beta(y) = \frac{1}{2}(\beta(x+y) + \beta(x-y)),$$

so  $\beta(0) = 1$ . This is d'Alembert's functional equation. By continuity,  $\text{Re } \beta(x) > 0$  if  $0 \leq x \leq x_0$  for some  $x_0 > 0$ . Then  $\beta(x_0) = \cosh c$  for some complex  $c = a + ib$  with  $a \geq 0$ ,  $-\frac{1}{2}\pi < b < \frac{1}{2}\pi$ . Now, following the proof in ACZEL [1, 2.4.1] it can be shown <sup>1)</sup> that for all integer  $n, m \geq 0$

$$\beta\left(\frac{n}{2^m}x_0\right) = \cosh\left(\frac{c}{x_0}\frac{n}{2^m}x_0\right).$$

So, by continuity and evenness of  $\beta$ :

$$\beta(x) = \cosh\left(\frac{c}{x_0}x\right) \text{ for all } x \in \mathbf{R}. \quad \square$$

## 5.6. NOTES

5.6.1. Some other examples of Gelfand pairs  $(G \times K, K^*)$  are provided by  $G = SO_0(n, 1)$ ,  $K = SO(n)$  and  $G = SU(n, 1)$ ,  $K = S(U(n) \times U(1))$ , cf. BOERNER [4, Ch. VII, §12; Ch. V, §6], DIXMIER [8] or KOORNWINDER [27, Theorems 5.7, 5.8].

5.6.2. The main Theorem 5.4, which was first proved in the case of unitary representations by BARGMANN [2], is a special case of the *subrepresentation*

<sup>1)</sup> I thank H. van Haeringen for this reference.