

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 28 (1982)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: NON-STANDARD MODELS OF PEANO ARITHMETIC
Autor: Kochen, Simon / Kripke, Saul
Kapitel: IV. Ramsey-type Theorems
DOI: <https://doi.org/10.5169/seals-52238>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 10.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

then we can construct the sequence $\{h_i\}$ and the set \mathcal{F} to satisfy the Stability and Closure Conditions via Theorem 2.

We could now proceed to show that the above condition is indeed satisfied in \mathbf{N} and thus construct a non-standard model of Peano arithmetic. However, our goal is the construction of a mathematically perspicuous model which is independent of the logical formulas. The functions $\{h_i\}$ given by the above condition require the logical calculus in their definition. Accordingly, we shall consider a larger class \mathcal{F} of functions than those defined above, which we shall construct independently of logical formulas. This class will be constructed from combinatorial principles derived from Ramsey's Partition Theorem.

IV. RAMSEY-TYPE THEOREMS

The infinite Ramsey Theorem states that for every partition $P : [\mathbf{N}]^e \rightarrow r^1$) there exists an infinite subset X of \mathbf{N} such that $P|_{[X]^e}$ is constant. In these circumstances one says that X is homogeneous for the partition P . This set-theoretic theorem has various combinatorial consequences which are formalizable in elementary arithmetic. One such immediate consequence which we shall prove independent of the Peano axioms is the following.

PROPOSITION 1. *Let $P : [\mathbf{N}]^e \rightarrow r$ be a primitive recursive partition. For every natural number k there exists a finite subset X of \mathbf{N} , with $\# X \geq k$ and $\# X \geq 2^{2^{\min X}}$, which is homogeneous for the partition P .*

In order to apply Theorem 2 we require the construction of a set which is simultaneously homogeneous for several partitions. This is easily done by the infinite Ramsey Theorem. Suppose $P_1 : [\mathbf{N}]^{e_1} \rightarrow r_1$ and $P_2 : [\mathbf{N}]^{e_2} \rightarrow r_2$ are two partitions. Let X_1 be an infinite subset of \mathbf{N} homogeneous for P_1 . Then $P_2|_{[X_1]^{e_2}}$ is a partition of $[X_1]^{e_2}$, and hence there is an infinite subset X_2 of X_1 which is homogeneous for P_2 (as well as P_1). This proof extends immediately to finitely many partitions. A direct consequence is the following generalization of Proposition 1.

PROPOSITION 2. *Let $P_i : [\mathbf{N}]^{e_i} \rightarrow r_i$, $i \leq i \leq n$ be a set of primitive recursive partitions. For every natural number k there exists a finite subset X of \mathbf{N} with $\# X \geq k$ and $\# X \geq 2^{2^{\min X}}$, which is simultaneously homogeneous for all the partitions P_1, \dots, P_n .*

¹⁾ We identify the number r with the set of all natural numbers $< r$.

Proposition 2 may be expressed by a \prod_2^0 formula. First it is clear that we can construct a \sum_1^0 -formula ϕ_i that expresses the properties that

1. $P_i : [\mathbb{N}]^{e_i} \rightarrow r_i$ is a primitive recursive partition
2. $z_1 < z_2 < \dots < z_{n_k}$
3. $\{z_1, \dots, z_{n_k}\}$ is homogeneous for P_i
4. $k \leq n_k$
5. $2^{2^{z_1}} \leq n_k$

Proposition 2 asserts that for every k

$$\mathbb{N} \models \exists z_1 \dots \exists z_{n_k} \bigwedge_{i \leq k} \phi_i .$$

V. CONSTRUCTION OF THE MODEL

We now have all the ingredients at hand to construct a non-standard model of Peano arithmetic, and we have only to assemble them according to the specifications of Section II.

Let P_i be an effective enumeration of all primitive recursive partitions $P_i : [\mathbb{N}]^{e_i} \rightarrow r_i$. By Proposition 2 we have that for every k

$$\mathbb{N} \models \exists z_1 \dots \exists z_{n_k} \bigwedge_{i \leq k} \phi_i$$

where ϕ_i is the \sum_1^0 -formula of Section IV expressing the conditions (1)-(5) satisfied by the partition P_i .

Following the prescription given in Section III we let a_{kn_k} be the smallest number such that a_{k_1}, \dots, a_{kn_k} is an increasing sequence satisfying the formula $\bigwedge_{j \leq k} \phi_j$. Now we define the functions h_j by

$$h_0(k) = n_k \quad \text{for every } k$$

and for $j > 0$

$$h_j(k) = \begin{cases} a_{kj} & \text{for } j \leq n_k \\ h_{j-1}(k)^2 & \text{for } j > n_k . \end{cases}$$

Let $\mathcal{F} = \{f \mid f \leq h_j\}$.

Since $\mathbf{1} \leq h_0$ the function $\mathbf{1}$ is automatically in \mathcal{F} .

By Theorem 2 the sequence $\{h_j\}$ satisfies $\bigwedge_{j < \infty} \phi_j$ in \mathcal{F}/D . We now prove that this implies that the sequence $\{h_j\}$ satisfies the Stability and