

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 28 (1982)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** NON-STANDARD MODELS OF PEANO ARITHMETIC  
**Autor:** Kochen, Simon / Kripke, Saul  
**Kapitel:** Note (Added in proof)  
**DOI:** <https://doi.org/10.5169/seals-52238>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 27.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

We now define the functions  $h_j$  as before by

$$h_j(k) = \begin{cases} a_{kj} & \text{for } j \leq k \\ h_{j-1}^2(k) & \text{for } j > k. \end{cases}$$

A simple induction argument now shows that (1) holds for the axioms and hence that  $\mathcal{F}/D$  is a model of Peano arithmetic. If as in Theorem 3 the  $a_{kk}$  is chosen as the least number so that the sequence  $a_{k1}, \dots, a_{kk}$  satisfies the above conditions then a true statement which is false in  $\mathcal{F}/D$  may be constructed via the method of Theorem 3.

The disadvantage of this direct approach is that the model  $\mathcal{F}/D$  constructed in this manner is dependent in its definition upon logical formulas and so is not as purely an algebraic construction. Moreover the independent statement which results has no simple combinatorial expression as have those given in Sections IV, V, and VI. Note that in this approach we have not used the property peculiar to Peano's axioms concerning the limited associates of the axioms which is expressed in the proof of Theorem 4. This shows that the method outlined here applies to any recursively enumerable set of axioms for arithmetic which is sufficient to allow the coding required for Theorem 3. Thus, we may prove a general form of Gödel's Incompleteness Theorem without the use of self-reference techniques. At the same time the very generality of the approach outlined here indicates that there is no hope by this method to avoid the use of metamathematics. It is only the above-mentioned property of the Peano axioms vis-à-vis limited formulas that allowed us the latitude to define suitable functions  $h_j$ , and hence the model  $\mathcal{F}/D$ , by means of a combinatorial principal without reference to logical formulas.

#### NOTE (ADDED IN PROOF)

The first sentence of the section entitled "Added in proof" of Kochen and Kripke [12] p. 294, which was inserted by the second author, is not correct and should be deleted in favor of the following corrected version. The first author proposed that the Paris-Harrington statement is false in an initial segment of any non-standard model, and this was verified jointly by the two authors. Adapting this idea, the second author defined the set  $\mathcal{F}$  of functions which result in the model of Section V. The first author subsequently found the new set  $\mathcal{F}$  of functions which define the simpler model of Section VI.

The devices used in Section III are an adaptation of the ideas in Paris-Harrington [3].

The approach outlined in (d) of Section VII is due to the second author and leads to a concept of 'satisfying' formulas by finite sequences called *fulfillability* which leads to model-theoretic proofs of many theorems (such as Gödel's and Rosser's theorems) usually proved proof-theoretically and to other applications to the model theory and proof theory of arithmetic. It will be developed in a subsequent paper of the second author.

#### BIBLIOGRAPHY

- [1] McDOWELL, R. und E. P. SPECKER. Modelle der Arithmetik. *Infinistic Methods. Proc. Symp. on Foundations of Math.* Warsaw (1959) Perg. Press, 257-263.
- [2] SPECKER, E. P. Ramsey's theorem does not hold in recursive set theory. *Logic Colloquium* (1969), 439-442.
- [3] PARIS, J. and L. HARRINGTON. A mathematical incompleteness in Peano arithmetic. *Handbook of Mathematical Logic*, North-Holland (1977), 1133-1142.
- [4] KIRBY, L. *Initial Segments of Models of Arithmetic*. Ph.D. Thesis (1977), Manchester University.
- [5] PARIS, J. Some independence results for Peano arithmetic. *JSL* (1979).
- [6] SCOTT, D. On constructing models of arithmetic. *Infinistic Methods, Proc. Symp. in Foundations of Math.*, Warsaw (1959), Perg. Press, 235-255.
- [7] SKOLEM, Th. Über die Nicht-Characterisierbarkeit der Zahlenreihe mittels endlich oder abzählbar unendlich vieler Aussagen mit ausschliesslich Zahlenvariablen. *Fund. Math.* 23 (1934).
- [8] SHOENFIELD, J. *Mathematical Logic*. Addison-Wesley, 1967.
- [9] KLEENE, S. *Introduction to Metamathematics*. Van Nostrand, 1962.
- [10] JOCKUSCH, C., Jr. Ramsey's Theorem and recursion theory. *JSL* (1972), 269-280.
- [11] RAMSEY, F. P. On a problem in formal logic. *Proc. London Math. Soc.* 30 (1930), 264-286.
- [12] KOCHEN, S. and S. KRIPKE. Non-standard models of Peano arithmetic. *Logic and algorithmic*, Monographie de l'Enseignement Mathématique N° 30, Université de Genève, 1982, 275-295.

(Reçu le 9 avril 1981)

Simon Kochen  
Saul Kripke

Princeton University  
Department of Mathematics  
Princeton, N.J. 08540  
USA