

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 28 (1982)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ON BOOLEAN ALGEBRAS WITH DISTINGUISHED SUBALGEBRAS  
**Autor:** Koppelberg, Sabine  
**Kapitel:** 1. The sheaf representation of Boolean algebra extensions  
**DOI:** <https://doi.org/10.5169/seals-52239>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 27.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

1. THE SHEAF REPRESENTATION OF BOOLEAN ALGEBRA EXTENSIONS

Let  $\mathcal{L}$  be any language for first-order predicate logic. Suppose  $X$  is a non-empty set and for every  $p \in X$  we have an  $\mathcal{L}$ -structure  $\mathcal{B}_p = (B_p, \dots)$ ; put  $S = \bigcup_{p \in X} B_p$ . Suppose  $\varphi(x_1 \dots x_n)$  is an  $\mathcal{L}$ -formula,  $u \subseteq X$  and  $f_1, \dots, f_n : u \rightarrow S$  are such that  $f_i(p) \in B_p$  for  $1 \leq i \leq n$  and  $p \in u$ . Then let

$$\| \varphi [f_1 \dots f_n] \| = \{ p \in u \mid \mathcal{B}_p \models \varphi [f_1(p) \dots f_n(p)] \} .$$

We may think of  $\| \varphi [f_1 \dots f_n] \| \subseteq X$  as being a (Boolean) truth value of  $\varphi [f_1 \dots f_n]$  in the power set of  $X$ .

A sheaf of  $\mathcal{L}$ -structures is a sequence

$$\mathcal{S} = (S, \pi, X, \mu)$$

such that a)  $S$  and  $X$  are topological spaces and  $\pi : S \rightarrow X$  is a continuous open local homeomorphism from  $S$  onto  $X$ , b)  $\mu$  is a function assigning to each  $p \in X$  an  $\mathcal{L}$ -structure  $\mathcal{B}_p = (B_p, \dots)$  such that the  $B_p$  are pairwise disjoint,  $S = \bigcup_{p \in X} B_p$  and  $\pi(s) = p$  iff  $s \in B_p$ , c) for every open subset  $u$  of  $X$  and continuous  $f_1, \dots, f_n : u \rightarrow S$  satisfying  $f_i(p) \in B_p$  for  $p \in u$  and every atomic  $\mathcal{L}$ -formula  $\varphi(x_1 \dots x_n)$ ,  $\| \varphi [f_1 \dots f_n] \|$  is an open subset of  $u$ .

The  $\mathcal{L}$ -structure  $\mathcal{B}_p$  is called the stalk of  $\mathcal{S}$  over  $p$ . — Let, if  $\mathcal{S}$  is a sheaf of  $\mathcal{L}$ -structures,  $\Gamma(\mathcal{S})$  be the set of all continuous functions  $f : X \rightarrow S$  satisfying  $f(p) \in B_p$  for  $p \in X$  (the set of “global sections” of  $\mathcal{S}$ ). So  $\Gamma(\mathcal{S})$  is, if non-empty, (the underlying set of) a substructure of the product structure  $\prod_{p \in X} \mathcal{B}_p$ , hence an  $\mathcal{L}$ -structure.

For the rest of the paper, let  $\mathcal{L} = \{ +, \cdot, -, 0, 1, U \}$  where  $U$  is a unary predicate. We indicate how, for a given  $BA$  extension  $(B, A)$ ,  $B$  may be represented by  $\Gamma(\mathcal{S})$  where  $\mathcal{S}$  is a sheaf of  $\mathcal{L}$ -structures over a Boolean space. We omit most of the proofs which are easy and entirely analogous to well-known representation theorems for lattices over Boolean spaces. Let  $X$  be the Stone space of  $A$ , i.e. the set of all ultrafilters of  $A$  with the usual topology. For  $p \in X$ , let  $\langle p \rangle$  be the filter of  $B$  generated by  $p$ . Let  $\pi_p : B \rightarrow B / \langle p \rangle = B_p$  be the canonical epimorphism. So  $B_p$  is a  $BA$  with at least two elements. For  $p, q \in X$  and  $p \neq q$ ,  $B_p$  and  $B_q$  are disjoint. Let  $S = \bigcup_{p \in X} B_p$  and  $\pi : S \rightarrow X$  be defined as stated in b) above. Let, for  $p \in X$ ,  $\mu(p)$  be the  $\mathcal{L}$ -structure  $(B_p, \dots, \{0, 1\})$ . For  $u \subseteq X$  open and  $b \in B$ , let  $M_{ub} = \{ \pi_p(b) \mid p \in u \}$ . The set of these  $M_{ub}$  constitutes a base

for a topology of  $S$ , and this makes  $\mathcal{S} = (S, \pi, X, \mu)$  a sheaf of  $\mathcal{L}$ -structures. Furthermore, for  $b \in B$ ,  $\sigma_b : X \rightarrow S$  defined by  $\sigma_b(p) = \pi_p(b)$  is a global section of  $\mathcal{S}$  and

$$\left. \begin{array}{l} \sigma : B \rightarrow \Gamma(\mathcal{S}) \\ b \mapsto \sigma_b \end{array} \right\}$$

is an isomorphism from  $B$  onto  $\Gamma(\mathcal{S})$ . We shall now identify  $B$  with  $\Gamma(\mathcal{S})$ ; so every  $b \in B$  is a function from  $X$  to  $S$ . This identifies  $A$  with those  $b \in B$  such that for every  $p \in X$   $b(p) = 0$  or  $b(p) = 1$ , i.e. with those  $b \in B$  satisfying  $\|U(b)\| = X$ . Let  $C$  be the  $BA$  of clopen subsets of  $X$  and  $e(c)$  the characteristic function of  $c$  for  $c \in C$ . Thus  $e$  is an isomorphism from  $C$  onto  $A \subseteq B$ .

In the rest of this section, we show that the property of being a Hausdorff sheaf for  $\mathcal{S}$  is equivalent to a property of the extension  $(B, A)$  which reflects, in a way which is first-order expressible in  $\mathcal{L}$ , completeness of the embedding of  $A$  into  $B$ . Recall that, for a sheaf  $\mathcal{S}$  over a Boolean space  $X$ ,  $S$  is a  $T_2$ -space iff, for any  $f, g \in \Gamma(\mathcal{S})$ ,  $\|f = g\|$  is a clopen subset of  $X$ ;  $\mathcal{S}$  is then said to be a Hausdorff sheaf. Call  $A$  relatively complete in  $B$  if, for every  $b \in B$ , there is a largest element  $a \in A$  such that  $a \leq b$ , equivalently: for  $b \in B$ , there is a largest  $a \in A$  such that  $a \cdot b = 0$  or: for  $b \in B$ , there is a smallest  $a \in A$  such that  $b \leq a$ .

1.1. PROPOSITION.  $\mathcal{S}$  is a Hausdorff sheaf iff  $A$  is relatively complete in  $B$ .

*Proof.* Suppose  $\mathcal{S}$  is Hausdorff and  $b \in B$ . Let  $d \in B$  such that  $d(p) = 0$  for every  $p \in X$ , let  $c = \|b = d\|$  and  $a = e(c)$ . Then  $a$  is the largest element of  $A$  satisfying  $a \cdot b = 0$ .

Conversely, let  $A$  be relatively complete in  $B$  and suppose  $f, g \in B$ . Let  $a$  be the largest element of  $A$  such that  $a \leq f \cdot g + -f \cdot -g$ . Let  $c \in C$  such that  $a = e(c)$ . Then  $\|f = g\| = c$  is a clopen subset of  $X$ .

1.2. REMARK. Let  $A$  be relatively complete in  $B$ . Then the inclusion map from  $A$  to  $B$  is a complete homomorphism.

*Proof.* Suppose  $M$  is a subset of  $A$  having a supremum  $a$  in  $A$ . We show that  $a$  is the supremum of  $M$  in  $B$ . Clearly,  $a$  is an upper bound for  $M$  in  $B$ . Suppose that  $b$  is another upper bound for  $M$  in  $B$ . Let  $\alpha \in A$  be the largest element of  $A$  such that  $\alpha \leq b$ . For every  $m \in M \subseteq A$ , we have  $m \leq b$ , hence  $m \leq \alpha$  and  $a \leq \alpha \leq b$ .

The following facts are trivial:

1.3. REMARK. a) Let  $A$  and the inclusion map from  $A$  to  $B$  be complete. Then  $A$  is relatively complete in  $B$ .

b) Suppose  $A$  is relatively complete in  $B$  and  $B$  is complete. Then  $A$  is complete.

## 2. RELATIVE AUTOMORPHISMS OF FINITE EXTENSIONS

We first give an internal description of a finite extension  $(B, A)$  where  $B = A(u_1 \dots u_n)$  and  $n \in \omega$ . We shall always assume that  $u_1, \dots, u_n$  are the atoms of the subalgebra of  $B$  generated by  $u_1, \dots, u_n$ ; i.e. that they are non-zero, pairwise disjoint and  $u_1 + \dots + u_n = 1$ . Let  $I_r = \{a \in A \mid a \cdot u_r = 0\}$  for  $1 \leq r \leq n$ . Clearly, each  $I_r$  is a proper ideal of  $A$  and  $I_1 \cap \dots \cap I_n = \{0\}$ . The family  $(I_r \mid 1 \leq r \leq n)$  completely characterizes the extension  $(B, A)$ :

2.1. REMARK. Suppose  $C = A(v_1 \dots v_n)$  is a finite extension of  $A$  where  $v_1, \dots, v_n$  are pairwise disjoint and  $1 = v_1 + \dots + v_n$ . Let  $B = A(u_1 \dots u_n)$  be as above. There is an isomorphism  $g$  from  $B$  onto  $C$  satisfying  $g(a) = a$  for  $a \in A$  and  $g(u_r) = v_r$  iff, for each  $r$ ,  $\{a \in A \mid a \cdot v_r = 0\} = I_r$ .

*Proof.* By Theorem 12.4 in [7].

2.2. REMARK.  $A$  is relatively complete in  $B = A(u_1 \dots u_n)$  iff, for each  $r$ ,  $I_r$  is a principal ideal.

*Proof.* The only-if part follows by the definition of relative completeness. Now suppose  $\alpha_r \in A$  generates  $I_r$ ; let  $b \in B$  and  $I = \{a \in A \mid a \cdot b = 0\}$ . There are  $a_1, \dots, a_n \in A$  such that  $b = a_1 \cdot u_1 + \dots + a_n \cdot u_n$ . It follows that  $I$  is the principal ideal generated by  $\alpha = (-a_1 + \alpha_1) \cdot \dots \cdot (-a_n + \alpha_n)$ .

Conversely, given any family  $(I_r \mid 1 \leq r \leq n)$  of proper ideals in  $A$  satisfying  $I_1 \cap \dots \cap I_n = \{0\}$ , there is an extension  $A(u_1 \dots u_n)$  of  $A$  such that  $I_r = \{a \in A \mid a \cdot u_r = 0\}$ : let  $D = A(x_1 \dots x_n)$  be the free product of  $A$  and a finite BA with atoms  $x_1, \dots, x_n$ . Let

$$K = \{i_1 \cdot x_1 + \dots + i_n \cdot x_n \mid i_1 \in I_1, \dots, i_n \in I_n\}.$$

$K$  is an ideal of  $D$ ; the canonical epimorphism  $\pi$  from  $D$  onto  $B = D/K$  is one-one on  $A$ , and for  $a \in A$ ,  $\pi(a) \cdot u_r = 0$  iff  $a \in I_r$  where  $u_r = \pi(x_r)$ . Now identify  $A$  with the subalgebra  $\pi(A)$  of  $B$ .

For the rest of this section we think, as in section 1, of  $B$  as being the set of global sections of a sheaf  $\mathcal{S} = (S, \pi, X, \mu)$  of Boolean algebras over a