

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 29 (1983)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON POLYLOGARITHMS, HURWITZ ZETA FUNCTIONS, AND THE KUBERT IDENTITIES
Autor: Milnor, John
Anhang: Appendix 1 Relations between polylogarithm and Hurwitz function
DOI: <https://doi.org/10.5169/seals-52983>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

is non-zero according to Dirichlet. Thus we obtain a contribution of $-1 + \varphi(m)/2$ to the rank coming from the non-trivial even characters.

On the other hand, for the eigenspace corresponding to the trivial character, using formula (10) of §4 we obtain a contribution equal to the number of primes dividing m . Lemmas 8 and 10 of §5 now complete the proof. \square

APPENDIX 1

RELATIONS BETWEEN POLYLOGARITHM AND HURWITZ FUNCTION

For every complex number s , it follows from Theorem 1 that there exists a linear relation between the even [or the odd] part of the function $l_s(x)$ and of the function $\zeta_{1-s}(x)$ or $\beta_s(x) = -s\zeta_{1-s}(x)$. This appendix will work out the precise form of these relations. Compare [3], [19], [27].

For integer values of s , the required relation can be obtained as follows. Recall from formula (9) of §2 that

$$l_0(x) = (-1 + i \cot \pi x)/2$$

hence

$$l_0(x) + l_0(1-x) + \beta_0(x) = 0.$$

Integrating, we see that

$$l_1(x) - l_1(1-x) + 2\pi i \beta_1(x)/1! = 0$$

$$l_2(x) + l_2(1-x) + (2\pi i)^2 \beta_2(x)/2! = 0$$

and so on, for $0 < x < 1$. For even values of the subscript, specializing to $x = 0$ as in §4, this yields Euler's formula

$$2\zeta(2k) + (2\pi i)^{2k} b_{2k}/(2k)! = 0.$$

In particular, it follows that $\zeta(0) = -\frac{1}{2}$, and that the numbers $b_2, -b_4, b_6, -b_8, \dots$ are strictly positive. On the other hand, differentiating the formula for $l_0(x)$, we obtain

$$l_{-1}(x) = -\operatorname{csc}^2(\pi x)/4.$$

This is an even function satisfying $(*_{-1})$, so it must be some multiple of $\zeta_2(x) + \zeta_2(1-x)$. Comparing asymptotic behavior as $x \rightarrow 0$, we obtain the classical formula

$$\zeta_2(x) + \zeta_2(1-x) = \pi^2/\sin^2 \pi x = (2\pi i)^2 l_{-1}(x)/1!.$$

Differentiating, we see that

$$\begin{aligned}
 -\zeta_3(x) + \zeta_3(1-x) &= (2\pi i)^3 l_{-2}(x)/2! \\
 \zeta_4(x) + \zeta_4(1-x) &= (2\pi i)^4 l_{-3}(x)/3!
 \end{aligned}$$

and so on.

For $s \neq 0, 1, 2, \dots$ we know from §3 that there is some relation of the form

$$(14) \quad l_s(x) = A_s \zeta_{1-s}(x) + B_s \zeta_{1-s}(1-x)$$

for $0 < x < 1$; where A_s and B_s are certain uniquely determined constants. Expressing each of these functions of x as the sum of an even part and an odd part, we see that

$$(15) \quad \begin{cases} l_s^{\text{even}}(x) = (A_s + B_s) \zeta_{1-s}^{\text{even}}(x) \\ l_s^{\text{odd}}(x) = (A_s - B_s) \zeta_{1-s}^{\text{odd}}(x) . \end{cases}$$

Evidently the functions $s \mapsto A_s \pm B_s$ are meromorphic, taking finite non-zero values for all $s \in \mathbf{C} - \mathbf{Z}$. Differentiating with respect to x , we see that

$$(16) \quad A_s \pm B_s = s(A_{s+1} \mp B_{s+1})/(2\pi i) .$$

For integral values of s , using the discussion above, we easily obtain the following table of values, where $0! = 1$.

s	...	-2	-1	0	1	2	3	...
$1, + B_s$...	0	$\frac{2 \cdot 1!}{(2\pi i)^2}$	0	∞	$\frac{(2\pi i)^2}{2 \cdot 1!}$	∞	...
$1, - B_s$...	$-\frac{2 \cdot 2!}{(2\pi i)^3}$	0	$-\frac{2 \cdot 0!}{2\pi i}$	$\frac{2\pi i}{2 \cdot 0!}$	∞	$\frac{(2\pi i)^3}{2 \cdot 2!}$...

Now suppose that we specialize to $x = 0$, by the procedure of §4. Then equation (14) reduces to a form

$$\zeta(s) = (A_s + B_s) \zeta(1-s)$$

of Riemann's functional equation. It follows that

$$(A_s + B_s) (A_{1-s} + B_{1-s}) = 1 ,$$

and hence using (16) that

$$(A_s - B_s) (A_{1-s} - B_{1-s}) = -1 .$$

This discussion gives all of the information about $A_s \pm B_s$ which we will need. However, it is possible to compute precise values as follows. Let $\zeta_{1-s}(e^{2\pi i}x)$ be the result of analytic continuation in a loop circling the origin. Then evidently

$$\zeta_{1-s}(e^{2\pi i}x) - \zeta_{1-s}(x) = (e^{2\pi is} - 1)x^{s-1}.$$

Using the integral formula (6), computation shows that

$$l_s(e^{2\pi i}x) - l_s(x) = -(2\pi i)^s x^{s-1} / \Gamma(s).$$

Comparing these two expressions, and noting that $\zeta_{1-s}(1-x)$ is holomorphic throughout a neighborhood of $x = 0$, we can solve for A_s . The result after some manipulation is

$$A_s = \frac{i(2\pi)^s e^{-\pi is/2}}{2\Gamma(s) \sin(\pi s)}.$$

Now comparing the behavior of l_s and ζ_{1-s} under complex conjugation we see easily that

$$B_s = \overline{A_s} = \frac{-i(2\pi)^s e^{\pi is/2}}{2\Gamma(s) \sin(\pi s)}.$$

In particular, it follows that

$$A_s + B_s = \frac{(2\pi)^s}{2\Gamma(s) \cos(\pi s/2)}, \quad A_s - B_s = \frac{i(2\pi)^s}{2\Gamma(s) \sin(\pi s/2)}.$$

As an application of formula (15), let us prove the corresponding functional equation for a Dirichlet L -function. Recall from Lemma 14 that for any primitive Dirichlet character χ modulo m the function

$$L(s, \chi) = \sum_1^m \chi(k) \zeta_s(k/m) / m^s$$

satisfies

$$L(s, \bar{\chi}) = \sum_1^m \chi(k) l_s(k/m) / \tau.$$

Here we may just as well use either the even or the odd parts of ζ_s and l_s according as $\chi(-1)$ is $+1$ or -1 . Therefore, it follows from (15) that

$$\begin{aligned} L(s, \bar{\chi}) &= (A_s \pm B_s) \sum_1^m \chi(k) \zeta_{1-s}(k/m) / \tau \\ &= m^s (A_s \pm B_s) L(1-s, \chi) / \tau. \end{aligned}$$

Thus we have proved the functional equation

$$(17) \quad L(s, \bar{\chi}) = m^{1-s} (A_s + \chi(-1)B_s) L(1-s, \chi) / \tau(\chi).$$

Here the factor m^{1-s}/τ is never zero or infinite, while $A_s \pm B_s$ is zero or infinite only at certain integer values, as indicated in the table above.

The proof of Lemma 13 can now easily be completed as follows. If $s \leq 0$ is an integer, then $L(1-s, \chi) \neq 0$, so it follows that $L(s, \bar{\chi})$ equals zero if and only if $A_s \pm B_s$ is zero, as indicated in the table. \square

APPENDIX 2

SOME RELATIVES OF THE GAMMA FUNCTION

This appendix will describe certain functions $\gamma_1(x), \gamma_2(x), \dots$ which satisfy a modified form of the Kubert identities, with a polynomial correction term. (See (22) below.) They are defined as partial derivatives of the Hurwitz function by the formula

$$(18) \quad \gamma_{1-t}(x) = \partial \zeta_t(x) / \partial t.$$

We will show that γ_1 is related to the classical gamma function via Lerch's identity

$$(19) \quad \gamma_1(x) = \log(\Gamma(x)/\sqrt{2\pi}).$$

(Compare [27, p. 60].) As a bonus, we will give a self-contained exposition of the basic properties of the gamma function, based on formulas (18) and (19).

Let us begin with equation (18), which defines $\gamma_s(x)$ as an analytic function of both variables for all $s \neq 0$ and all $x > 0$. Recall that the Hurwitz function $\zeta_t(x) = x^{-t} + (x+1)^{-t} + \dots$ (analytically extended in t for $t \neq 1$) satisfies

$$\zeta_t(x+1) = \zeta_t(x) - x^{-t}.$$

Differentiating with respect to t , and then substituting $t = 1 - s$, we obtain

$$(20) \quad \gamma_s(x+1) = \gamma_s(x) + x^{s-1} \log x.$$

In particular,

$$\gamma_1(x+1) = \gamma_1(x) + \log x.$$

Note that

$$\zeta'_t(x) = -t\zeta_{t+1}(x)$$

hence

$$\zeta''_t(x) = t(t+1)\zeta_{t+2}(x),$$