

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 29 (1983)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON POLYLOGARITHMS, HURWITZ ZETA FUNCTIONS, AND THE KUBERT IDENTITIES
Autor: Milnor, John
Anhang: Appendix 2 SOME RELATIVES OF THE GAMMA FUNCTION
DOI: <https://doi.org/10.5169/seals-52983>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 26.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Here the factor m^{1-s}/τ is never zero or infinite, while $A_s \pm B_s$ is zero or infinite only at certain integer values, as indicated in the table above.

The proof of Lemma 13 can now easily be completed as follows. If $s \leq 0$ is an integer, then $L(1-s, \chi) \neq 0$, so it follows that $L(s, \bar{\chi})$ equals zero if and only if $A_s \pm B_s$ is zero, as indicated in the table. \square

APPENDIX 2

SOME RELATIVES OF THE GAMMA FUNCTION

This appendix will describe certain functions $\gamma_1(x)$, $\gamma_2(x)$, ... which satisfy a modified form of the Kubert identities, with a polynomial correction term. (See (22) below.) They are defined as partial derivatives of the Hurwitz function by the formula

$$(18) \quad \gamma_{1-t}(x) = \partial \zeta_t(x) / \partial t.$$

We will show that γ_1 is related to the classical gamma function via Lerch's identity

$$(19) \quad \gamma_1(x) = \log(\Gamma(x)/\sqrt{2\pi}).$$

(Compare [27, p. 60].) As a bonus, we will give a self-contained exposition of the basic properties of the gamma function, based on formulas (18) and (19).

Let us begin with equation (18), which defines $\gamma_s(x)$ as an analytic function of both variables for all $s \neq 0$ and all $x > 0$. Recall that the Hurwitz function $\zeta_t(x) = x^{-t} + (x+1)^{-t} + \dots$ (analytically extended in t for $t \neq 1$) satisfies

$$\zeta_t(x+1) = \zeta_t(x) - x^{-t}.$$

Differentiating with respect to t , and then substituting $t = 1 - s$, we obtain

$$(20) \quad \gamma_s(x+1) = \gamma_s(x) + x^{s-1} \log x.$$

In particular,

$$\gamma_1(x+1) = \gamma_1(x) + \log x.$$

Note that

$$\zeta'_t(x) = -t\zeta_{t+1}(x)$$

hence

$$\zeta''_t(x) = t(t+1)\zeta_{t+2}(x),$$

where the prime stands for the derivative with respect to x . By analytic continuation, this last equation holds also at $t = 0$. Differentiating with respect to t at $t = 0$, we obtain

$$(21) \quad \gamma_1''(x) = \zeta_2(x).$$

In particular, it follows that $\gamma_1''(x) > 0$ for all $x > 0$.

Let us define the gamma function as follows. (Compare Artin [1].)

LEMMA 15 (Bohr and Mollerup). *There is one and only one twice continuously differentiable function $\Gamma(x) > 0$ for $x > 0$ which satisfies*

$$\Gamma(x+1) = x\Gamma(x), \quad \Gamma(1) = 1, \quad \text{and} \quad (\log \Gamma(x))'' \geq 0.$$

Proof. Evidently it suffices to show that there is one and, up to an additive constant, only one C^2 -function

$$f(x) = \log \Gamma(x) + c$$

for $x > 0$ which satisfies the two conditions

$$f(x+1) = f(x) + \log x$$

and

$$f''(x) \geq 0.$$

Existence is clear, since the equation $\gamma_1(x)$ satisfies both of these conditions. To prove uniqueness, let us differentiate twice to obtain

$$f''(x+1) = f''(x) - 1/x^2,$$

hence

$$f''(x+n+1) = f''(x) - x^{-2} - (x+1)^{-2} - \dots - (x+n)^{-2} \geq 0.$$

Taking the limit as $n \rightarrow \infty$, it follows that

$$f''(x) \geq \zeta_2(x).$$

On the other hand, note that the difference $f(x) - \gamma_1(x)$ is periodic, of period 1. Hence its second derivative $f''(x) - \zeta_2(x)$ is periodic, and has average $\int_0^1 (f''(x) - \zeta_2(x))dx$ equal to zero. Clearly it follows that $f''(x) = \zeta_2(x)$ everywhere. Integrating twice, we see that

$$f(x) = \gamma_1(x) + ax + b.$$

Subtracting the corresponding equation for $f(x+1)$, we see that $a = 0$, which completes the proof. \square

This argument shows that

$$\gamma_1(x) = \log(\Gamma(x)/C)$$

for some constant C , whose precise value will be computed later.

Remark: The customary definition of the gamma function is the expression

$$\Gamma(x) = \int_0^\infty e^{-t} t^{x-1} dt$$

which was used in §2 and Appendix 1. Here is an outline proof that this expression does indeed satisfy the conditions of Lemma 15. Integration by parts shows that $\Gamma(x+1) = x\Gamma(x)$. Note that a twice differentiable positive function satisfies $(\log f(x))'' \geq 0$ if and only if the matrix

$$\begin{bmatrix} f(x) & f'(x) \\ f'(x) & f''(x) \end{bmatrix}$$

is positive semi-definite, for all x . But the collection of all 2×2 positive semi-definite matrices forms a convex cone. It follows that the sum $f(x) + g(x)$ of any two functions which satisfy this condition will also satisfy it. Similarly the integral

$$\begin{bmatrix} \Gamma(x) & \Gamma'(x) \\ \Gamma'(x) & \Gamma''(x) \end{bmatrix} = \int_0^\infty \begin{bmatrix} 1 & \log t \\ \log t & (\log t)^2 \end{bmatrix} e^{-t} t^{x-1} dt$$

is a positive semi-definite matrix. Hence $(\log \Gamma(x))'' \geq 0$ as required. \square

Now consider the Kubert identity

$$m^t \zeta_t(x) = \sum_0^{m-1} \zeta_t((x+k)/m).$$

If we differentiate both sides with respect to t , then substitute $t = 1 - s$ and $\zeta_t = -\beta_s/s$, we obtain

$$(22) \quad \gamma_s(x) = (\log m)\beta_s(x)/s + m^{s-1} \sum_0^m \gamma_s((x+k)/m).$$

Thus γ_s satisfies the Kubert identity $(*)_s$, except for a correction term involving the Bernoulli polynomial $\beta_s(x)$, for $s = 1, 2, 3, \dots$.

If we work modulo the logarithms of positive rational numbers, then the function

$$\mathbf{Q}/\mathbf{Z} \rightarrow \mathbf{R}/\mathbf{Q} \log \mathbf{Q}^+$$

induced by γ_s actually satisfies $(*)_s$. It seems natural to conjecture that this is a universal Kubert function on \mathbf{Q}/\mathbf{Z} for integers $s \geq 1$.

For $s = 1$, the "even" part of this conjecture can easily be proved using Bass' theorem, together with the classical identity

$$\gamma_1(x) + \gamma_1(1-x) + \log(2 \sin \pi x) = 0$$

for $0 < x < 1$, which is proved below, and the fact that $\gamma_1(1) = \log(1/\sqrt{2\pi})$ where π is transcendental. For the odd part of γ_1 , Rohrlich has conjectured universality even if we work modulo the logarithms of *all* algebraic numbers. See [17, p. 66].

In the case $s = 1$, formula (22) takes the form

$$(23) \quad \gamma_1(x) = (\log m) \left(x - \frac{1}{2} \right) + \sum_{k=0}^{m-1} \gamma_1((x+k)/m).$$

Hence the derivative $\gamma'_1(x) = \Gamma'(x)/\Gamma(x)$ satisfies

$$(24) \quad \gamma'_1(x) = \log m + m^{-1} \sum_{k=0}^{m-1} \gamma'_1((x+k)/m).$$

Note that $\gamma'_1(x+1) = \gamma'_1(x) + 1/x \equiv \gamma'_1(x) \pmod{\mathbf{Q}}$, if x is positive and rational. We may conjecture that γ'_1 induces a universal function $\mathbf{Q}/\mathbf{Z} \rightarrow \mathbf{R}/(\mathbf{Q} + \mathbf{Q} \log \mathbf{Q}^+)$ satisfying $(*)_0$. (It can be shown that $\gamma'_1(1)$ is equal to the negative of Euler's constant. Thus even at $x = 1$ the number theoretic properties of $\gamma'_1(x)$ are not known.)

As a typical application of (23), taking $x = 1$ we obtain the equation

$$\gamma_1(1/m) + \gamma_1(2/m) + \dots + \gamma_1((m-1)/m) = \log(1/\sqrt{m}).$$

In particular, $\gamma_1(1/2) = \log(1/\sqrt{2})$.

As a further application of (23), we will prove the classical formula

$$(25) \quad \gamma_1(x) + \gamma_1(1-x) + \log(2 \sin \pi x) = 0$$

for $0 < x < 1$. If we add (23) to the corresponding formula for $\gamma_1(1-x)$, then the correction terms cancel out. Hence the sum $\gamma_1(x) + \gamma_1(1-x)$ satisfies the Kubert identities $(*)_1$ in their original form. By Theorem 1, this implies that

$$\gamma_1(x) + \gamma_1(1-x) = c \log(2 \sin \pi x)$$

for some constant c . One way to evaluate c would be to differentiate twice:

$$\zeta_2(x) + \zeta_2(1-x) = -c\pi^2/\sin^2 \pi x,$$

and to note that both $\zeta_2(x)$ and $\pi^2/\sin^2 \pi x$ are asymptotic to $1/x^2$ as $x \rightarrow 0$. (Compare Appendix 1.) Another would be to substitute $x = 1/2$, noting that

$$\gamma_1(1/2) = -\frac{1}{2} \log 2 \text{ while } \log(2 \sin \pi/2) = \log 2. \text{ Using either method, one}$$

finds that $c = -1$, proving equation (25). \square

Next let us prove Lerch's identity (19). We showed during the proof of Lemma 15 that $\gamma_1(x) = \log(\Gamma(x)/C)$ for some constant $C > 0$. Exponentiating (25), we obtain

$$\frac{\Gamma(x)}{C} \frac{\Gamma(1-x)}{C} 2 \sin \pi x = 1.$$

Since

$$\Gamma(x) \sim x^{-1}, \quad \Gamma(1-x) \sim 1, \quad \text{and} \quad 2 \sin \pi x \sim 2\pi x$$

as $x \rightarrow 0$, it follows that $C = \sqrt{2\pi}$, as required. \square

This argument also proves the classical *Euler functional equation*

$$\Gamma(x)\Gamma(1-x) = \pi/\sin \pi x.$$

Taking $x = 1/2$, it proves that $\Gamma(1/2) = \sqrt{\pi}$.

Similarly, exponentiating (23), we obtain the classical *Gauss multiplication formula*

$$\frac{\Gamma(x)}{\sqrt{2\pi}} = m^{x-1/2} \prod_{k=0}^{m-1} \frac{\Gamma((x+k)/m)}{\sqrt{2\pi}}.$$

As an example, taking $x = 1$ and $m = 2$, we obtain another proof that $\Gamma(1/2) = \sqrt{\pi}$.

Note that each γ_{s+1} is essentially just an indefinite integral of γ_s , up to a constant factor and a polynomial summand. More precisely, differentiating the equation

$$\zeta'_t(x) = -t\zeta_{t+1}(x)$$

with respect to t and setting $s = -t$, we find that

$$(26) \quad \gamma'_{s+1}(x) = \partial \gamma_{s+1}(x) / \partial x = s\gamma_s(x) + \beta_s(x)/s.$$

The function $\exp(\gamma_s(x))$ can be thought of as a kind of higher order gamma function, satisfying

$$\exp(\gamma_s(n+1) - \gamma_s(1)) = 1^{1^{s-1}} 2^{2^{s-1}} \dots n^{n^{s-1}}.$$

(Compare Shintani [24].)

As a final remark, let us apply these methods to derive the Stirling asymptotic series for $\gamma_1(x)$ as $x \rightarrow \infty$. Using (26), together with (3) and (20), we have

$$\int_x^{x+1} \gamma_1(u) du = x \log x - x.$$

As in the discussion of Bernoulli polynomials in §2, the left side of this equation can be expanded as a Taylor series

$$\frac{e^D - I}{D} \gamma_1(x) = \sum_0^\infty D^n \gamma_1(x) / (n+1)!,$$

which converges whenever $\gamma_1(x)$ is analytic throughout a unit disk centered at x , or in other words whenever $x > 1$. Here D stands for d/dx . Recall from §2 that the inverse operator is given formally by

$$\frac{D}{e^D - I} = \sum_0^{\infty} b_n D^n / n!.$$

Hence, applying this inverse operator to both sides of the equation

$$\frac{e^D - I}{D} \gamma_1(x) = x \log x - x,$$

we might hope that

$$\gamma_1(x) \stackrel{?}{=} \frac{D}{e^D - I} (x \log x - x) = \sum_0^{\infty} b_n D^n (x \log x - x) / n!.$$

Unfortunately, this series does not converge. However, if we truncate, setting

$$s_N(x) = \sum_0^N b_n D^n (x \log x - x) / n!$$

for some integer $N \geq 1$, then we will prove that

$$\gamma_1(x) = s_N(x) + O(x^{-N})$$

as $x \rightarrow \infty$. This is the required asymptotic series. More explicitly, we can write it as

$$(27) \quad \gamma_1(x) = (x \log x - x) - \frac{1}{2} \log x + \sum_2^N \frac{b_n x^{1-n}}{n(n-1)} + O(x^{-N}).$$

(For a more precise description of the error term, see [1, p. 31]. Using (19) this yields the corresponding asymptotic formula for $\Gamma(x)$.)

To prove this formula, substitute the identity

$$x \log x - x = \sum_0^{\infty} \frac{D^m}{(m+1)!} \gamma_1(x)$$

in the definition of $s_N(x)$ to obtain a double series

$$s_N(x) = \sum_{n=0}^N \sum_{m=0}^{\infty} \frac{b_n D^n}{n!} \frac{D^m}{(m+1)!} \gamma_1(x),$$

which converges absolutely whenever $x > 1$. If we collect terms involving the same total power of D , then evidently all the terms involving D^1, D^2, \dots, D^N must cancel. Since

$$D^n \gamma_1(x) = \pm (n-1)! \zeta_n(x)$$

for $n \geq 2$, it follows that the resulting series has the form

$$s_N(x) = \gamma_1(x) + \sum_{n=N+1}^{\infty} a_n \zeta_n(x)$$

for suitable constants a_n . Setting

$$E(x) = \sum_{n=N+1}^{\infty} a_n x^{-n},$$

we can write the error term as

$$s_N(x) - \gamma_1(x) = E(x) + E(x+1) + \dots$$

Note that all of these series converge absolutely for $x > 1$. Evidently

$$E(x) = O(x^{-N-1})$$

as $x \rightarrow \infty$, for any fixed N , so

$$s_N(x) - \gamma_1(x) = O(x^{-N})$$

as required. \square

This argument yields similar asymptotic series for related functions such as $\zeta_s(x)$, $\gamma_s(x)$, and $\gamma'_s(x)$. Such estimates work also for complex values of x , as long as x stays well away from the negative real axis.

APPENDIX 3

VOLUME AND THE DEHN INVARIANT IN HYPERBOLIC 3-SPACE

We will describe some constructions in hyperbolic space involving the dilogarithm function $\mathcal{L}_2(z)$ and its Kubert identity (7). Further details may be found in the paper "Scissors Congruences, II" by J. L. Dupont and C.-H. Sah (*J. Pure Appl. Algebra* 25 (1982), 159-195).

Using the upper half-space model for hyperbolic 3-space, consider a totally asymptotic 3-simplex Δ . In other words, we assume that the vertices a, b, c, d of Δ all lie on the 2-sphere of points at infinity, which we identify with the extended complex plane $\mathbb{C} \cup \infty$. Then Δ is determined up to orientation preserving isometry by the cross ratio

$$z = (a, b; c, d) = (c-a)(d-b)/(c-b)(d-a).$$

[The semicolon is inserted in our cross ratio symbol as a remainder of its symmetry properties, which are similar to those of the four index symbol R_{hijk} in Riemannian geometry.] In particular, the volume of Δ can be expressed as a function of the cross ratio z .