

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 29 (1983)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON POLYLOGARITHMS, HURWITZ ZETA FUNCTIONS, AND THE KUBERT IDENTITIES
Autor: Milnor, John
Bibliographie
DOI: <https://doi.org/10.5169/seals-52983>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

Download PDF: 25.05.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Dupont and Sah show that the volume function and the sharpened Dehn invariant can be incorporated into a single function ρ , as follows. Let

$$\rho(z) = 1 \wedge L(z) - 1 \wedge L(1-z) + l(z) \wedge l(1-z),$$

with values in $\wedge^2 \mathbf{C}$, where $l(z) = \log(z)/2\pi i$ and

$$L(z) = \mathcal{L}_2(z)/4\pi^2 = \int_0^z l(1-t)dl(t).$$

This expression is certainly well defined in the strip $0 < \operatorname{Re}(z) < 1$, and satisfies $\rho(z) + \rho(1-z) = 0$. If we analytically continue each of its constituent functions in a loop around zero or one, then the expression $\rho(z)$ remains unchanged. Hence ρ is well defined as a mapping from $\mathbf{C} - \{0, 1\}$ to $\wedge^2 \mathbf{C}$. They show that ρ also satisfies the symmetry condition (31), the Kubert identity (32), and the cocycle equation (33).

REFERENCES

- [1] ARTIN, E. *The Gamma Function*. Holt, Rinehart and Winston 1964.
- [2] BASS, H. Generators and relations for cyclotomic units. *Nagoya Math. J.* 27 (1966), 401-407.
- [3] BATEMAN, H., edited by Erdélyi *et al.* *Higher Transcendental Functions, vol. I.* McGraw-Hill, 1953. (See pp. 24-39.)
- [4] BLOCH, S. Applications of the dilogarithm function in algebraic K -theory and algebraic geometry. *Proc. Conf. Algebraic Geom. Kyoto Univ.* 1977. (See also “Algebraic K -theory and zeta functions of elliptic curves”, *Proc. Int. Congr. Math. Helsinki* 1978, II, 511-515; and “The dilogarithm and extensions of Lie algebras”, preprint, U. Chicago 1981.)
- [5] BOREVICH, Z. I. and I. R. SHAFAREVICH. *Number Theory*. Academic Press 1966.
- [6] CHUDNOVSKY, G. V. Padé approximations to the generalized hypergeometric functions. I, *J. Math. pures appl.* 58 (1979), 445-476.
- [7] EWING, J. Spheres as fixed point sets. *Quart. J. Math., Oxford*, 27 (1976), 445-455.
- [8] HILBERT, D. Die Theorie der algebraischen Zahlkörper. *Jahresb. D. Math.-Ver.* 4 (1897), 175-546. (See §120.)
- [9] HURWITZ, A. Einige Eigenschaften der Dirichlet'schen Funktionen $F(s) = \sum \left(\frac{D}{n}\right) \frac{1}{n^s}$, die bei der Bestimmung der Klassenanzahlen binärer quadratischer Formen auftreten. *Zeits. Math. Phys.* 27 (1882), 86-101.
- [10] IWASAWA, K. *Lectures on p -Adic L -Functions*. Princeton University Press, Annals Studies 74 (1972).
- [11] JONQUIÈRE, A. Note sur le série $\sum x^n/n^s$. *Bull. Soc. Math. France* 17 (1889), 142-152.
- [12] KUBERT, D. The universal ordinary distribution. *Bull. Soc. Math. France* 107 (1979), 179-202.
- [13] KUBERT, D. and S. LANG. Units in the modular function field III. Distribution relations. *Math. Ann.* 218 (1975), 273-285.
- [14] —— Distributions on toroidal groups. *Math. Zeits.* 148 (1976), 33-51.
- [15] —— *Modular Units*. Grundlehren math. Wiss. 244, Springer, 1981.

- [16] LANG, S. Relations de distributions et exemples classiques. *Sémin. Delange-Pisot-Poitou* 19 (1977-78), n° 40.
- [17] ——— *Cyclotomic Fields*. Springer, 1978. (See pp. 32, 65.)
- [18] ——— *Cyclotomic Fields 2*. Springer, 1980. (See Ch. 17.)
- [19] LERCH, M. Note sur la fonction $\mathfrak{R}(w, x, s) = \sum_0^{\infty} \frac{e^{2knix}}{(w+k)^s}$. *Acta Math.* 11 (1887-88), 19-24.
- [20] LEWIN, L. *Polylogarithms and Associated Functions*. North-Holland, 1981.
- [21] MILNOR, J. Hyperbolic geometry : The first 150 years. *Bull. Amer. Math. Soc.* 6 (1982), 9-24.
- [22] RAMAKRISHNAN, D. A regulator for curves via the Heisenberg group. *Bull. Amer. Math. Soc.* 5 (1981), 191-195. (See also "On the monodromy of higher logarithms", preprint, U. Chicago, 1981.)
- [23] SERRE, J.-P. *A Course in Arithmetic*. Springer, 1973.
- [24] SHINTANI, T. On a Kronecker limit formula for real quadratic fields. *J. Fac. Sci. Univ. Tokyo, sect. IA Math.* 24 (1977), 167-199. (See p. 178.)
- [25] SINNOTT, W. On the Stickelberger ideal and the circular units of a cyclotomic field. *Annals of Math.* 108 (1978), 107-134.
- [26] TRUESDELL, C. On a function which occurs in the theory of the structure of polymers. *Annals of Math.* 46 (1945), 144-157.
- [27] WEIL, A. *Elliptic functions according to Eisenstein and Kronecker*. Springer, 1976. (See pp. 53-60.)

(Reçu le 11 novembre 1982)

John Milnor

Institute for Advanced Study
Princeton, New Jersey 08540
U.S.A.