Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 29 (1983)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE CLEBSCH-GORDAN FORMULAS
Autor: Flath, Daniel
Kapitel: 2. The Weyl algebra A
DOI: https://doi.org/10.5169/seals-52986

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 11.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot X = aX + cY \qquad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot Y = bX + dY \tag{1.1}$$

$$g \cdot X^a Y^b = (g \cdot X)^a (g \cdot Y)^b \qquad \text{for } g \in SL_2(\mathbf{C}) . \tag{1.2}$$

Each V_m is an $SL_2(\mathbf{C})$ subrepresentation of V.

By $\mathfrak{sl}_2$ we denote the Lie algebra of 2×2 complex matrices with trace 0. The representation of $SL_2(\mathbf{C})$ on V gives rise, through differentiation, to a representation of $\mathfrak{sl}_2$ on V.

$$L \cdot v = \frac{d}{dt}\bigg|_{t=0} \exp(tL) \cdot v \qquad \text{for } L \in \mathfrak{sl}_2, v \in V . \tag{1.3}$$

Choose a basis E_+, E_-, H of $\mathfrak{sl}_2$ as follows:

$$E_+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \qquad E_- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \qquad H = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \tag{1.4}$$

An easy calculation establishes the following equalities of linear endomorphisms of V.

$$E_+ = X\partial_Y , \qquad E_- = Y\partial_X , \tag{1.5}$$

$$H = X\partial_X - Y\partial_Y . \tag{1.6}$$

From (1.5) and (1.6) one easily deduces that each V_m is an *irreducible* representation of $\mathfrak{sl}_2$ (and of $SL_2(\mathbf{C})$).

We define for integers m, n a representation τ of $\mathfrak{sl}_2$ on $\mathrm{Hom}_{\mathbf{C}}(V_m, V_n)$ by means of formula (1.7).

$$(\tau(L) \cdot T)v = L(Tv) - T(Lv)$$
$$\text{for } L \in \mathfrak{sl}_2, T \in \mathrm{Hom}_{\mathbf{C}}(V_m, V_n), v \in V_m . \tag{1.7}$$

The principal result of this article is the explicit decomposition of the $\mathfrak{sl}_2$-representations $\mathrm{Hom}_{\mathbf{C}}(V_m, V_n)$.

2. The Weyl algebra $\mathscr{A}$

Let $\mathscr{A}$ be the subalgebra of $\mathrm{End}_{\mathbf{C}}(V)$ consisting of polynomial differential operators on $V = \mathbf{C}[X, Y]$. The algebra $\mathscr{A}$ is spanned by the elements

$$D(i, j, a, b) = X^i Y^j \partial_X{}^a \partial_Y{}^b . \tag{2.1}$$

The Euler operator J, which acts as scalar multiplication by m on V_m, lies in $\mathscr{A}$.

$$J = X\partial_X + Y\partial_Y . \tag{2.2}$$

The next lemma assures us that $\mathscr{A}$ is large enough for the study of all spaces $\mathrm{Hom}_{\mathbf{C}}(V_m, V_n)$.

LEMMA 2.3. *Let U be a finite dimensional vector subspace of V and let $T \in \mathrm{End}_{\mathbf{C}}(U)$. Then there exists an element of $\mathscr{A}$ whose restriction to U equals T.*

Proof: The element $S = X^c Y^d (\partial_X)^a (\partial_Y)^b \prod_{\substack{m=0 \\ m \neq a+b}}^{N} (J-m)$ of $\mathscr{A}$ maps $X^a Y^b$ to a nonzero multiple of $X^c Y^d$ and kills all other monomials of degree at most N. But by enlarging U we may assume that $\mathrm{End}_{\mathbf{C}}(U)$ is spanned by restrictions of elements of the form S. □

We use the inclusion of $\mathfrak{sl}_2$ in $\mathscr{A}$ to define a representation ρ of $\mathfrak{sl}_2$ on $\mathscr{A}$.

$$\rho(L)a = [L, a] \qquad \text{for } L \in \mathfrak{sl}_2, a \in \mathscr{A} . \tag{2.4}$$

For integers n let $\mathscr{A}^n$ be the set of T in $\mathscr{A}$ such that $T(V_m) \subset V_{m+n}$ for all m.

This defines a grading of $\mathscr{A}$ which is preserved by the action of $\mathfrak{sl}_2$.

$$\mathscr{A} = \bigoplus_{n \in \mathbf{Z}} \mathscr{A}^n , \qquad \mathscr{A}^m \cdot \mathscr{A}^n \subset \mathscr{A}^{m+n} , \tag{2.5}$$

$$\rho(L)\mathscr{A}^n \subset \mathscr{A}^n \qquad \text{for all } L \in \mathfrak{sl}_2 . \tag{2.6}$$

The algebra $\mathscr{A}$ and representation ρ have been defined just so that the next lemma, which is an immediate consequence of Lemma 2.3, will be true.

LEMMA 2.7. *For each m, n the restriction map*

$$\mathrm{res}: \mathscr{A}^n \to \mathrm{Hom}_{\mathbf{C}}(V_m, V_{m+n})$$

is a surjective homomorphism of $\mathfrak{sl}_2$ representations. □

The method of this paper is to deduce the decomposition of the representations $\mathrm{Hom}_{\mathbf{C}}(V_m, V_{m+n})$ from the decomposition of the representation ρ on $\mathscr{A}$ by means of Lemma 2.7.