Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 29 (1983)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE CLEBSCH-GORDAN FORMULAS
Autor: Flath, Daniel
Kapitel: 4. The decomposition of A
DOI: https://doi.org/10.5169/seals-52986

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 11.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

3. The theory of the highest weight

Before decomposing the $\mathfrak{sl}_2$-space $\mathscr{A}$ we must review the finite dimensional representation theory of $\mathfrak{sl}_2$.

The *weight vectors* of an $\mathfrak{sl}_2$-representation W are the eigenvectors of H in W. The *weights* of W are the eigenvalues of its nonzero weight vectors.

Every finite dimensional $\mathfrak{sl}_2$-module is spanned by its weight vectors. The weights of such a representation are all integers and are thus ordered by the usual order on $\mathbf{R}$. The largest of a finite set of integral weights is traditionally referred to as the *highest* weight.

Two finite dimensional irreducible $\mathfrak{sl}_2$-representations are isomorphic if and only if they have the same highest weights, which are necessarily nonnegative.

The element $X^a Y^b$ of V is a weight vector of weight a-b. This shows that X^m is a vector of highest weight m in V_m and therefore that the V_m for $m \geqslant 0$ form a set of representatives of the equivalence classes of finite dimensional irreducible $\mathfrak{sl}_2$-representations; which is precisely why we are studying them is this paper.

The last general fact which we will recall without proof is this: every finite dimensional representation of $\mathfrak{sl}_2$ is a direct sum of irreducible representations.

Given a representation W of $\mathfrak{sl}_2$ which is a sum of finite dimensional representations one often wishes to write it explicitly as a direct sum of irreducible representations, that is, of representations isomorphic to the V_m. A method for doing this is provided by the observation that the space of weight vectors of highest weight in V_m is the space annihilated by E_+ and is one dimensional. Thus for each $v \in W$ of weight m such that $E_+ v = 0$, there is a unique $\mathfrak{sl}_2$-homomorphism from V_m to W taking X^m to v. The explicit decomposition of W therefore amounts to the determination of a basis consisting of weight vectors of the kernel of E_+ in W.

4. The decomposition of $\mathscr{A}$

We apply the procedure of the last paragraph to the representation of $\mathfrak{sl}_2$ on $\mathscr{A}$. By definition of ρ the kernel of $\rho(E_+)$ is just the commutant of E_+ in $\mathscr{A}$.

Let $\mathscr{B}$ be the subalgebra of $\mathscr{A}$ generated by X, ∂_Y, and J.

PROPOSITION 4.1. *$\mathscr{B}$ is the commutant of E_+ in $\mathscr{A}$.*

Proof: One easily verifies that E_+ commutes with X, ∂_Y, and J, which shows that $\mathscr{B}$ is contained in the commutant of E_+.

Let U be the $\mathfrak{sl}_2$-subrepresentation of $\mathscr{A}$ generated by $\mathscr{B}$. The considerations of Section 3 show that the inclusion of the commutant of E_+ in $\mathscr{B}$ is equivalent to the assertion that U equals all of $\mathscr{A}$. We proceed to establish that equality.

The algebra $\mathscr{B}$ is spanned as a vector space by the elements

$$J^a X^b (\partial_Y)^c \qquad \text{with } a, b, c \geqslant 0 . \tag{4.2}$$

We present two calculations.

$$\begin{aligned} &[E_-, J^a X^b (\partial_Y)^{c+1}] \\ &= -(b+c+1) J^a X^b (\partial_Y)^c \partial_X + b(J+1-b) J^a X^{b-1} (\partial_Y)^c \end{aligned} \tag{4.3}$$

$$\begin{aligned} &[E_-, J^a X^{b+1} (\partial_Y)^c] \\ &= (b+1) J^a X^b (\partial_Y)^c Y - c J^a X^b (\partial_Y)^{c-1} (b+1+X\partial_X) . \end{aligned} \tag{4.4}$$

From (4.3) one concludes that $\mathscr{B} \cdot \partial_X \subset U$. From that and (4.4) one concludes that $\mathscr{B} \cdot Y \subset U$.

Because E_- commutes with ∂_X and Y, one has that

$$\rho(E_-)^n (\mathscr{B} \partial_X) = \big(\rho(E_-)^n \mathscr{B}\big) \cdot \partial_X$$

and that

$$\rho(E_-)^n (\mathscr{B} \cdot Y) = \big(\rho(E_-)^n \mathscr{B}\big) \cdot Y .$$

Because $V_m = \bigoplus_{n=0}^{\infty} E_-{}^n(\mathbf{C}X^m)$ one knows that $U = \bigoplus_{n=0}^{\infty} \rho(E_-)^n \mathscr{B}$. And thus

$$U \cdot \partial_X \subset U , \qquad U \cdot Y \subset U . \tag{4.5}$$

Iterating, we have

$$U Y^d (\partial_X)^e \subset U \qquad \text{for } d, e, \geqslant 0 . \tag{4.6}$$

But $\mathscr{A}$ is generated as an algebra by X, Y, ∂_X, and ∂_Y and so (4.2) and (4.6) prove that $U = \mathscr{A}$. □

COROLLARY 4.7. *$\mathscr{A}^0$ is the subalgebra of $\mathscr{A}$ generated by $\mathfrak{sl}_2$ and J.*

Proof: $\mathscr{A}^0$ is the $\mathfrak{sl}_2$-subrepresentation of $\mathscr{A}$ generated by $\mathscr{A}^0 \cap \mathscr{B}$.

$\mathscr{A}^0 \cap \mathscr{B}$ is spanned by the elements (4.2) such that $b = c$, all of which are of the form $J^a E_+{}^b$. □

We remark that the subalgebra of $\mathscr{A}$ generated by $\mathfrak{sl}_2$ is canonically isomorphic to the universal enveloping algebra of $\mathfrak{sl}_2$. The element $J(J+2)$ equals $H^2 + 2(E_+E_- + E_-E_+)$, the Casimir element for $\mathfrak{sl}_2$. Thus $\mathscr{A}^0$ is a little larger than the enveloping algebra of $\mathfrak{sl}_2$.

For integers l, n define $\mathscr{B}\binom{n}{l}$ to be the set of $T \in \mathscr{B} \cap \mathscr{A}^n$ such that $\rho(H)T = lT$.

This defines a grading of $\mathscr{B}$:

$$\mathscr{B} = \oplus\, \mathscr{B}\binom{n}{l}, \qquad \mathscr{B}\binom{n}{l} \cdot \mathscr{B}\binom{n'}{l'} \subset \mathscr{B}\binom{n+n'}{l+l'}. \tag{4.8}$$

The generators of $\mathscr{B}$ fit in as follows:

$$J \in \mathscr{B}\binom{0}{0}, \qquad X \in \mathscr{B}\binom{1}{1} \qquad \partial_Y \in \mathscr{B}\binom{-1}{1}. \tag{4.9}$$

PROPOSITION 4.10. *i)* $\mathscr{B}\binom{0}{0} = \mathbf{C}[J]$.

ii) $\mathscr{B}\binom{n}{l} \neq 0$ *if and only if* $l \geqslant 0, |n| \leqslant l$, *and* $l \equiv n \pmod 2$. *If these conditions are met, then*

$$\mathscr{B}\binom{n}{l} = \mathbf{C}[J] \cdot X^{\frac{l+n}{2}} (\partial_Y)^{\frac{l-n}{2}} \tag{4.11}$$

Proof: Immediate. □

We note that the condition that $\mathscr{B}\binom{n}{l} \neq (0)$ may be rephrased thus: $l \geqslant 0$ and n is a weight of V_l.

5. DECOMPOSITION OF $\mathrm{Hom}(V_m, V_{m+n})$

THEOREM 5.1. *Let* l, m, n *be integers with* $l, m, m+n \geqslant 0$. *There is an* $\mathfrak{sl}_2$*-subrepresentation of* $\mathrm{Hom}_{\mathbf{C}}(V_m, V_{m+n})$ *which is isomorphic to* V_l *if and only if* $|n| \leqslant l, n \equiv l \pmod 2$, *and* $m \geqslant \dfrac{l-n}{2}$.