

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 29 (1983)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SOME PARADOXICAL SETS WITH APPLICATIONS IN THE GEOMETRIC THEORY OF REAL VARIABLE
Autor: de Guzmán, Miguel
Kapitel: 4. A SMALL TREE WITH MANY FRUITS
DOI: <https://doi.org/10.5169/seals-52969>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

3. A TOURIST COLONY NOT TO BE RECOMMENDED

In 1927 Nikodym, in order to explore the geometric structure of the measurable sets in the plane, showed how to construct, inside a square Q , a set N that fills it (i.e. the measure of $Q-N$ is zero) and so that for each point x of N there is a straight line $l(x)$ passing through it and not hitting any other point of N (Fig. 4).




FIGURE 4

The architects of tourist colonies have not yet learned about this magnificent business possibility, but the day somebody tells you about the marvels of a colony in an island which offers a free view over the ocean from each one of its apartments, beware!

Although it seems incredible one can still make it better. R. O. Davies in 1953 constructed a set N in Q filling Q and such that each x of N has infinitely many directions in which one can see the ocean... inside any arbitrarily small angle one may fix!

4. A SMALL TREE WITH MANY FRUITS

In 1928 Besicovitch was informed about the needle problem and published its solution. In 1929 Perron simplified the somewhat laborious construction of Besicovitch. It has been further simplified later on. The final product of the line of

thought, that we shall call *the Perron tree*, has proved to be an extraordinarily fruitful tool for the solution of certain deep problems of recent mathematical analysis.

The result is as follows: Given an arbitrary $\varepsilon > 0$ and an arbitrary triangle ABC of area that we denote by $S(ABC)$, we can divide the triangle ABC into small triangles T_1, T_2, \dots, T_n as Figure 5 shows (i.e. dividing the basis a into a finite number of equal intervals I_1, I_2, \dots, I_n) and one can translate appropriately the small triangles T_1, T_2, \dots, T_n parallelly to the basis a in such a way that the area of the union of the translated triangles is less than $\varepsilon S(ABC)$. (See Fig. 6.)




FIGURE 5




FIGURE 6

5. HOW THE PERRON TREE SPROUTS

Following an idea of Rademacher (1962), the construction of the Perron tree can be easily understood as follows. Let us divide first a triangle T, MNP , of area $S(T)$, into two triangles T_1, T_2 , with bases J_1, J_2 , of the same length. If we wish to move T_1 and T_2 , parallelly to NP so that the shifted triangles cover less area we can do it by pushing T_2 towards T_1 as Figure 7 shows. The area covered by T_1 and T'_2 can be easily measured by elementary geometry and is (see Fig. 7, we take $1/2 < \alpha < 1$)

$$\alpha^2 S(T) + 2(1 - \alpha)^2 S(T)$$