

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 29 (1983)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SOME PARADOXICAL SETS WITH APPLICATIONS IN THE GEOMETRIC THEORY OF REAL VARIABLE
Autor: de Guzmán, Miguel
Kapitel: 5. How the Perron tree sprouts
DOI: <https://doi.org/10.5169/seals-52969>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

thought, that we shall call *the Perron tree*, has proved to be an extraordinarily fruitful tool for the solution of certain deep problems of recent mathematical analysis.

The result is as follows: Given an arbitrary $\varepsilon > 0$ and an arbitrary triangle ABC of area that we denote by $S(ABC)$, we can divide the triangle ABC into small triangles T_1, T_2, \dots, T_n as Figure 5 shows (i.e. dividing the basis a into a finite number of equal intervals I_1, I_2, \dots, I_n) and one can translate appropriately the small triangles T_1, T_2, \dots, T_n parallelly to the basis a in such a way that the area of the union of the translated triangles is less than $\varepsilon S(ABC)$. (See Fig. 6.)




FIGURE 5




FIGURE 6

5. HOW THE PERRON TREE SPROUTS

Following an idea of Rademacher (1962), the construction of the Perron tree can be easily understood as follows. Let us divide first a triangle T, MNP , of area $S(T)$, into two triangles T_1, T_2 , with bases J_1, J_2 , of the same length. If we wish to move T_1 and T_2 , parallelly to NP so that the shifted triangles cover less area we can do it by pushing T_2 towards T_1 as Figure 7 shows. The area covered by T_1 and T'_2 can be easily measured by elementary geometry and is (see Fig. 7, we take $1/2 < \alpha < 1$)

$$\alpha^2 S(T) + 2(1-\alpha)^2 S(T)$$




FIGURE 7

If the triangle MNP is divided into four parts, instead of two, as Figure 8 indicates, we can first subject the pair of triangles MNL_1 and ML_1L_2 on the one hand to the above indicated operation with an α , $1/2 < \alpha < 1$, and, on the other hand we can do the same, with the same α , to the other pair of adjacent triangles ML_2L_3 , ML_3P . The result is indicated in Figure 8.




FIGURE 8

It is easy to see that the area of the figure now covered by the so translated triangles is less than

$$(*) \quad \alpha^2 S(T) + 2(1-\alpha)^2 S(T)$$

If we now shift in a solidary way the figure formed by the union of the two triangles T_3 and T'_4 towards the left until L_2 coincides with L'_2 , the new formed figure covered by the four triangles can be considered (see Fig. 9) as consisting of

a triangle HNP'' similar to the first one MNP with a similarity ratio α plus four peak triangles that overlap more than before. The area of this figure is therefore less than (*).




FIGURE 9

In the triangle HNP'' we have the basis divided into equal portions NL'_2 and L'_2P'' and so we can submit HNP'' to the initial operation, i.e. shifting the right hand triangle towards the left one with the same constant α that measures the magnitude of this shift and shifting thereby solidarily the triangles T''_3 and T''_4 that constitute the right hand portion of the triangle HNP'' . The result is shown in Figure 10.




FIGURE 10

The final result is a triangle similar to the initial one with similarity ration α^2 , its area therefore being $\alpha^4 S(T)$, plus four peaks that cover an area smaller than

$$2(1-\alpha)^2 \alpha^2 S(T) + 2(1-\alpha)^2 S(T)$$

i.e. the total area now covered is not greater than

$$\alpha^4 S(T) + 2(1-\alpha)^2 (1+\alpha^2) S(T)$$

It is now not difficult to realize that if we initiate our process with 2^n equal portions of the basis and we proceed in a similar way, at the end, i.e. after n repetitions of the process consisting of (a) a shift of the right triangle of each pair of adjacent triangles towards the left triangle (shift constant = α), and (b) gluing together the resulting figures to compose a triangle similar to the original one with one half the number of divisions on the basis, we obtain a figure with an area not greater than

$$\begin{aligned} & \alpha^{2n} S(T) + 2(1-\alpha)^2 (1+\alpha^2 + \dots + \alpha^{2n-2}) S(T) \\ & \leq \alpha^{2n} S(T) + 2(1-\alpha)^2 (1+\alpha^2 + \dots + \alpha^{2n-2} + \dots) S(T) \\ & = \left(\alpha^{2n} + \frac{2(1-\alpha)^2}{1-\alpha^2} \right) S(T) \leq (\alpha^{2n} + 2(1-\alpha)) S(T) \end{aligned}$$

Therefore, given $\varepsilon > 0$, we first choose α , $1/2 < \alpha < 1$ such that $1 - \alpha < \varepsilon/2$ and then n so that $\alpha^{2n} < \varepsilon/2$. In this way we obtain a Perron tree. Its name is justified by the fact that the final figure consists of a trunk (a triangle similar to the initial one with similarity ration α^n) plus many sharp branches that seem to rest on it. Its area is less than $\varepsilon S(T)$. (See Fig. 11.)




FIGURE 11

The reader interested in more details can consult Guzmán (1975).