L'Enseignement Mathématique
Commission Internationale de l'Enseignement Mathématique
29 (1983)
1-2: L'ENSEIGNEMENT MATHÉMATIQUE
SOME PARADOXICAL SETS WITH APPLICATIONS IN THE GEOMETRIC THEORY OF REAL VARIABLE
de Guzmán, Miguel
7. The construction of the Besocovitch set
https://doi.org/10.5169/seals-52969

## Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

## **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

## Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

hand side of  $T'_1$ . Therefore it can move, using the above construction, covering an area J. Within  $T'_2$ , and so within P, it can move to the right hand side of  $T'_2$ . From there to the left hand side of  $T'_3$  and so on until it comes to AC, covering area less than  $\eta/3$ .

It is clear that with three equilateral triangles and three repetitions of this process we can turn the needle around covering area smaller than  $\eta$ .

## 7. The construction of the Besocovitch set

The Besicovitch set is also easily built starting from the Perron tree by means of the following auxiliary construction:

(\*\*) Given an arbitrary parallelogram ABCD and  $\varepsilon > 0$ , it is possible to construct a finite number of closed parallelograms  $\omega_1, \omega_2, ..., \omega_n$  so that (see Fig. 13):



FIGURE 13

- (a) Each one has one basis on AB and another one on CD.
- (b) The area of their union is less than  $\varepsilon$ .
- (c) For each segment joining a point of AB to another one of CD there exists inside some  $\omega_j$  a segment parallel to it of the same length.

To see this, given ABCD and  $\varepsilon > 0$  we first take two strips  $\omega_1$  and  $\omega_2$  as indicated in Figure 14 such that  $S(\omega_1) + S(\omega_2) < \varepsilon/4$ . We take now a point L of UV so that LC is parallel to UT. Then we divide VC into intervals with the same length smaller than that of DV and we join L to the extreme points of these intervals. A typical triangle of the ones so obtained is LMN. Let p be



FIGURE 14

the number of such triangles. From LMN we construct a Perron tree with area less than  $\varepsilon/4p$  and the same is done with each of the p triangles we have. The union of  $\omega_1$ ,  $\omega_2$  and all the small triangles of the p Perron trees has an area less than  $\varepsilon/4 + p \varepsilon/4p = \varepsilon/2$ . One of these small triangles composing one of the Perron trees has the situation indicated in Figure 15 (it has been enlarged to make the figure more easily understandable).



FIGURE 15

We wish to substitute the trapezoid HKRW by strips, what we can easily do without augmenting the area in the way schematically indicated in Figure 15, where four parallelograms have sufficed to cover the portion of *ABCD* covered by *HKRW*.

So we obtain  $\omega_1$ ,  $\omega_2$  and a number of small strips covering together an area smaller than  $\varepsilon/2$ . It is easy to see that for each segment joining A to a point of DC, there is another one parallel to it of the same length inside one of the strips we have obtained. If we now perform an analogous construction starting from the other side BC of ABCD we obtain a finite number of strips satisfying all the properties indicated (a), (b), (c).

The Besicovitch set is now very easily obtained as follows. We take a square MNPQ of side length 1 and apply to it the above auxiliary construction (\*\*) with  $\varepsilon = 1/2$ . We obtain a number of strips  $\omega_1^1, \omega_2^1, ..., \omega_{r_1}^1$ , covering an area smaller than 1/2 and such that for each segment determined by a point of MN and another of PQ there is a segment of the same length and direction inside  $\Omega^1 = \omega_2^1 \cup \omega_2^1 \cup ... \cup \omega_{r_1}^1$ .

Now we consider each of the parallelograms  $\omega_j^1$  and apply to it the same construction (\*\*) with  $\varepsilon = 1/2^2 r_1$ . Collecting all parallelograms corresponding to each  $\omega_j^1$ ,  $j = 1, 2, ..., r_1$ , we obtain a second family of parallelograms  $\omega_1^2, \omega_2^2, ..., \omega_{r_2}^2$ . Their union  $\Omega^2 = \omega_1^2 \cup \omega_2^2 \cup ... \cup \omega_{r_2}^2$  has area less than 1/2, is contained in  $\Omega^1$  and, again, for each segment joining a point of MN to another of PQ there is another one of the same length and direction inside  $\Omega^2$ . We proceed with the parallelograms  $\omega_j^2$  as we did with the  $\omega_j^1$ , now with  $\varepsilon = 1/2^3 r_2$ , and so on. Thus we obtain

,  $\Omega^1 \supset \Omega^2 \supset \Omega^3 \supset ...$ 

of areas

$$S(\Omega^1) < 1/2, S(\Omega^2) < 1/2^2, S(\Omega^3) < 1/2^3, \dots$$

The sets  $\Omega^{j}$  are compact and have the property of containing a parallel translation of each segment with one extremity on MN and the other on PQ. The intersection

$$B = \Omega^1 \cap \Omega^2 \cap ... \cap \Omega^j \cap ...$$

is of null area and has this same property. We now proceed with the square MNPQ in the same way in the other direction and obtain a compact set of measure zero containing a segment of length one in each direction, i.e. the Besicovitch set.

# 8. The Nikodym set

The Nikodym set can be obtained from the Perron tree in a similar way through the following auxiliary construction, also surprising in itself.