

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 29 (1983)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: REPRESENTATIONS OF THE SYMMETRIC GROUP, THE SPECIALIZATION ORDER, SYSTEMS AND GRASSMANN MANIFOLDS
Autor: Hazewinkel, Michiel / Martin, Clyde F.
Kapitel: 4. Schubert Cells
DOI: <https://doi.org/10.5169/seals-52973>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

the point of $\mathbf{G}_n(\mathbf{C}^{n+m})$ represented by $\text{Ker}(V \rightarrow E(x))$. This gives a holomorphic map $\Psi_E: M \rightarrow \mathbf{G}_n(\mathbf{C}^{n+m})$ such that the pullback of ξ_m by means of Ψ_E is isomorphic to E , $\Psi_E^! \xi_m \simeq E$. It is universality properties such as this one which account for the importance of the bundles ξ_m and η_n in differential and algebraic topology [16], algebraic geometry and also system and control theory (cf. [22, 23] and the references therein for the last mentioned).

The bundle ξ_m has a number of obvious holomorphic sections, viz. the sections defined by $\varepsilon_i(x) = e_i \bmod x$ where e_i is the i -th standard basis vector of \mathbf{C}^{n+m} , $i = 1, \dots, n+m$. And, as a matter of fact, it is not difficult to show that $\Gamma(\xi_m, \mathbf{G}_n(\mathbf{C}^{n+m}))$ is $(n+m)$ -dimensional and that the $\varepsilon_1, \dots, \varepsilon_{n+m}$ form a basis for this space of holomorphic sections; cf. subsection 8.1 below.

4. SCHUBERT CELLS

4.1. *Schubert Cells.* Consider again the Grassmann manifold $\mathbf{G}_n(\mathbf{C}^{m+n})$. Let $\underline{A} = (A_1, \dots, A_n)$ be a sequence of n -subspaces of \mathbf{C}^{m+n} such that $0 \neq A_1 \subset A_2 \subset \dots \subset A_n$ with each containment strict. To each such sequence \underline{A} we associate the closed subset

$$(4.2) \quad SC(\underline{A}) = \{x \in \mathbf{G}_n(\mathbf{C}^{m+n}) \mid \dim(x \cap A_i) \geq i\}$$

and call it the closed Schubert-cell of the sequence \underline{A} . In particular if

$$0 < \gamma_1 < \gamma_2 < \dots < \gamma_n \leq n + m$$

is a strictly increasing sequence of natural numbers less than or equal to $n + m$ then we define (setting $\gamma = (\gamma_1, \dots, \gamma_n)$)

$$(4.3) \quad SC(\gamma) = SC(\mathbf{C}^{\gamma_1}, \dots, \mathbf{C}^{\gamma_n})$$

where \mathbf{C}^r is viewed as the subspace of all vectors in \mathbf{C}^{n+m} whose last $n + m - r$ coordinates are zero.

4.4 *Flag Manifolds and the Bruhat Decomposition.* A flag in \mathbf{C}^{n+m} is a sequence of subspaces $\underline{F} = F_1 \subset \dots \subset F_{n+m} \subset \mathbf{C}^{n+m}$ such that $\dim F_i = i$. Let $Fl(\mathbf{C}^{n+m})$ denote the analytic manifold of all flags in \mathbf{C}^{n+m} . There is a natural holomorphic mapping $Fl(\mathbf{C}^{n+m}) \rightarrow \mathbf{G}_n(\mathbf{C}^{n+m})$ given by associating to a flag \underline{F} its n -th element F_n . The flag manifold can be seen as the space of all cosets Bg , $g \in \mathbf{GL}_{n+m}(\mathbf{C})$ where B is the Borel subgroup of all lower triangular matrices

where the last * in each row is nonzero. The closure of this open Schubert-cell is the Schubert-cell $SC(\gamma)$ defined in (4.3) above.

One easily checks that

$$(4.8) \quad SC(\mu) \subset SC(\gamma) \leftrightarrow \mu_i \leq \gamma_i, i = 1, \dots, n$$

and this order on the Schubert cells $SC(\gamma)$, or the equivalent ordering on n -tuples of natural numbers, is therefore a quotient of the Bruhat order on the Weyl group S_{n+m} . It is the induced order on the set of cosets $(S_n \times S_m)\sigma$, $\sigma \in S_{n+m}$. (Obviously if $\tau \in S_n \times S_m$, then $\tau\sigma(e_{\gamma_i}) \in \{e_1, \dots, e_n\}$ if $\sigma(e_{\gamma_i}) \in \{e_1, \dots, e_n\}$.) (And inversely the Bruhat order is determined by the associated orders of Schubert cells in the sense that $\sigma > \tau$ in S_n iff for all $k = 1, \dots, n - 1$ we have for the associated Schubert cells in $G_k(\mathbb{C}^n)$ that $SC(\sigma) \supset SC(\tau)$; this is a rather efficient way of calculating the Bruhat order on the Weyl group S_n .)

5. INTERRELATIONS

Now that we have defined the concepts we need we can start to describe some interrelations between the various manifestations of the specialization order we discussed in section 2 above.

5.1. *Overview of the Various Relations.* A schematic overview of the various interconnections is given by the following diagram. In this diagram we

