Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 29 (1983)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: REPRESENTATIONS OF THE SYMMETRIC GROUP, THE SPECIALIZATION ORDER, SYSTEMS AND GRASSMANN MANIFOLDS

Autor: Hazewinkel, Michiel / Martin, Clyde F.

Kapitel: 8. VECTORBUNDLES AND SYSTEMS

DOI: https://doi.org/10.5169/seals-52973

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 11.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Then using the results above one shows that

$$\underline{t}\,\underline{s}\overline{(0(\kappa))} = \overline{0(\kappa)},\ \underline{s}\,\underline{t}\overline{(U(\kappa))} = \overline{U(\kappa)}$$

so that $\underline{t}$ and $\underline{s}$ set up a bijective correspondence between the closures of orbits in the two cases and hence induce a bijective order preserving correspondence between the sets of orbits themselves.

8. Vectorbundles and systems

This section contains a modified version of the construction of Hermann-Martin [14] associating a vectorbundle $E(\Sigma)$ over the Riemann sphere $\mathbf{P}^1(\mathbf{C})$ to every $\Sigma = (A, B) \in L^{cr}_{m,n}$. This version makes it almost trivial to see that $E(\Sigma)$ splits as a direct sum of line bundles $L(\kappa_i)$, $i = 1, ..., m$ where $\kappa = (\kappa_1, ..., \kappa_m)$ is the set of Kronecker indices of Σ.

The first thing needed is some more information on the universal bundle ξ_m.

8.1. *On the Universal Bundle* $\xi_m \to \mathbf{G}_n(\mathbf{C}^{n+m})$. Let V be a complex $n + m$ dimensional vector space and $V^* = \mathrm{Hom}_{\mathbf{C}}(V, \mathbf{C})$ its dual vector space. Given $x \in \mathbf{G}_n(\mathbf{C}^{n+m})$ define $x^* = \{y \in V^* \mid < y, v > = 0 \text{ for all } x \in V\}$ where $< , >$ denotes the usual pairing $V^* \times V \to \mathbf{C}$. Then x^* is m-dimensional and $x \mapsto x^*$ defines a holomorphic isomorphism

$$d : \mathbf{G}_n(V) \to \mathbf{G}_m(V^*)\,. \tag{8.2}$$

Now $v \in V/x$ defines a unique homomorphism $v^T : x^* \to \mathbf{C}$ as follows:

$v^T(a) = < a, \tilde{v} >$ for all $a \in x^*$, where $\tilde{v} \in V$ is any representative of v. This is well defined because $< a, b > = 0$ for all $b \in x$ if $a \in x^*$. This defines an isomorphism between the pullback $(d^{-1})\,\xi_m$ and the dual of the subbundle η_m on $G_m(V^*)$ defined by

$$\eta_m = \{(x^*, w) \in \mathbf{G}_m(V^*) \times V^* \mid w \in x^*\}$$

It follows that ξ_m is a subbundle of an $n + m$ dimensional trivial bundle $\mathbf{G}_n(\mathbf{C}^{n+m}) \times \mathbf{C}^{n+m}$. Because $\mathbf{G}_n(\mathbf{C}^{n+m})$ is projective (compact) all holomorphic maps $\mathbf{G}_n(\mathbf{C}^{n+m}) \to \mathbf{C}$ are constant so that the space of holomorphic sections $\Gamma(\mathbf{G}_n(\mathbf{C}^{n+m}) \times \mathbf{C}^{n+m}, \mathbf{G}_n(\mathbf{C}^{n+m}))$ is of dimension $n + m$. As a subbundle of a trivial $(n+m)$-dimensional bundle ξ_m can therefore have at most $(n+m)$ linearly

independent holomorphic sections. But we have already found $(n+m)$ linearly independent sections viz. the $\varepsilon_1, \ldots, \varepsilon_{n+m}$ defined by $\varepsilon_i(x) = e_i \bmod x$ where e_i is the i-th standard basis vector of $\mathbf{C}^{n+m}$. Therefore

$$\dim \Gamma(\xi_m, \mathbf{G}_n(\mathbf{C}^{n+m})) = n + m \tag{8.3}$$

Now let $A \in \mathbf{GL}_{n+m}(\mathbf{C})$. Then A induces a holomorphic automorphism A^* of $\mathbf{G}_m(\mathbf{C}^{n+m})$ defined by $x \mapsto Ax$. Then, of course, there is an induced isomorphism $A^{-1} : \mathbf{C}^{n+m}/Ax \to \mathbf{C}^{n+m}/x$ which for varying x induces an isomorphism

$$A \overset{!}{*} \xi_m \simeq \xi_m, \; A \in \mathbf{GL}_{n+m}(\mathbf{C}) \tag{8.4}$$

8.5. *The Line Bundles $L(i)$ over $\mathbf{P}^1(\mathbf{C})$.* The Riemann sphere $\mathbf{P}^1(\mathbf{C}) = S^2$ can be obtained by gluing together two copies of $\mathbf{C}$ along the open subsets $\mathbf{C}\backslash\{0\}$ by the isomorphism

$$\mathbf{C}\backslash\{0\} \to \mathbf{C}\backslash\{0\}, \; s \mapsto t = s^{-1}$$

A line bundle over $\mathbf{P}^1(\mathbf{C})$ is then obtained by giving a holomorphic isomorphism $\mathbf{C}\backslash\{0\} \times \mathbf{C} \to \mathbf{C}\backslash\{0\} \times \mathbf{C}$ linear in the second variable compatible with the above isomorphism. Obviously the only possibilities are $(s, v) \to (s^{-1}, s^i v)$ for $i \in \mathbf{Z}$. This gives us the following commutative diagram identifications

$$\begin{tikzcd}
\mathbf{C} \times \mathbf{C} \arrow[d] & \supset \mathbf{C}\backslash\{0\} \times \mathbf{C} \arrow[d] \arrow[rr, "{(s,v) \to (s^{-1}, s^i v)}", "{s \to s^{-1} = t}"'] & & \mathbf{C}\backslash\{0\} \times \mathbf{C} \arrow[d] & \subset \mathbf{C} \times \mathbf{C} \arrow[d] \\
\mathbf{C} \arrow[u, dashed, "s_1"] & \supset \mathbf{C}\backslash\{0\} \arrow[rr] & & \mathbf{C}\backslash\{0\} & \subset \mathbf{C} \arrow[u, dashed, "s_2"']
\end{tikzcd}$$

The corresponding holomorphic line bundle is denoted $L(-i)$. A section of $L(-i)$ consists of two holomorphic mappings s_1, s_2 of the form $s \to (s, f(s))$, $t \to (t, g(t))$ such that $s^i f(s) = g(s^{-1})$. It readily follows that $f(s)$ must be a polynomial of degree $\leqslant -i$. Thus

$$\dim \Gamma(L(i)) = 0 \qquad \text{if } i < 0 \tag{8.6}$$

$$\dim \Gamma(L(i)) = i + 1 \qquad \text{if } i \geqslant 0 \tag{8.7}$$

8.8. *The (modified) Hermann-Martin vectorbundle of a system.* Let $\Sigma = (A, B)$ be a pair of real or complex matrices of sizes $n \times n$ and $n \times m$. Then (A, B) is completely reachable (cr) iff the $n \times (n+m)$ matrix $(sI - A\,;\, B)$ is of rank n for all complex values of s. So if $\Sigma = (A, B)$ is cr one can define a holomorphic map ψ_Σ by

(8.9) $\quad \psi_\Sigma : \mathbf{P}^1(\mathbf{C}) \to \mathbf{G}_n(\mathbf{C}^{n+m}),\ s \mapsto \mathrm{Row}(sI - A\,;\, B),\ \infty \mapsto \mathrm{Row}(I\,;\,0)$

where $\mathrm{Row}(M)$ for an $n \times (m+n)$ matrix M denotes the subspace of $\mathbf{C}^{n+m}$ generated by the rows of M. The vectorbundle $E(\Sigma)$ over $\mathbf{P}^1(\mathbf{C})$ is now defined by

(8.10) $$E(\Sigma) = \psi_\Sigma^{!}\xi_m$$

8.11. *Proposition.* $E(\Sigma)$ depends only on the feedback orbit of Σ.

Indeed one easily checks that $\Sigma = (A, B)$, $\Sigma' = (A', B') \in L^{cr}_{m,n}$ are feedback equivalent (cf. 2.6 above) iff there are constant invertible matrices P, Q such that

$$P(sI - A\,;\, B)Q = (sI - A'\,;\, B')\,.$$

Now $\mathrm{Row}(PM) = \mathrm{Row}(M)$ and postmultiplication with Q changes ψ_Σ to $Q_* \circ \psi_\Sigma$ and

$$E(\Sigma') = \psi_\Sigma^{!}(\xi_m) = \psi_\Sigma^{!}(Q_*^{!}\xi_m) \simeq \left(\psi_\Sigma^{!}(\xi_m) = E(\Sigma)\right)$$

by 8.4 above, proving the proposition.

Thus to determine $E(\Sigma)$ we can assume that $\Sigma = (A, B)$ is in Brunowsky canonical form which means that A, B takes the form

$$\left[\begin{array}{ccc|ccc|ccc}
0 & 1 & 0 & & & & & & \\
 & & & & 0 & & & 0 & \\
 & & 1 & & & & & & \\
0 & & 0 & & & & & & \\
\hline
 & & & 0 & 1 & 0 & & & \\
 & & & & & & & 0 & \\
 & & & & & 1 & & & \\
 & & & 0 & & 0 & & & \\
\hline
 & & & & & & 0 & 1 & 0 \\
 & 0 & & & 0 & & & & \\
 & & & & & & & & 1 \\
 & & & & & & 0 & & 0
\end{array}\right]
\quad
\begin{array}{c}
\left[\begin{array}{ccc}
0 & 0 & 0 \\
 & & \\
0 & & \\
1 & 0 & 0 \\
\hline
0 & 0 & 0 \\
 & & \\
 & 0 & \\
0 & 1 & 0 \\
\hline
0 & 0 & 0 \\
 & & \\
 & & 0 \\
0 & 0 & 1
\end{array}\right]
\end{array}
\begin{array}{c}
\\ \kappa_1 \\ \\ \\ \\ \kappa_2 \\ \\ \\ \\ \kappa_3 \\ \\ \\
\end{array}$$

in case $m = 3$, where $(\kappa_1, \kappa_2, \kappa_3) = \kappa(A, B)$ are the Kronecker indices of $\Sigma = (A, B)$. (The general case is evident from this example); every $(A, B) \in U(\kappa)$ is feedback equivalent to such a pair [30, 9]. The matrix $(sI - A\,;\, B)$ is now easily written down, and one observes that for all

$$s \neq 0, \infty, e_1 \equiv e_2 \equiv \ldots \equiv e_{\kappa_1} \equiv e_{n+1} \bmod \mathrm{Row}(sI - A\,;\, B)\,,$$

i.e. mod $\psi_\Sigma(s)$ and for $s = 0$, $e_2 \equiv \ldots \equiv e_{\kappa_1} \equiv e_{n+1} \equiv 0$ but $e_1 \neq 0$ and for $s \doteq \infty$, $e_1 \equiv \ldots \equiv e_{\kappa_1} \equiv 0$ and $e_{n+1} \neq 0$. It follows that the vectors

$$\varepsilon_1(\psi_\Sigma(s)), \ldots, \varepsilon_{\kappa_1}(\psi_\Sigma(s)), \varepsilon_{n+1}(\psi_\Sigma(s))$$

span a one-dimensional subspace of $\xi_m(\psi_\Sigma(s))$ for all s so that $E(\Sigma) \simeq \psi_\Sigma^! \xi_m$ contains a line bundle L_1 which admits at least $\kappa_1 + 1$ linearly independent holomorphic sections viz. the $\varepsilon_1, \ldots, \varepsilon_{\kappa_1}, \varepsilon_{n+1}$. Similar relations hold for

$$\varepsilon_{\kappa_1 + \ldots + \kappa_{i-1} + 1}, \ldots, \varepsilon_{\kappa_1 + \ldots + \kappa_i}, \varepsilon_{n+1}$$

for all $i = 1, \ldots, m$ giving us subbundles L_i, $i = 1, \ldots, m$ which admit at least $\kappa_i + 1$ linearly independent holomorphic sections. This exhausts the ε_i and because the $\varepsilon_1(x), \ldots, \varepsilon_{n+m}(x)$ span $\xi_m(x)$ for all $x \in \mathbf{G}_n(\mathbf{C}^{n+m})$ it follows that $E(\Sigma) = \oplus L_i$. As the pullback of the bundle ξ_m, $E(\Sigma)$ itself is a subbundle of an $(n+m)$-dimensional trivial bundle. Because $\mathbf{P}^1(\mathbf{C})$ is projective it follows (as before) that $E(\Sigma)$ has at most $n + m$ linearly independent holomorphic sections. But L_i has at least $\kappa_i + 1$ linearly independent sections, hence $\oplus L_i$ has at least $\Sigma(\kappa_i + 1) = n + m$ linearly independent sections which proves that L_i has precisely $\kappa_i + 1$ linearly independent sections and hence identifies L_i as the bundle $L(\kappa_i)$ described above in (8.5). We have reproved the theorem of Hermann and Martin [14].

8.12. *Theorem.* Keeping the notations introduced above in (8.10) and (8.5) we have $E(\Sigma) \simeq \bigoplus_{i=1}^{m} L(\kappa_i)$.

Still another proof of this theorem, using the Riemann-Roch theorem is found in Byrnes [33].

8.13. *The Correspondence B.* (cf. the diagram in section 5 above). The mapping $\Sigma \mapsto E(\Sigma)$ is obviously continuous. Thus the result $\overline{U(\kappa)} \supset U(\lambda) \leftrightarrow \kappa > \lambda$ can be deduced from Shatz's theorem (cf. 2.9). Inversely Shatz's theorem for positive bundles over $\mathbf{P}^1(\mathbf{C})$ can be deduced from the result on feedback orbits because every positive bundle arises as an $E(\Sigma)$. By tensoring with a suitable $L(r)$, r high enough, the result is then extended to arbitrary bundles over $\mathbf{P}^1(\mathbf{C})$.

9. Vectorbundles, systems and Schubert cells

9.1. *Partitions and Schubert-cells.* Let κ be a partition of n. To κ we associate the following increasing sequence of n numbers $\tau(\kappa)$.