Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 29 (1983)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: REPRESENTATIONS OF THE SYMMETRIC GROUP, THE SPECIALIZATION ORDER, SYSTEMS AND GRASSMANN MANIFOLDS

Autor: Hazewinkel, Michiel / Martin, Clyde F.

Kapitel: 10. Deformations of representation homomorphisms and subrepresentations

DOI: https://doi.org/10.5169/seals-52973

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 11.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

So we have the following

(1) $t < i + p(i)$ (by hypothesis)

(2) $n - i \geqslant n + m - t - l$ or equivalently $i \leqslant t + l - m$

(3) $\kappa_m + \ldots + \kappa_{m-l+1} + l \leqslant n + m - t$

(4) $\kappa_1 + \ldots + \kappa_{p(i)} < i \leqslant \kappa_1 + \ldots + \kappa_{p(i)+1}$.

Using (2) and (3) we have that

$$\kappa_m + \ldots + \kappa_{m-l+1} \leqslant n - i = \kappa_1 + \ldots + \kappa_m - i$$

so we have $i \leqslant \kappa_1 + \ldots + \kappa_{m-l}$ which implies $m - l \geqslant p(i) + 1$ thus

$$p(i) + i \leqslant m - l - 1 + i \leqslant (m-l-1) + (t+l-m) = t - 1$$

which contradicts (1). This proves the theorem.

9.7. *Vectorbundles and Schubert cells.* Because every positive vectorbundle over $\mathbf{P}^1(\mathbf{C})$ arises as the bundle $E(\Sigma)$ of some system Σ one has the obvious analogues of theorems 9.5 and 9.6 for positive bundles over $\mathbf{P}^1(\mathbf{C})$. Here the morphism ψ_Σ must, of course, be replaced by the classifying morphism (cf. section 3.2 above) of a positive vector bundle E, and $n + m$ and m are determined respectively as $\dim \Gamma(E, \mathbf{P}^1(\mathbf{C}))$ and $\dim E$.

10. Deformations of representation homomorphisms and subrepresentations

10.1 *On proving Inclusion Results for Representations.* Suppose we have given a continuous family of homomorphisms of finite dimensional representations over $\mathbf{C}$ of a finite group G

$$\pi_t : M \to V \tag{10.2}$$

Suppose that $Im\, \pi_t \simeq \rho$ for $t \neq 0$ (and small) and that $Im\, \pi_0 \simeq \rho_0$. Then the representation ρ_0 is a direct summand of the representation ρ. This is seen as follows. Because the category of finite dimensional complex representations of G is semisimple there is a homomorphism of representations $\phi_0 : Im\, \pi_0 \to M$ such that $\pi_0 \circ \phi_0 = id$. Then $\pi_t \circ \phi_0 : Im\, \phi_0 \to Im\, \pi_t$ is still injective for small t (by the continuity of π_t) which gives us ρ_0 as a subrepresentation and hence a direct summand of ρ.

It is almost equally easy to construct a surjective homomorphism $Im\, \pi_t \to Im\, \pi_0$.

10.3. *The Inverse Result.* Inversely if ρ_0 is a subrepresentation of ρ then there is a family of representations (10.3) such that $Im\ \pi_t \simeq \rho$ for $t \neq 0$ and $Im\ \pi_0 \simeq \rho_0$, and if ρ is generated (as a $\mathbf{C}[G]$-module) by one element one can take for M in (10.2) the regular representation. Indeed if ρ_0 is a subrepresentation of ρ then $\rho = \rho_0 \oplus \rho_1$. Let $\pi : M \to \rho = \rho_0 \oplus \rho_1$ be a surjective map of representations. Let π_0, π_1 be the two components of π. Let $s = (s_0, s_1)$ be a section of π. Then $\pi_0 s_0 = id, \pi_1 s_1 = id, \pi_0 s_1 = 0, \pi_1 s_0 = 0$ and it follows that $\pi(t)$ consisting of the components π_0 and $t\pi_1$ is still surjective. Hence $Im\ \pi(t) = \rho$ and $Im\ \pi(0) = \rho_0$.

11. A FAMILY OF REPRESENTATIONS OF S_{n+m} PARAMETRIZED BY $\mathbf{G}_n(\mathbf{C}^{n+m})$

11.1. *Construction of the Family.* Let M be the regular representation of S_{n+m}. That is M has a basis e_σ, $\sigma \in S_{n+m}$ and S_{n+m} acts on M by the formula $\tau(e_\sigma) = e_{\tau\sigma}$, for all $\tau \in S_{n+m}$. Now consider the universal bundle ξ_m over $\mathbf{G}(\mathbf{C}^{n+m})$ and the $n + m$ holomorphic section $\varepsilon_1, ..., \varepsilon_{n+m}$ defined by

$$\varepsilon_i(x) = e_i \bmod x \in \mathbf{C}^{n+m}/x\,,$$

where e_i is the i-th standard basis vector. Take the $(m+n)$-fold tensor product of ξ_m and define a family of homomorphisms parametrized by $\mathbf{G}_n(\mathbf{C}^{n+m})$ by

$$\pi_x : M \to \xi_m(x)^{\otimes(n+m)},\ e_{\sigma^{-1}} \mapsto \varepsilon_{\sigma(1)}(x) \otimes ... \otimes \varepsilon_{\sigma(n)}(x) \tag{11.2}$$

More precisely (11.2) defines a homomorphism of vectorbundles

$$\mathbf{G}_n(\mathbf{C}^{n+m}) \times M \to \xi_m^{\otimes(n+m)} \tag{11.3}$$

The group S_{n+m} acts on $\xi_m(x)^{\otimes(n+m)}$ by permuting the factors and it is a routine exercise to see that π_x is equivariant with respect to this action, i.e. that $\pi_x(\tau v) = \tau\pi_x(v)$ for all $v \in M, \tau \in S_{n+m}$. (Here the product $\tau\sigma \in S_{n+m}$ is interpreted as first the automorphism σ of $1, ..., n + m$ and then the automorphism τ.)

Thus $Im\ \pi_x = \pi(x)$ is a representation of S_{n+m} for all x giving us a family of representations parametrized by $\mathbf{G}_n(\mathbf{C}^{n+m})$. Fixing a point $x_0 \in \mathbf{G}_n(\mathbf{C}^{n+m})$ and choosing m independent sections of ξ_m in a neighbourhood U of x_0, this gives us families of homomorphisms of representations