Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	29 (1983)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	FREE GROUPS IN LINEAR GROUPS
Autor:	de la Harpe, Pierre
Kapitel:	4. Free subgroups of GL(2, R) and of GL(2, C)
DOI:	https://doi.org/10.5169/seals-52975

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

also elliptic, the foot of the perpendicular from the fixed point of g onto the invariant line of g would be fixed by g, and this cannot be. If g was at the same time elliptic with fixed point $a \in H^{n+1}$ and parabolic with fixed point $b \in S^n$, the line from a towards b would have two points at infinity b and b' both fixed by g, and this cannot be.

That any $g \in GM(n)_0$ belongs to one of the three classes follows for example from Brouwer's fixed point theorem. (See also 4.9.3 in [Th].)

Observe that an hyperbolic isometry $g \in GM(n)_0$ has a unique invariant line δ . Suppose indeed that there are two of them, say δ and δ' . If $\delta \cap \delta' \neq \phi$, the intersection point (which is unique) is fixed by g, and this cannot be. If $\delta \cap \delta' = \phi$ and if δ , δ' have no common point at infinity, there is a unique line perpendicular to both δ and δ' ; but this line intersects δ in a point fixed by g, and this cannot be. Assume finally that $\delta \cap \delta' = \phi$ and that δ and δ' have a common point at infinity; choose some number $\rho > 0$ and consider the set C_{ρ} of points in H^{n+1} at a distance of ρ from δ' ; the intersection $C_{\rho} \cap \delta$ is a point fixed by g, and again this cannot be. One may consequently also define an isometry $g \in GM(n)_0$ to be

elliptic if d(a, g(a)) = 0 for some $a \in H^{n+1}$,

parabolic if $\inf d(a, g(a)) = 0$, with the infimum over $a \in H^{n+1}$ not attained,

hyperbolic if $\inf d(a, g(a)) > 0$ (and the infimum is then attained exactly on the invariant line of g).

We shall need below the following dynamical description. An hyperbolic isometry $g \in GM(n)_0$ has on \mathbb{S}^n one attracting point P_a and one repulsing point P_r . This means that, for any neighborhood U of P_a in \mathbb{S}^n and for any compact subset K of $S^n - \{P_r\}$, one has $g^k(K) \subset U$ for k large enough. (And similarly with g^{-1} instead of g when exchanging P_a and P_r .) Consider now a parabolic isometry $g \in GM(n)_0$ with fixed point $P \in \mathbb{S}^n$. Let U be a neighborhood of P in \mathbb{S}^n and let K be compact in $S^n - \{P\}$; then $g^k(K) \subset U$ for any $k \in \mathbb{Z}$ with |k| large enough. (This is obvious when g is a translation in $\mathbb{R}^n \times \mathbb{R}^*_+$ by some vector in \mathbb{R}^n , and any parabolic isometry of H^{n+1} is conjugate to such a translation.)

4. Free subgroups of $GL(2, \mathbf{R})$ and of $GL(2, \mathbf{C})$

We show in this section that a subgroup of the proper Mæbius group $G = PGL(2, \mathbf{R})$ which is not almost solvable contains free groups; the same fact for $GL(2, \mathbf{R})$ follows straightforwardly. We discuss also the case of $GL(2, \mathbf{C})$.

PROPOSITION. Let $g, h \in G - \{1\}$ be without any common fixed point in $H^2 \cup S^1$. Then the group Γ generated by g and h contains free groups, up to two exceptions.

The first of these happens when $g^2 = h^2 = 1$. The second when one element is an involution, say $g^2 = 1$, when h is hyperbolic, and when g exchanges the two fixed points of h on S^1 . In these two cases, Γ is the infinite dihedral group, and is thus solvable.

Proof. We check below in each of the non exceptional cases that Γ contains a free group.

Case 1. One element, say g, is parabolic with fixed point $P \in S^1$.

Consider the parabolic $k = hgh^{-1}$, with fixed point $Q = h(P) \neq P$ in S¹. Let S_1 [respectively S_2] be a compact neighborhood of P [resp. Q] in S¹ with $S_1 \cap S_2 = \phi$. The end of section 3 shows that there exists a positive integer n_0 such that $g^n(S_2) \subset S_1$ and $k^n(S_1) \subset S_2$ for any $n \in \mathbb{Z}$ with $|n| \ge n_0$. It follows from Klein's criterium that g^{n_0} and k^{n_0} generate a free subgroup of G.

Case 2. Both g and h are hyperbolic.

Let S_1 [respectively S_2] be a compact neighborhood of the fixed points of g [resp. of h] in S^1 with $S_1 \cap S_2 = \phi$, and proceed as in case 1.

Case 3. One of the elements, say h, is hyperbolic with fixed points $P, Q \in S^1$ and g does not exchange them, say $R = g(Q) \notin \{P, Q\}$.

If $g(P) \in \{P, Q\}$ then h and ghg^{-1} are as in case 2. We may thus assume that g(P) = Q. If $g(R) \neq P$ then h and g^2hg^{-2} are again as in case 2. We may thus also assume g(R) = P. Consider then $h' = g^{-1}hg$, an hyperbolic with fixed points R and P, as well as $h'' = ghg^{-1}hgh^{-1}g^{-1}$, an hyperbolic with fixed points $Q = ghg^{-1}(Q)$ and $S = ghg^{-1}(P)$. One has $h(R) \neq Q$ and thus $S = gh(R) \neq g(Q) = R$; one has also $h(R) \neq R$ and $S \neq g(R) = P$. Consequently h' and h'' are as in case 2.

Case 4. Both g and h are elliptic with $g^2 \neq 1$.

Possibly after conjugation within G, one may assume that $g = r_{\alpha}$ is a rotation around the origin of the disc H^2 by some angle $\alpha \in [0, 2\pi[-\{\pi\}]$. Then $k = hgh^{-1} \neq g$, otherwise h would also fix the origin.

In the average, any point of S^1 is rotated by k of an angle α . More precisely, if $\tilde{k}: \mathbf{R} \to \mathbf{R}$ is the lifting of k to the universal covering of S^1 with $0 \leq \tilde{k}(0) < 1$, then $\lim_{n \to \infty} \frac{1}{n} (\tilde{k}^n(x) - x)$ exists for all $x \in \mathbf{R}$ and this limit is α . Moreover

 $\min_{x \in \mathbf{R}} \left(\tilde{k}(x) - x \right) \leq \alpha \leq \max_{x \in \mathbf{R}} \left(\tilde{k}(x) - x \right).$

(See any exposition of the rotation number, for example chapter 17 in [CL] or section 1 in [Ka].) It follows that there exists $P \in S^1$ with k(P) = g(P), so that $g^{-1}k$ has a fixed point in S^1 and one of the previous cases applies.

Exceptional cases. If $g^2 = h^2 = 1$, then gh generate an infinite cyclic subgroup of index 2 in Γ and Γ is isomorphic to the infinite dihedral group. If h is hyperbolic and if g exchanges its fixed points, then $ghg^{-1} = h^{-1}$ so that $g^2 = (gh)^2 = 1$ and Γ is as in the previous case.

The proof is now complete.

The proposition above is well known, and may essentially be found in any of the following papers: [LU1], [Md], [Ro] (see corollary 1). One should also mention Magnus' surveys [Ms1], [Ms2].

As two elements of G having a common fixed point in $H^2 \cup S^1$ generate a solvable subgroup, we have proved the 2-generators particular case of the following fact.

THEOREM 1. A subgroup Γ of $G = PGL(2, \mathbf{R})$ (or of $GL(2, \mathbf{R})$) which is not solvable contains free groups.

Proof. We assume that Γ does not contain free groups, and check that Γ is solvable. If Γ contains at least one parabolic isometry, this follows from case 1 of the proof above. If it contains at least one hyperbolic isometry, then all hyperbolics in Γ have a common fixed point (see case 2) and then either all elements in Γ have a common fixed point or Γ is dihedral (see case 3). Finally, if Γ is an elliptic group, it follows from case 4 that Γ is abelian.

This covers in particular the case of Fuchsian groups. The next theorem covers that of Kleinian groups.

THEOREM 2. Let Γ be a subgroup of $SL(2, \mathbb{C})$ which is not solvable. Assume moreover that Γ is not relatively compact (or equivalently that Γ is not conjugate to a subgroup of the maximal compact subgroup SU(2) of $SL(2, \mathbb{C})$). Then Γ contains free groups.

In particular, a discrete subgroup of $PGL(2, \mathbb{C})$ which is not almost solvable contains free groups.

Proof. The group Γ acts on \mathbb{C}^2 ; as Γ is not solvable, the representation is irreducible. Easy arguments à la Burnside show that Γ does not contain elliptic elements only; indeed, Γ does contain a hyperbolic element (see [CG], or corollary 1.8 in [B]). The first statement follows now as theorem 1.

The second follows from this: a discrete subgroup of $PGL(2, \mathbb{C})$ containing elliptic elements only is finite. Indeed, such a group is periodic. If Γ is a priori

known to be finitely generated, then Γ is finite by a theorem of Schur (§36 in [CR]) so that the hyperbolic subspace $F(\Gamma) = \{x \in H^3 \mid \Gamma x = \{x\}\}$ is non empty. In general, to any finitely generated subgroup Γ_t of Γ corresponds a non empty subspace $F_t \subset H^n$; it is easy to check that $F(\Gamma) = \bigcap F_t$ is non empty so that Γ lies in a compact subgroup of the Mœbius group; it follows again that Γ is finite.

Instead of the assumption of theorem 2, assume the following: there exists $g \in \Gamma$ with two distinct eigenvalues of same modulus, say $\mu_1 = \rho \exp(i\theta_1)$ and $\mu_2 = \rho \exp(i\theta_2)$ where ρ , θ_1 , $\theta_2 \in \mathbf{R}$ satisfy $\rho > 0$ and $\theta_1 \neq \theta_2 \pmod{2\pi}$, and there exists an automorphism α of \mathbf{C} with $|\alpha(\mu_1)| \neq |\alpha(\mu_2)|$. Then α induces an automorphism $\tilde{\alpha}$ of $GL(2, \mathbf{C})$ and the proof applies to $\tilde{\alpha}(\Gamma)$. But this procedure has its limits, because there exist complex numbers μ (such as $\frac{1}{5}(3+4i)$, see the remark below) such that $|\alpha(\mu)| = 1$ for any automorphism α of \mathbf{C} but which are not roots of 1; then the argument above fails ¹) for example for $g = \begin{pmatrix} \mu & 0 \\ 0 & \mu^{-1} \end{pmatrix}$.

Something is true however: let k be a finitely generated field of characteristic 0, let $\mu \in k - \{0\}$ and assume μ is not a root of 1. Then there exists a locally compact field k' endowed with an absolute value ω and there exists a homomorphism $\sigma: k \to k'$ such that $\omega(\sigma(\mu)) \neq 1$; this is lemma 4.1 of [T]. It follows that the argument above may be recuperated, but one has to consider other fields than subfields of **C**.

For self-consistency, let us end with the announced remark. For any automorphism α of C, one has clearly

$$\left| \alpha \left(\frac{3+4i}{5} \right) \right| = \left| \frac{3 \pm 4i}{5} \right| = 1;$$

we check now that $\frac{3+4i}{5}$ is not a root of one.

Let p, q be coprime integers and let $\mu = \exp\left(i2\pi \frac{p}{q}\right)$ be a root of 1. Then μ is an algebraic number of degree $\varphi(q)$, where φ is Euler's function. It follows that $\cos\left(2\pi \frac{p}{q}\right)$ is an algebraic number of degree $d \ge \frac{1}{2}\varphi(q)$: because if F is a polynomial of degree d in Z[X] with $F\left(\cos\left(2\pi \frac{p}{q}\right)\right) = 0$, then μ is a root of

¹) This shows that one point on page 50 of [D] is incorrect.

$$X^{d}F\left(\frac{1}{2}X+\frac{1}{2}X^{-1}\right)$$
, which is of degree 2d in $Z[X]$, so that $2d \ge \varphi(q)$. If $q \in \{1, 2, 3, 4, 6\}$, one checks easily that $\exp\left(i2\pi\frac{p}{q}\right) \ne \frac{3+4i}{5}$. If $q = 5$ or if $q \ge 7$, then $\varphi(q) > 2$ so that $\cos\left(2\pi\frac{p}{q}\right)$ is not rational. Thus the root of unity μ cannot be equal to $\frac{3+4i}{5}$.

5. Some other cases of Tits' theorem

Let *n* be an integer with $n \ge 2$.

Define a subgroup Γ of $GL(n, \mathbb{C})$ [respectively of $PGL(n, \mathbb{C})$] to be *irreducible* if any linear subspace of \mathbb{C}^n [resp. of $P_{\mathbb{C}}^{n-1}$] invariant by Γ is trivial, and *not almost reducible* if any subgroup of Γ of finite index is irreducible. When referring to the Zariski topology on $PGL(n, \mathbb{C})$, we use below the letter Z.

Reduction. Tits' theorem for complex linear groups is equivalent to the following statements (one for each $n \ge 2$):

Let Γ be a subgroup of $PGL(n, \mathbb{C})$ which is not almost solvable. Assume that (i) is not almost reducible;

(ii) the Z-closure G of Γ in $PGL(n, \mathbb{C})$ is Z-connected. Then Γ contains free groups.

That one may assume (i) without loss of generality is an easy exercise on reducibility, and one may assume (ii) because the Z-closure of any subgroup of $PGL(n, \mathbb{C})$ has finitely many Z-connected components. (The hypothesis of the reduced statement are redundant: (i) and (ii) imply by Lie's theorem that G is not solvable, so that Γ is not almost solvable!)

Now let $g \in PGL(n, \mathbb{C})$ and choose a representative $\tilde{g} \in GL(n, \mathbb{C})$ of g. Let us define g to be

elliptic if \tilde{g} is semi-simple with all eigenvalues of equal moduli,

parabolic if \tilde{g} is not semi-simple and has all its eigenvalues of equal moduli,

hyperbolic if \tilde{g} has at least two eigenvalues of distinct moduli.

These definitions are obviously independent on the choice of \tilde{g} . They generalize those of section 3 as follows from [Gr]. The meaning of "hyperbolic" fits with current use in dynamical systems theory (see e.g. definition 5.1 in [Sh]).

Let g be hyperbolic and let \tilde{g} be as above. Let $\tilde{A}(g)$ [respectively $\tilde{A}'(g)$] be the direct sum of the nilspaces of \tilde{g} corresponding to all eigenvalues of maximal modulus [resp. to all other eigenvalues] of \tilde{g} . Let A(g) [resp. A'(g)] be the canonical image of $\tilde{A}(g) - \{0\}$ [resp. $\tilde{A}'(g) - \{0\}$] in $\mathbf{P} = P_{\mathbf{C}}^{n-1}$. Then $A(g) \cap A'(g) = \emptyset$ and the smallest linear subspace of \mathbf{P} containing both A(g) and A'(g) is \mathbf{P} itself. Tits calls A(g) [resp. $A(g^{-1})$] the attracting space [resp. repulsing space] of g. We say that g is sharp if A(g) is a point and that g is very sharp if both A(g) and $A(g^{-1})$ are points. For each $k \in \{1, 2, ..., n-1\}$, the fundamental representation of GL(n, C) in $\wedge^k \mathbf{C}^n$ induces an injection

$$\lambda_k : PGL(n, \mathbf{C}) \to PGL(\binom{n}{k}, \mathbf{C});$$

as g is hyperbolic, $\lambda_k(g)$ is sharp for some k. We also say that two hyperbolic elements $g, h \in PGL(n, \mathbb{C})$ are in general position if

$$A(g) \cup A(g^{-1}) \subset \mathbf{P} - \{A'(h) \cup A'(h^{-1})\}$$
$$A(h) \cup A(h^{-1}) \subset \mathbf{P} - \{A'(g) \cup A'(g^{-1})\}.$$

Observe that any hyperbolic element of $PGL(2, \mathbb{C})$ is very sharp, and that two hyperbolic elements of $PGL(2, \mathbb{C})$ are in general position if and only if they do not have any common fixed point on \mathbb{S}^2 .

Recall that an element of $PGL(n, \mathbb{C})$ is semi-simple if its inverse image in $GL(n, \mathbb{C})$ contains diagonalisable matrices.

LEMMA 1. Let Γ be an irreducible subgroup of $PGL(n, \mathbb{C})$ having a Z-connected Z-closure. If Γ contains a sharp semi-simple element g, then Γ contains a very sharp element.

About the proof. Let $\tilde{g} \in GL(n, \mathbb{C})$ be some representative of g having an eigenvalue of "large" modulus and all other eigenvalues with moduli "near" 1. For suitable $h, u \in \Gamma$ and for $j \in N$ large enough, one may hope that $g^{-j}hg^{j}h^{-1}u$ has a representative in $GL(n, \mathbb{C})$ with one eigenvalue of very large modulus (look at $hg^{j}h^{-1}u$), one eigenvalue of very small modulus (look at g^{-j}), and other eigenvalues of moduli "near" 1. Section 3 of [T] shows that this hope is realistic. (See also below, after the theorem.)

LEMMA 2. Let Γ be an irreducible subgroup of PGL(n, C) having a Zconnected Z-closure. If Γ contains a very sharp element, then Γ contains two very sharp elements in general position.

Proof. Let P_1 , P_2 be two linear subspaces of **P** with $P_1 \neq \emptyset$ and $P_2 \neq \mathbf{P}$. Then $\{x \in G \mid x(P_1) \notin P_2\}$ is obviously a Z-open subset of G. It is not empty: Choose indeed $p \in P_1$; then the subspace of **P** spanned by the orbit Gp is stable under G and must therefore coincide with **P**; hence there exists $x \in G$ with $x(p) \notin P_2$ and, a fortiori, $x(P_1) \notin P_2$.

Let g be a very sharp element in Γ . It follows from above that

$$X = \left\{ x \in G \mid A(g) \text{ and } A(g^{-1}) \text{ are not contained in any of } xA'(g), \\ xA'(g^{-1}), x^{-1}A'(g), x^{-1}A'(g^{-1}) \right\}$$

is a non empty Z-open subset of G. Let $y \in X \cap \Gamma$. Then g and ygy^{-1} are both very sharp and are in general position.

For the next lemma, we choose as above k with $1 \le k \le n-1$ and we consider the k^{th} fundamental representation $\lambda_k : SL(n, \mathbb{C}) \to SL(\binom{n}{k}, \mathbb{C})$ of $SL(n, \mathbb{C})$.

LEMMA. Let Γ be a group and let $\rho: \Gamma \to SL(n, \mathbb{C})$ be an irreducible representation. Then the Z-closure G of $\rho(\Gamma)$ in $SL(n, \mathbb{C})$ is semi-simple and the representation $\sigma = \lambda_k \rho: \Gamma \to SL(\binom{n}{k}, \mathbb{C})$ is completely reducible.

Proof. We show first that G is semi-simple. Consider the solvable radical R of G. By Lie's theorem, there exists an eigenvector for R, namely there exist $v \in \mathbb{C}^n - \{0\}$ and $\alpha \in \text{Hom}(R, \mathbb{C}^*)$ with $r(v) = \alpha(r)v$ for all $r \in R$. As R is normal in G, any vector g(v) ($g \in G$) is also an eigenvector for R. By irreductibility, any vector in \mathbb{C}^n is also an eigenvector, so that R is made up of dilations. But R is connected and is in $SL(n, \mathbb{C})$, so that R = 1.

Now $\lambda_k: G \to SL(\binom{n}{k}, \mathbb{C})$ is completely reducible; denote by $\lambda_{k,j}: G \to SL(W_j)$ the components of a decomposition $\lambda_k = \bigoplus_{j \in J} \lambda_{k,j}$ and define σ_j = $\lambda_{k,j}\rho$ ($j\in J$). One has clearly $\sigma = \bigoplus_{j \in J} \sigma_j$, and each $\sigma_j: \Gamma \to SL(W_j)$ is irreducible (this because $\lambda_{k,j}$ is irreducible and by Schur's lemma).

THEOREM. Let Γ be a subgroup of $PGL(n, \mathbb{C})$ and assume (i) Γ is neither almost solvable nor almost reducible,

(ii) Γ contains a semi-simple hyperbolic element.

Then Γ contains free groups.

Proof. As one may consider instead of Γ a subgroup of finite index, there is no loss of generality if we assume that the Z-closure of Γ is Z-connected. We denote by $\tilde{\Gamma}$ the inverse image of Γ in $SL(n, \mathbb{C})$. By (ii), there exists $k \in \{1, ..., n-1\}$ and a semi-simple element $\tilde{\gamma} \in \tilde{\Gamma}$ having eigenvalues $\mu_1, ..., \mu_n$ with $|\mu_1| = ...$ $= |\mu_k| > |\mu_j|$ for j = k + 1, ..., n. Let $N = \binom{n}{k}$, and denote by λ_k both the fundamental representation $GL(n, \mathbb{C}) \rightarrow GL(N, \mathbb{C})$ and the induced homomorphism $PGL(n, \mathbb{C}) \to PGL(N, \mathbb{C})$. Then $\lambda_k(\tilde{\gamma})$ has eigenvalues $v_1, ..., v_N$ with $|v_1| > |v_j|$ for j = 2, ..., N. By lemma 3, there exists a $\lambda_k(\tilde{\Gamma})$ -irreducible subspace W_0 of \mathbb{C}^N , associated to a representation $\sigma_0 \colon \tilde{\Gamma} \to GL(W_0)$, such that v_1 is an eigenvalue of $\sigma_0(\tilde{\gamma})$. As the Z-closure \tilde{G} of $\tilde{\Gamma}$ in $SL(n, \mathbb{C})$ is semi-simple, the group \tilde{G} is perfect and $\sigma_0(\tilde{\Gamma})$ lies in $SL(W_0)$. As $|v_1| > 1$, one has $\dim_{\mathbb{C}} W_0 \ge 2$.

Thus one may assume from the start that Γ contains a sharp semi-simple element, and indeed by lemmas 1 and 2 two very sharp elements in general position. The conclusion follows as in case 2 of the proof of the proposition in section 4.

Now lemma 1 remains true without the hypothesis "semi-simple". This has been announced by Y. Guivarch', who uses ideas of H. Fürstenberg to show the following: given an appropriate subset S of Γ containing a sharp element, then almost any "long" word in the letters of S is very sharp. Using this, one may replace (ii) in the theorem above by the following a priori weaker hypothesis

(ii') Γ is not relatively compact.

Then, one first checks as for theorem 2 of section 4 that Γ contains hyperbolic elements; one concludes as in the previous proof, with Guivarch's version of lemma 1.

For subgroups of PU(n), one may repeat the discussion at the end of section 4.

REFERENCES

- [A] AHLFORS, L. V. Möbius transformations in several dimensions. School of Mathematics, University of Minnesota, 1981.
- [Ba] BASS, H. Groups of integral representation type. Pacific J. Math. 86 (1980), 15-51.
- [BL] BASS, H. and A. LUBOTZKY. Automorphisms of groups and of schemes of finite type. *Preprint*.
- [B] BOURBAKI, N. Eléments d'histoire des mathématiques. Hermann 1969.
- [CL] CODDINGTON, E. A. and N. LEVINSON. Theory of ordinary differential equations. McGraw Hill, 1955.
- [CG] CONZE, J. P. and Y. GUIVARCH'. Remarques sur la distalité dans les espaces vectoriels. C. R. Acad. Sc. Paris, Sér. A, 278 (1974), 1083-1086.
- [CR] CURTIS, C. and I. REINER. Representation theory of finite groups and associative algebras. Interscience, 1962.
- [DE] DUBINS, L. E. and M. EMERY. Le paradoxe de Hausdorff-Banach-Tarski. Gazette des Mathématiciens 12 (1979), 71-76.
- [D] DIXON, J. D. Free subgroups of linear groups. Lecture Notes in Math. 319 (Springer, 1973), 45-56.
- [E] EPSTEIN, D. B. A. Almost all subgroups of a Lie group are free. J. of Algebra 19 (1971), 261-262.
- [FK] FRICKE, R. and F. KLEIN. Vorlesungen über die Theorie der automorphen Functionen, vol. I. Teubner, 1897.

4