

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 29 (1983)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: FREE GROUPS IN LINEAR GROUPS
Autor: de la Harpe, Pierre

Bibliographie

DOI: <https://doi.org/10.5169/seals-52975>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

homomorphism $PGL(n, \mathbf{C}) \rightarrow PGL(N, \mathbf{C})$. Then $\lambda_k(\tilde{\gamma})$ has eigenvalues v_1, \dots, v_N with $|v_1| > |v_j|$ for $j = 2, \dots, N$. By lemma 3, there exists a $\lambda_k(\tilde{\Gamma})$ -irreducible subspace W_0 of \mathbf{C}^N , associated to a representation $\sigma_0: \tilde{\Gamma} \rightarrow GL(W_0)$, such that v_1 is an eigenvalue of $\sigma_0(\tilde{\gamma})$. As the Z -closure \tilde{G} of $\tilde{\Gamma}$ in $SL(n, \mathbf{C})$ is semi-simple, the group \tilde{G} is perfect and $\sigma_0(\tilde{\Gamma})$ lies in $SL(W_0)$. As $|v_1| > 1$, one has $\dim_{\mathbf{C}} W_0 \geq 2$.

Thus one may assume from the start that Γ contains a sharp semi-simple element, and indeed by lemmas 1 and 2 two very sharp elements in general position. The conclusion follows as in case 2 of the proof of the proposition in section 4. \square

Now lemma 1 remains true without the hypothesis "semi-simple". This has been announced by Y. Guivarch', who uses ideas of H. Fürstenberg to show the following: given an appropriate subset S of Γ containing a sharp element, then almost any "long" word in the letters of S is very sharp. Using this, one may replace (ii) in the theorem above by the following a priori weaker hypothesis

(ii') Γ is not relatively compact.

Then, one first checks as for theorem 2 of section 4 that Γ contains hyperbolic elements; one concludes as in the previous proof, with Guivarch's version of lemma 1.

For subgroups of $PU(n)$, one may repeat the discussion at the end of section 4.

REFERENCES

- [A] AHLFORS, L. V. *Möbius transformations in several dimensions*. School of Mathematics, University of Minnesota, 1981.
- [Ba] BASS, H. Groups of integral representation type. *Pacific J. Math.* 86 (1980), 15-51.
- [BL] BASS, H. and A. LUBOTZKY. Automorphisms of groups and of schemes of finite type. *Preprint*.
- [B] BOURBAKI, N. *Eléments d'histoire des mathématiques*. Hermann 1969.
- [CL] CODDINGTON, E. A. and N. LEVINSON. *Theory of ordinary differential equations*. McGraw Hill, 1955.
- [CG] CONZE, J. P. and Y. GUIVARCH'. Remarques sur la distalité dans les espaces vectoriels. *C. R. Acad. Sc. Paris, Sér. A*, 278 (1974), 1083-1086.
- [CR] CURTIS, C. and I. REINER. *Representation theory of finite groups and associative algebras*. Interscience, 1962.
- [DE] DUBINS, L. E. and M. EMERY. Le paradoxe de Hausdorff-Banach-Tarski. *Gazette des Mathématiciens* 12 (1979), 71-76.
- [D] DIXON, J. D. Free subgroups of linear groups. *Lecture Notes in Math.* 319 (Springer, 1973), 45-56.
- [E] EPSTEIN, D. B. A. Almost all subgroups of a Lie group are free. *J. of Algebra* 19 (1971), 261-262.
- [FK] FRICKE, R. and F. KLEIN. *Vorlesungen über die Theorie der automorphen Functionen*, vol. I. Teubner, 1897.

- [Gr] GREENBERG, L. Discrete subgroups of the Lorentz group. *Math. Scand.* 10 (1962), 85-107.
- [G] GROMOV, M. Groups of polynomial growth and expanding maps. *Publ. Math. I.H.E.S.* 53 (1981), 53-78.
- [H] HAUSDORFF, F. *Grundzüge der Mengenlehre*. Veit, 1914.
- [Hm] HAUSMANN, J. C. Sur l'usage de critères pour reconnaître un groupe libre, un produit amalgamé ou une HNN -extension. *L'Enseignement math.* 27 (1981), 221-242.
- [Ig] IGNATOV, Ju. A. Free and nonfree subgroups of $PSL_2(\mathbb{C})$ generated by two parabolic elements. *Math. USSR Sbornik* 35 (1) (1979), 49-55.
- [Ka] KATZNELSON, Y. Sigma-finite invariant measure for smooth mappings of the circle. *J. d'analyse math.* 31 (1977), 1-18.
- [Ku] KURANISHI, M. Two elements generations on semi-simple Lie groups. *Kōda Math. Sem. rep.* 5-6 (1949), 9-10.
- [L] LEUTBECHER, A. Über die Heckeschen Gruppen $G(\lambda)$. *Abh. Math. Sem. Un. Hamburg* 31 (1967), 199-205.
- [L1] LICHTMAN, A. On subgroups of the multiplicative group of skew fields. *Proc. Amer. Math. Soc.* 63 (1971), 15-16.
- [L2] ——— On normal subgroups of multiplicative group of skew fields generated by a polycyclic-by-finite group. *J. of Algebra* 78 (1982), 548-577.
- LU1] LYNDON, R. C. and J. L. ULLMAN. Pairs of real 2-by-2 matrices that generate free products. *Michig. Math. J.* 15 (1968), 161-166.
- [LU2] ——— Groups generated by two parabolic linear fractional transformations. *Can. J. Math.* 21 (1969), 1388-1403.
- [LS] LYNDON, R. C. and P. E. SCHUPP. *Combinatorial group theory*. Springer, 1977.
- [Ms1] MAGNUS, W. Two generator subgroups of $PSL(2, \mathbb{C})$. *Nachr. Akad. Wiss. Göttingen, II Math. Phys. Kl.* 7 (1975), 81-94.
- [Ms2] ——— The uses of 2 by 2 matrices in combinatorial group theory. A survey. *Res. der Math.* 4 (1981), 171-192.
- [Md] MAJEED, A. Freeness of the group $\langle a^n, b^n \rangle$ for some integer n , $a, b \in SL(2, \mathbb{C})$. *J. Math. Soc. Japan* 29 (1977), 29-33.
- [M] MAL'CEV, A. I. On the faithful representations of infinite groups by matrices. *Amer. Math. Soc. Transl. (2)* 45 (1965), 1-18.
- [Mt] MASKIT, B. On Poincaré's theorem for fundamental polygons. *Adv. in Math.* 7 (1971), 219-230.
- [Mi] MATELSKI, J. P. The classification of discrete 2-generator subgroups of $PSL(2, \mathbb{R})$. *Israel J. of Math.* 42 (1982), 309-317.
- [vN] VON NEUMANN, J. Zur allgemeinen Theorie des Masses. *Fund. Math.* 13 (1929), 143-173.
- [N] NEWMAN, M. Pairs of matrices generating discrete free groups and free products. *Michig. Math. J.* 15 (1968), 155-160.
- [Ro] ROSENBERGER, G. On discrete free subgroups of linear groups. *J. London Math. Soc.* 17 (1978), 79-85.
- [S] SERRE, J. P. Arbres, amalgames, SL_2 . *Astérisque* 46 (Soc. math. France, 1977).
- [Sh] SHUB, M. Stabilité globale des systèmes dynamiques. *Astérisque* 56 (Soc. math. France, 1978).
- [Si] SIEGEL, C. L. *Topics in complex function theory*, volume II. Wiley-Interscience 1971.
- [Th] THURSTON, W. P. *The geometry and topology of 3-manifolds*. Notes from lectures at Princeton, 1978-1979, chapters 1-9.
- [T] TITS, J. Free subgroups in linear groups. *J. of Algebra* 20 (1972), 250-270.

- [Wa] WANG, S. P. A note on free subgroups in linear groups. *J. of Algebra* 71 (1981), 232-234.
- [Wh] WEHRFRITZ, B. A. F. 2-generator conditions in linear groups. *Archiv. Math.* 22 (1971), 237-240.
- [Wi] WEIL, A. *L'intégration dans les groupes topologiques et ses applications*. Hermann, 1940.

ADDED IN PROOF: See moreover the report by Yu. I. Merzlyakov "Linear groups", *J. Soviet Math.* 14 (1980), 887-921.

(Reçu le 18 janvier 1983)

Pierre de la Harpe

Section de mathématiques

C.P. 240

CH-1211 Genève 24