

# DIVISION ALGEBRAS AND THE HAUSDORFF-BANACH-TARSKI PARADOX

Autor(en): **Deligne, Pierre / Sullivan, Dennis**

Objektyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **29 (1983)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **26.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-52976>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

## DIVISION ALGEBRAS AND THE HAUSDORFF-BANACH-TARSKI PARADOX

by Pierre DELIGNE and Dennis SULLIVAN


In this note we observe that a question raised by Dekker (1956) about rotations inspired by the Hausdorff-Banach-Tarski paradox can be answered using algebraic number theory. For motivation, we recall a form of the paradox.

Partition the free group in two generators  $F$  into the five sets  $E, X, \bar{X}, Y, \bar{Y}$  consisting respectively of the identity and of the elements which, when written in reduced form, begin with  $x, x^{-1}, y$  or  $y^{-1}$ . If  $F$  acts freely on a sphere  $S$ , by rigid rotations, and if, using the axiom of choice, we choose a transversal  $T$ , i.e. a set with exactly one point in each orbit, the five subsets  $T, XT, \bar{X}T, YT, \bar{Y}T$  form a partition of  $S$ . For economy of notation, we will write again  $E, X, \bar{X}, Y, \bar{Y}$  for  $T, XT, \bar{X}T, YT, \bar{Y}T$ . The rotation by  $x$  moves  $\bar{X}$  onto

$$S - X = E \cup \bar{X} \cup Y \cup \bar{Y}.$$

Similarly,  $y$  moves  $\bar{Y}$  onto

$$S - Y = E \cup X \cup \bar{X} \cup \bar{Y}.$$



Thus we can reassemble from these 11 (actually 5) pieces 2 congruent spheres plus one congruent copy of the set  $E$ . This is a form of the Hausdorff-Banach-Tarski paradox which comes quickly from a free action of a free group

on two generators (see Appendix A for the more precise form). Thus we have the question of Dekker (communicated by Jan Mycielski): do such actions really exist? The sphere must be odd-dimensional for topological reasons: a fixed point free map  $f : S^d \rightarrow S^d$  must have vanishing Lefschetz number

$$L(f) = \sum (-1)^i \text{Trace}(f ; H_i(S^d)) = 1 + (-1)^d \text{deg}(f).$$

For a free action of a free group, the square of a non-trivial element will act by a map of degree  $+1$  and this forces  $d$  odd. The dimension must be  $> 1$ . These are the only conditions.

**THEOREM.** *For  $n \geq 2$ , there is a free non-abelian group of rigid rotations acting freely on the odd dimensional sphere  $S^{2n-1}$ .*

*Remark.* The corresponding orthogonal matrices can be chosen to have algebraic entries, and the group of matrices corresponds to a subgroup of the non-zero elements in a division algebra over a number field.

*Remark.* The theorem was proved by Dekker for  $n$  even [D].

*Remark.* Let  $u : O(p) \times O(q) \rightarrow O(p+q)$  be the natural embedding:

$$u(A, B) = \text{matrix} \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}.$$

If  $A \in O(p)$  and  $B \in O(q)$  don't have any nonzero fixed vector, neither has  $u(A, B)$ . If  $\sigma_1 : F \rightarrow O(p)$  and  $\sigma_2 : F \rightarrow O(q)$  define free actions on  $S^{p-1}$  and  $S^{q-1}$ ,  $u(\sigma_1, \sigma_2)$  hence defines a free action of  $F$  on  $S^{p+q-1}$ . Using this remark, one could reduce the theorem to the two particular cases  $n = 2$  and  $n = 3$ .

*Proof.* Let  $k \subset \mathbf{R}$  be a real algebraic number field, and  $k' \subset \mathbf{C}$  be a quadratic extension of  $k$ . We assume that  $k' \not\subset \mathbf{R}$ , i.e. that  $k' \otimes_k \mathbf{R} = \mathbf{C}$ . Let  $D$  be a division algebra of dimension  $n^2$  over its center  $k'$ , equipped with an anti-involution  $*$  inducing on  $k'$  the complex conjugation. The  $\mathbf{R}$ -algebra  $D \otimes_k \mathbf{R}$  is a simple algebra over its center  $k' \otimes_k \mathbf{R} = \mathbf{C}$ , hence isomorphic to  $M(n, \mathbf{C})$ . We assume that, for a suitable isomorphism between  $D \otimes_k \mathbf{R}$  and  $M(n, \mathbf{C})$ ,  $*$  becomes transpose conjugate.

In term of an isomorphism as above, the elements  $u$  of  $D$  satisfying  $uu^* = 1$  become unitary matrices. They operate on the unit sphere in  $\mathbf{C}^n$ . Furthermore, if  $u \neq 1$ ,  $u - 1$  is invertible in  $D$  so that the corresponding matrix does not have 1 as an eigenvalue. It hence acts without fixed point on the sphere.

The group  $\Gamma := \{u \in D \mid uu^* = 1\}$  is the group of  $k$ -rational points of a  $k$ -form of the real algebraic group  $U(n)$ . For  $n > 1$ , the perfect subgroup  $SU(n)$

of  $U(n)$  is not trivial, and  $U(n)$  is not solvable. The group  $\Gamma$  is dense in  $U(n)$ : skew adjoints elements of  $D$  are dense in the skew adjoint matrices in  $M(n, \mathbf{C})$ , and the Cayley transform  $t \mapsto \frac{t - 1}{t + 1}$  is an homeomorphism from the space of skew-adjoint matrices in  $M(n, \mathbf{C})$  to an open dense subset of  $U(n)$ , carrying skew adjoint elements of  $D$  into  $\Gamma$ . From this density, it results that, if  $n > 1$ , the linear group  $\Gamma$  is not solvable. By [Tits], it contains a non abelian free subgroup.

It remains to construct pairs  $(D, *)$ . A division algebra  $D$  with center  $k'$  admits an anti-involution  $*$  inducing on  $k'$  the non trivial element  $\text{Gal}(k'/k)$ , if and only if its class  $\text{cl}(D)$  in the Brauer group  $\text{Br}(k')$  of  $k'$  has a trivial image by the norm map  $N_{k'/k} : \text{Br}(k') \rightarrow \text{Br}(k)$ —see Appendix B. Class field theory provides an explicit computation of  $\text{Br}(k)$ , and of  $N_{k'/k}$ , and tells which elements of  $\text{Br}(k')$  come from division algebras. From the explicit description it provides, existence of such  $D$  follows. A direct construction is given in Appendix C. When we choose an isomorphism of  $D \otimes_k \mathbf{R}$  with  $M(n, \mathbf{C})$ , the involution  $*$  becomes adjunction with respect to some hermitian form  $\phi$  on  $\mathbf{C}^n$ , not necessarily positive definite:  $\phi(ax, y) = \phi(x, a^*y)$ . If  $h$  is self adjoint in  $D$ ,  $\text{int}(h^{-1}) \circ *$  is adjunction, with respect to the form  $\phi_h(x, y) = \phi(hx, y)$ . For suitable  $h$ ,  $\phi_h$  is positive definite and  $(D, \text{int}(h^{-1}) \circ *)$  is of the type sought.

### APPENDIX A

Consider  $\phi : S' \cup S'' \rightarrow S - E$  as in the introduction, with  $S'$  and  $S''$  two copies of the sphere  $S$ , and  $\psi : S \rightarrow S'$  the obvious bijection. Consider as in the Schröder-Bernstein theorem the set  $S_e$  of points  $p$  in  $S$  with an even number of ancestors, namely for which there exists an integer  $n \geq 0$  with  $p \in \text{Im}(\phi \circ \psi)^n$  and  $p \notin \text{Im}(\psi \circ (\phi \circ \psi)^n)$ . Consider also the set  $S_0$  of those  $p$  in  $S$  for which there exists  $n \geq 0$  with  $p \in \text{Im}(\psi \circ (\phi \circ \psi)^n)$  and  $p \notin \text{Im}(\phi \circ \psi)^{n+1}$ , and finally the set  $S_\infty$  of those  $p$  such that  $p \in \text{Im}(\phi \circ \psi)^n$  for any  $n \geq 0$ . Consider similarly

$$S' \cup S'' = (S' \cup S'')_e \cup (S' \cup S'')_0 \cup (S' \cup S'')_\infty.$$

Then  $\psi$  induces a bijection from  $S_e \cup S_\infty$  onto  $(S' \cup S'')_0 \cup (S' \cup S'')_\infty$  and  $\phi^{-1}$  from  $S_0$  onto  $(S' \cup S'')_e$ . Combining these two we have a bijection  $\chi : S \rightarrow S' \cup S''$  and a partition of  $S$  into finitely many pieces, the restriction of  $\chi$  to each of these being a rotation.

## APPENDIX B

Let  $K$  be a separable quadratic extension of a field  $k$ . We denote  $x \mapsto \bar{x}$  the non trivial element  $\text{Gal}(K/k)$ . Let  $D$  be a simple algebra with dimension  $n^2$  over its center  $K$ . We will check the criterion of the text, for the existence of an involution of the second kind on  $D$ , i.e. of an anti-involution  $*$  of  $D$ , inducing  $x \mapsto \bar{x}$  on  $K$ . The criterion is that  $N_{K/k} \text{cl}(D) = 0$  in  $\text{Br}(k)$ .

Let us localize, for the étale topology, over  $\text{Spec}(k)$ . This means making large enough étale extensions of scalars  $\otimes_k k'$ , and keeping track of the functoriality in  $k'$ . The field  $K$  becomes the separable quadratic extension  $K' = K \otimes_k k'$  of  $k'$ . The algebra  $D$  becomes  $D' = D \otimes_k k'$ , and is of the form  $D' = \text{End}_{K'}(V')$ , for  $V'$  a free module  $K'$ . The module  $V'$  is not determined uniquely by  $D'$ , only up to homotheties (the corresponding projective space is uniquely determined).

For any  $K$ -module  $M$ , let  $M^-$  be the module deduced from  $M$  by the extension of scalars  $\bar{\phantom{x}} : K \rightarrow K$ , i.e. the module, unique up to unique isomorphism, provided with an anti-linear isomorphism  $x \mapsto \bar{x} : M \xrightarrow{\sim} M^-$ . Similarly for  $K'$ -modules. If  $D' = \text{End}(V')$ , then  $D'^- = \text{End}(V'^-)$ , and

$$(D \otimes_k D^-)' = \text{End}(V' \otimes V'^-).$$

Let  $W'$  be the fixed subspace of the anti-linear automorphism of  $V' \otimes V'^-$  defined by  $v \otimes \bar{w} \mapsto w \otimes \bar{v}$ . It is the space of Hermitian forms on the dual of  $V'$ . One has  $W' \otimes_{k'} K' = V' \otimes V'^-$ . If  $D_1 \subset D \otimes_k D^-$  is the fixed subspace of the anti-linear automorphism of  $D \otimes_k D^-$  defined by  $x \otimes \bar{y} \mapsto y \otimes \bar{x}$ , then  $D'_1$  is the  $k'$ -form of the  $K'$ -algebra  $(D \otimes_k D^-)' = \text{End}(V' \otimes V'^-)$  deduced from the  $k'$ -form  $W'$  of the  $K'$ -module  $V' \otimes V'^- : D'_1 = \text{End}_{k'}(W')$ .

Involutions of the second kind on  $D'$  correspond one to one to non degenerate Hermitian forms on  $V'$ , taken up to a factor (in  $k'^*$ ). Those, in turn, by the “dual form” construction, correspond to “non degenerate” elements of  $W'$ . Again, one has to take them up to a factor. The projective space  $\mathbf{P}(W')$  over  $k'$  is determined up to unique isomorphism by  $D'$ . It is hence (this is the point of localisation) defined over  $k : \mathbf{P}(W') = P \otimes_k k'$ , functorially in  $k'$ . The  $k$ -points of  $P$  (rather, the non degenerate points) parametrize the involutions of the second kind on  $D$ .

The functorial isomorphism  $D'_1 = \text{End}_{k'}(W')$  shows that  $P$  is the form of projective space (Severi-Brauer variety) attached to  $D_1$ . It has a rational point, and is then the ordinary projective space, if and only if  $D_1$  is a matrix algebra.

This shows that  $D$  has involutions of the second kind if and only if the class of  $D_1$  in  $\text{Br}(k)$  is trivial. This class is the required norm  $N_{K/k}(\text{cl}(D))$ . In the localization spirit, this can be deduced from the fact that the homothety by  $\lambda \in K'^*$  of  $V'$  induces on  $W'$  the homothety by  $N_{K'/k'}(\lambda) \in k'^*$ .

### APPENDIX C

For  $n \geq 3$ , examples can be obtained as follows: take  $k' = \mathbf{Q}[\zeta]$ , with  $\zeta = \exp(2\pi i/n)$ , and  $k = k' \cap \mathbf{R}$ . Fix  $a, b \in k'^*$  and let  $D$  be the  $k'$ -algebra generated by  $X, Y$ , subject to

$$\begin{aligned} X^n &= a, & Y^n &= b \\ XY &= \zeta YX. \end{aligned}$$

It admits the anti-involution  $*$ , inducing complex conjugation on  $k'$ , defined by  $\zeta^* = \zeta^{-1}$ ,  $X^* = X$ ,  $Y^* = Y$ . The algebra  $D$  is of the type we require, provided it is a division algebra. This happens already with  $a, b \in \mathbf{Z}$ : take for  $a$  a prime congruent to 1 mod  $n$ , and for  $b$  an integer whose residue mod  $a$  has in the cyclic group of order  $n$   $(\mathbf{Z}/(a))^*/(\mathbf{Z}/(a))^{*n}$  an image of exact order  $n$ . For instance  $n = 3$ ,  $a = 7$ ,  $b = 2$ . For  $n = 2$ , one proceeds similarly with  $k' = \mathbf{Q}[i]$ ,  $\zeta = -1$ ,  $a$  congruent to 1 mod 4 and  $b$  not a square mod  $a$ . For instance,  $a = 5$ , and  $b = 2$ . In each case, the assumption on  $a$  ensures that  $k'$  embed in the  $a$ -adic completion  $\mathbf{Q}_a$  of  $\mathbf{Q}$ , and the fact that  $D$  is a division algebra can be seen locally at  $a$ :  $D \otimes_{k'} \mathbf{Q}_a$  is a division algebra with center  $\mathbf{Q}_a$ .

## BIBLIOGRAPHY

- [A] ALBERT, A. *Structure of algebras*. A.M.S. Colloquium Publ. 24 (1939).  
[Sch] SCHARLAU, W. Zur Existenz von Involutionen auf einfachen Algebren. *Math. Zeitschr.* 145 (1975), 29-32.  
[Myc] MYCIELSKI, Jan. Can one solve equations in groups? *Amer. Math. Monthly* 84 (1977), 723-726.  
[Tits] TITS, J. Free subgroups in linear groups. *Journal of Algebra* 20 (1972), 250-270.  
[D] DEKKER, Th. J. Decomposition of sets and spaces I, II. *Indig. Math.* 18 (1956), 581-595, and 19 (1957), 104-107.

(Reçu le 8 mars 1982)

Pierre Deligne  
Dennis Sullivan

Institut des Hautes Etudes Scientifiques  
35, route de Chartres  
91440 Bures-sur-Yvette  
France