Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 29 (1983)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: DIVISION ALGEBRAS AND THE HAUSDORFF-BANACH-TARSKI PARADOX

Autor: Deligne, Pierre / Sullivan, Dennis

Anhang: Appendix A

DOI: https://doi.org/10.5169/seals-52976

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 11.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

of $U(n)$ is not trivial, and $U(n)$ is not solvable. The group Γ is dense in $U(n)$: skew adjoints elements of D are dense in the skew adjoint matrices in $M(n, \mathbf{C})$, and the Cayley transform $t \mapsto \dfrac{t-1}{t+1}$ is an homeomorphism from the space of skew-adjoint matrices in $M(n, \mathbf{C})$ to an open dense subset of $U(n)$, carrying skew adjoint elements of D into Γ. From this density, it results that, if $n > 1$, the linear group Γ is not solvable. By [Tits], it contains a non abelian free subgroup.

It remains to construct pairs $(D, *)$. A division algebra D with center k' admits an anti-involution $*$ inducing on k' the non trivial element $\mathrm{Gal}(k'/k)$, if and only if its class $\mathrm{cl}(D)$ in the Brauer group $\mathrm{Br}(k')$ of k' has a trivial image by the norm map $N_{k'/k} : \mathrm{Br}(k') \to \mathrm{Br}(k)$—see Appendix B. Class field theory provides an explicit computation of $\mathrm{Br}(k)$, and of $N_{k'/k}$, and tells which elements of $\mathrm{Br}(k')$ come from division algebras. From the explicit description it provides, existence of such D follows. A direct construction is given in Appendix C. When we choose an isomorphism of $D \otimes_k \mathbf{R}$ with $M(n, \mathbf{C})$, the involution $*$ becomes adjunction with respect to some hermitian form ϕ on $\mathbf{C}^n$, not necessarily positive definite: $\phi(ax, y) = \phi(x, a^*y)$. If h is self adjoint in D, $\mathrm{int}(h^{-1}) \circ *$ is adjunction, with respect to the form $\phi_h(x, y) = \phi(hx, y)$. For suitable h, ϕ_h is positive definite and $(D, \mathrm{int}(h^{-1}) \circ *)$ is of the type sought.

Appendix A

Consider $\phi : S' \cup S'' \to S - E$ as in the introduction, with S' and S'' two copies of the sphere S, and $\psi : S \to S'$ the obvious bijection. Consider as in the Schröder-Bernstein theorem the set S_e of points p in S with an even number of ancestors, namely for which there exists an integer $n \geqq 0$ with $p \in \mathrm{Im}(\phi \circ \psi)^n$ and $p \notin \mathrm{Im}(\psi \circ (\phi \circ \psi)^n)$. Consider also the set S_0 of those p in S for which there exists $n \geqq 0$ with $p \in \mathrm{Im}(\psi \circ (\phi \circ \psi)^n)$ and $p \notin \mathrm{Im}(\phi \circ \psi)^{n+1}$, and finally the set S_∞ of those p such that $p \in \mathrm{Im}(\phi \circ \psi)^n$ for any $n \geqq 0$. Consider similarly

$$S' \cup S'' = (S' \cup S'')_e \cup (S' \cup S'')_0 \cup (S' \cup S'')_\infty .$$

Then ψ induces a bijection from $S_e \cup S_\infty$ onto $(S' \cup S'')_0 \cup (S' \cup S'')_\infty$ and ϕ^{-1} from S_0 onto $(S' \cup S'')_e$. Combining these two we have a bijection $\chi : S \to S' \cup S''$ and a partition of S into finitely many pieces, the restriction of χ to each of these being a rotation.